

Norwegian University of Life Sciences

Faculty of Environmental Sciences and Natural Resource Management


2026

ISSN 2535-2806

MINA fagrappport 113

Estimates of wolf density, abundance, and population dynamics in Sweden and Norway, 2016/17 - 2025/26

Pierre Dupont
Cyril Milleret
Asunción Semper-Pascual
Henrik Brøseth
Øystein Flagstad
Jonas Kindberg
Linn Svensson
Richard Bischof



Dupont, P., Milleret, C., Semper-Pascual, A., Brøseth, H., Flagstad, Ø., Kindberg, J., Svensson, L., and Bischof, R., 2026. **Estimates of wolf density, abundance, and population dynamics in Sweden and Norway, 2016/17 - 2025/26** - MINA fagrapport 113. 37 pp.

Ås, June 2026

ISSN: 2535-2806

COPYRIGHT

© Norwegian University of Life Sciences (NMBU)

The publication may be freely cited where the source is acknowledged

AVAILABILITY

Open

PUBLICATION TYPE

Digital document (pdf)

QUALITY CONTROLLED BY

The Research committee (FU), MINA, NMBU

PRINCIPAL

Naturvårdsverket, Ref: NV-366-23-005, Contact person: Robert Ekblom

Miljødirektoratet, Ref: 23047064 Contact person: Terje Bø

COVER PICTURE

Ridgeline plot of wolf density in 2025/2026.

NØKKEORD

Canis lupus, ulv, tetthet, populasjonsdynamikk, deteksjonssannsynlighet, ikke-invaderende innsamling av genetisk materiale, åpen populasjon romlig fangst-gjenfangst, rovdyrforvaltning

KEY WORDS

Canis lupus, wolf, population density, population dynamics, detection probability, non-invasive genetic sampling, open-population spatial capture-recapture, carnivore management

Pierre Dupont, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Cyril Milleret, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Asunción Semper-Pascual, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Henrik Brøseth, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway.

Øystein Flagstad, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway

Jonas Kindberg, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway.

Linn Svensson, Grimsö Wildlife Research Station, Department of Ecology, Swedish University of Agricultural Sciences, Grimsö 152, SE-739 93 Riddarhyttan, Sweden


Richard Bischof (richard.bischof@nmbu.no), Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Summary

Background The Scandinavian wolf (*Canis lupus*) population is being monitored annually using non-invasive genetic sampling (NGS) and recovery of dead individuals. DNA extracted from faeces, urine, hair, and tissue is used to identify the species, sex, and individual from which each sample originated. These data are compiled in the Scandinavian large carnivore database Rovbase 3.0.

Approach Using the Bayesian open-population spatial capture-recapture (OPSCR) model developed by RovQuant, we estimated annual density and vital rates of the Scandinavian wolf population for ten consecutive monitoring seasons from 2016/2017 to 2025/2026.

Results We generated annual density maps and estimated total and jurisdiction-specific population sizes for wolf from the winter 2016/2017 to 2025/2026 within the main population range. Based on the OPSCR model, the size of the Scandinavian wolf population was likely (95% credible interval) between 402 and 453 individuals in 2025/2026, with 350 to 394 individuals attributed to Sweden and 48 to 65 to Norway. These estimates represent a small increase after three years of continuous decline. In addition to annual density and jurisdiction-specific abundance estimates, we report annual estimates of cause-specific mortality, recruitment, and detection probabilities.




Density map and estimated wolf abundance range in 2025-2026

Sammendrag

Bakgrunn Den skandinaviske ulvebestanden (*Canis lupus*) blir overvåket årlig ved bruk av ikke-invasiv genetisk prøveinnsamling (NGS) og gjenfunn av døde individer. DNA isolert fra skit, urin, hår og vev brukes til å identifisere art, kjønn og individ for hver enkelt prøve. Denne informasjonen samles og ivaretas i den skandinaviske databasen for store rovdyr; Rovbase 3.0.

Tilnærming Ved bruk av en Bayesiansk åpen romlig fangst-gjenfangst populasjonsmodell (OP-SCR), utviklet av RovQuant, estimerte vi årlige tettheter og demografiske rater i den skandinaviske ulvebestanden i ti sesonger fra 2016/2017 til 2025/2026.

Resultater Innenfor hovedutbredelsesområdet til ulvebestanden laget vi årlige kart over tettheten av ulv fra vinteren 2016/2017 til 2025/2026, som viser både total bestandsstørrelse og bestandsstørrelser innenfor ulike administrative enheter. Basert på OPSCR-modellen var den skandinaviske ulvebestanden på mellom 402 og 453 individer i 2025/2026 (95% kredibelt intervall), med 350 til 394 individer i Sverige og 48 til 65 individer i Norge. Disse estimatene representerer en liten økning etter tre år med kontinuerlig nedgang. I tillegg til årlige tettheter og områdespesifikke bestandsestimater, gir rapporten estimater på dødelighetsfaktorer, rekruttering og deteksjonssannsynlighet.



Kart som viser tetthet av ulven med intervall for estimert antall ulven i 2025-2026

Contents

1	Introduction	6
2	Methods	8
2.1	Data	8
2.2	Open-population spatial capture-recapture model	9
3	Results	13
3.1	Non-invasive genetic samples and dead recoveries	13
3.2	Density and abundance	13
3.3	Vital rates	17
3.4	Detection probability	18
4	Discussion	19
5	Summary of improvements made	19
6	Suggestions for future improvements	19
7	Acknowledgements	20
8	Data availability	20
	References	22
	Appendices	24

1 Introduction

Non-invasive genetic sampling (NGS) and dead recoveries are central components of national and regional large carnivore monitoring in Norway and Sweden. Both countries have accumulated an extensive individual-based data set for wolf (*Canis lupus*) and plan to continue such transnational monitoring in the future.

Since 2017, project RovQuant has been developing statistical methods that allow a comprehensive assessment of the status and dynamics of large carnivore populations using NGS data and other sources of information collected and stored in Rovbase (www.rovbase.se, www.rovbase.no) by the national monitoring programs in Sweden and Norway. At the core of the analytical framework developed by RovQuant (Bischof et al., 2019b, 2020) are Bayesian open-population spatial capture-recapture (OPSCR) models (Ergon and Gardner, 2014; Bischof et al., 2016; Chandler et al., 2018). These models use the spatial and temporal information contained in the repeated genetic detections of individuals to estimate various population parameters, including spatially-explicit abundance and vital rates. Importantly, the approach accounts for imperfect detection (i.e., the fact that some individuals are not detected at all) and animal movement (i.e., the fact that individuals may use and be detected in multiple management units or countries).

The conventional approach for wolf population size estimation in Scandinavia since 2013/2014 entails the use of a static conversion factor to calculate the size of the wolf population from the number of observed reproductions (Wabakken et al., 2024). The conversion factor approach assumes that the relationship between total population size and the number of observed reproductions remains unchanged over time and between regions, which has led to criticism of the method (Chapron et al., 2016; Bischof et al., 2019a). Aside from relaxing this and several other assumptions (Bischof et al., 2019a), the OPSCR method brings along several advantages. These include the ability to map density, derive jurisdiction-specific abundance, estimate survival and recruitment (which are needed for making population projections), and yield tractable measures of uncertainty (Bischof et al., 2019a, 2020).

RovQuant reported its first results for wolves in 2019 (Bischof et al., 2019a,b, 2020) and on an annual basis since 2021 (Milleret et al., 2021, 2022b, 2023b, 2024, 2025b). In the present report we summarize the analysis of a 10-year time series (2016/2017–2025/2026) using the latest available wolf monitoring data and the most recent version of the OPSCR model. We provide the following information:

- Annual estimates of the number of wolves within the main population range (with Bayesian credible intervals) for Sweden, Norway, and both countries combined, as well as estimates by county and by large carnivore management region in both countries.
- Annual maps of wolf density throughout the species’ primary range in Scandinavia.
- Annual estimates of survival, cause-specific mortality, recruitment, and population growth rate (with Bayesian credible intervals).
- Estimated proportion of individuals detected through non-invasive genetic sampling (with Bayesian credible intervals).

Box 1: Terms and acronyms used

AC: Activity center. Model-based equivalent to the center of an individual's home range during the monitoring period. "AC location" refers to the spatial coordinates of an individual AC in a given year, and "AC movement" to the movement of an individual AC between consecutive years.

CrI: 95% Bayesian credible interval associated with a posterior sample distribution.

Detectors: Potential detection locations in the spatial capture-recapture framework. These can refer to fixed locations (e.g., camera-trap locations) or to areas searched (e.g., habitat grid cells where searches for genetic samples were conducted; used in this project).

Habitat buffer: Buffer surrounding the searched area that is considered suitable habitat but was not searched.

Legal culling: Lethal removal of individuals by legal means including licensed recreational hunting, management removals, defense of life and property.

Länsstyrelserna: Swedish County Administrative Boards, in charge of the monitoring of large carnivores at the county level.

MCMC: Markov Chain Monte Carlo.

NGS: Non-invasive genetic sampling.

OPSCR: Open-population spatial capture-recapture.

p0: Baseline detection probability; probability of detecting an individual at a given detector, if the individual's AC is located exactly at the detector location.

RovQuant: Research group at the Norwegian University of Life Sciences (Ås, Norway) that develops and applies OPSCR models.

σ : Scale parameter of the detection function; related to the size of the circular home-range.

SCR: Spatial capture-recapture.

SNO: Statens naturoppsyn (Norwegian Nature Inspectorate) is the operative field branch of the Norwegian Environment Directorate (Miljødirektoratet).

Statsforvalteren: Norwegian state's representative in the county, responsible for following up decisions, goals, and guidelines from the legislature and the government.

2 Methods

2.1 Data

We included data from multiple sources, the primary one being the Scandinavian large carnivore database Rovbase 3.0 (rovbase.se and rovbase.no; last extraction: 2026-04-18). This database is used jointly by Norway and Sweden to record detailed information associated with large carnivore monitoring, including, but not limited to, non-invasive genetic sampling (NGS) data, dead recoveries, and GPS search tracks. In the following sections, we describe the various types of data used in the analysis. We used data collected during winters 2016/2017 to 2025/2026.

Non-invasive genetic sampling Swedish and Norwegian management authorities, SNO (Norway) and Länsstyrelserna (Sweden) conduct annual searches for sources of DNA (primarily scats and urine) throughout the Scandinavian wolf range (Svensson et al., 2023). Although samples may be collected throughout the year, the official survey period starting with the 2015/2016 season has been Oct 1 – Mar 31. About one third (30%) of DNA samples originated from opportunistic searches (without associated information on search effort) conducted by hunters or other members of the public. See Liberg et al. (2012) and Bischof et al. (2019a) for further details about the data collection procedure. For individual identification, at least 17 microsatellite markers were amplified with PCR throughout the study period. Samples were amplified four times to account for the occurrence of allelic dropout and false alleles (Taberlet et al., 1996). Consensus genotypes were constructed from the replicated PCR runs using the threshold rule that the same alleles had to appear at least twice for a heterozygous genotype and three times for a homozygous genotype (Åkesson et al., 2016). Since 2017/2018, 90 single nucleotide polymorphisms (SNPs) were used for individual identification, in addition to microsatellites. Samples were amplified and genotypes visualized twice using nanofluid arrays (Fluidigm Inc.). Consensus SNP genotypes consisted of the markers with the same allele composition from the two amplifications (Åkesson et al., 2018).

Dead recoveries In Scandinavia, all dead large carnivores killed legally (e.g., legal hunting, management kills, defense of life and property) have to be reported to the state authorities (Statsforvalteren or SNO in Norway and Länsstyrelserna or the police in Sweden). All wolves found dead due to other reasons (e.g., natural deaths, vehicle and train collisions, illegal hunting) also have to be reported, but an unknown proportion remains undetected. Tissue is collected from all reported dead carnivores for DNA extraction and analysis. DNA from dead carnivores, if genotyped, can be linked with NGS data via individual IDs and provide definite information about the fate of individuals. Dead recoveries associated with GPS-coordinates also provide additional information for the estimation of individual locations, especially valuable for individuals with few or no NGS detections (Dupont et al., 2021).

GPS search tracks Government employees involved in systematic searches for wolf DNA along roads and following wolf tracks (e.g., via snowmobiles, skis, snowshoes) documented their effort with GPS track logs which were registered in Rovbase 3.0. GPS search tracks were included in the OPSCR model to account for spatial and temporal variation in search effort.

Observation reports in Skandobs We used all observation records available from Skandobs (skandobs.se; skandobs.no) that were recorded during the wolf monitoring seasons since Oct 1, 2016, (last extraction: 2026-04-18). Skandobs is a web application that allows anyone to anonymously register observations (e.g., visual, tracks, faeces) of bears, lynx, wolves, and wolverines in Scandinavia. This database consists of around 120 000 records of possible large carnivore observations. Although most observations are not verified, they offer the best available proxy for spatio-temporal variation in opportunistic search effort.

Parentage analysis and social status assignation Each year, field observations, genetic sampling data, dead recoveries, and parentage analysis (which resulted in a near-complete pedigree of the Scandinavian wolf population) are used to establish counts of pairs, packs, and reproductions (Svensson et al., 2025; Wabakken et al., 2026). The social status (adult scent-marking member in a pack/pair, or subadult member of a pack and vagrant individuals) was determined on a yearly basis for genetically identified individuals using the pedigree and tracking information from the monitoring (Åkesson et al., 2022). The identity of known adult scent-marking individuals was reported in the yearly monitoring reports (Svensson et al., 2025; Wabakken et al., 2026) and a detailed description of the method used for parental assignment was provided in Åkesson et al. (2016). We assigned individuals to two different states depending on whether the individual had been identified as an adult scent-marking member of a pair or a pack (“adult scent-marking”) or not (“other”). See Bischof et al. (2019a) for further details.

Study area Although wolves can occur throughout the Scandinavian peninsula, we restricted the analysis to the core area (Figure 1). Following the recent wolf range expansion towards the south of Sweden, the Swedish Environmental Protection Agency requested a corresponding expansion of the study area to include southern Sweden (Milleret et al., 2022b, 2023b). Since 2022, the RovQuant models therefore include an additional 68 400 km² in the current study area (total = 322 400 km²).

2.2 Open-population spatial capture-recapture model

We analysed the data using a Bayesian open-population spatial capture-recapture (OPSCR) model (Bischof et al., 2019b), which addresses three challenges associated with population-level wildlife inventories:

1. Detection is imperfect and sampling effort heterogeneous in space and time: not all individuals present in the study area are detected (Kéry and Schaub, 2012).
2. Individuals that reside primarily outside the surveyed area may be detected within it. Without an explicit link between the population size parameter and geographic space or area, density cannot be estimated and population size is ill-defined (Efford, 2004).
3. Non-spatial population dynamic models usually estimate “apparent” survival and recruitment, as these parameters include the probability of permanent emigration and immigration, respectively. By explicitly modelling movement of individuals between years, the OPSCR model can help return unbiased estimates of demographic parameters (Ergon and Gardner, 2014; Schaub and Royle, 2014; Gardner et al., 2018; Efford and Schofield, 2022).

The OPSCR model (Ergon and Gardner, 2014; Bischof et al., 2016; Chandler et al., 2018) is composed of three sub-models:

1. A model for population dynamics and population size.
2. A model for density and movements.
3. A model for detections during DNA searches.

Population dynamics and population size sub-model We used a multi-state formulation (Lebreton and Pradel, 2002) where each individual life history is represented by a succession of up to five discrete states $z_{i,t}$: 1) “unborn” if individual i has not yet been recruited in the population (state “unborn” is required for the data augmentation procedure, see below) by the start of monitoring season t ; 2) “alive other” if it is alive and not assigned as an “adult scent-marking” individual; 3) “alive adult scent-marking” if it is alive and assigned as an “adult scent-marking” individual; 4) “culled” if it was culled and therefore recovered dead between the start of the previous and current monitoring season; 5) “Dead from other causes and recovered” if it died from

other causes of mortality (e.g. disease, car collision, ...) and was reported to the authorities; and 6) "dead": if it has died but was not recovered. We then modelled the transition between states from one monitoring season to the next (t to $t + 1$) to estimate vital rates (recruitment and mortality). More details are available in Bischof et al. (2019b), and Bischof et al. (2020). We used data augmentation (Royle and Dorazio, 2012), whereby additional, undetected individuals are available for inclusion in the population at each time step.

Density and movement sub-model We used a Bernoulli point process to model the distribution of individual activity centers. In the first year, individuals were located according to an intensity surface, which was a function of the locations of known packs at time $t - 1$ (see Bischof et al., 2019b and Bischof et al., 2020 for more details). For all subsequent years ($t > 1$), the location of individual activity centers was a function of the distance from previous activity centers (at time $t - 1$), as well as a function of the locations of known packs (at time $t - 1$). As in previous analyses (Milleret et al., 2022b, 2023b, 2024, 2025b), we used an exponential model to describe the movement of individuals between years, as it better accommodates distributions with long tails (i.e., a few individuals that make exceptionally long dispersal movements).

Detection sub-model SCR models take into account the spatial-variation in individual detection probability based on the distance between activity center locations (estimated by the density sub-model) and a given detector. A half-normal function was used to express the declining probability of detection with increasing distance between the activity center and the detector (Efford, 2011).

In Scandinavia, DNA material from live wolves is collected following two main processes. First, authorities collect genetic samples and record the corresponding search effort during official searches ("structured sampling" thereafter). Second, DNA material can be collected by any member of the public (e.g., hunters) or by the authorities in a more or less opportunistic manner and associated search effort is not directly available ("unstructured" sampling thereafter). Currently, it is not possible to unambiguously distinguish between samples collected by the authorities during the structured or unstructured sampling in Rovbase. We therefore assigned each sample to structured or unstructured sampling based on whether a given sample matched in time and space with recorded search tracks: a sample was assigned to the "structured" sampling if it was collected by the authorities (marked as collected by "Statsforvalteren" or "SNO" in Rovbase) and if it was located within 500 m from a search track recorded on the same day. All remaining samples were assigned to the unstructured sampling.

We assumed that both sampling processes could in theory occur within the entire study area and therefore used the same 10 km detector grid for both observation processes. Samples were then assigned to the closest detector (see details in Bischof et al., 2019b, and Bischof et al., 2020). However, spatial and temporal variation in the probability to detect a sample during the structured or unstructured sampling were assumed to be driven by different processes.

We accounted for spatio-temporal heterogeneity in detectability during *structured sampling* using:

- Spatial-temporal variation in effort using the length of GPS search tracks in each detector grid cell.
- Spatio-temporal variation in snow cover.
- Spatio-temporal variation in monitoring regimes between jurisdictions (groups of counties in Sweden, monitoring regions in Norway).

- Individual variation linked with the state of the individual (i.e., adult scent-marking individuals, other individuals).
- Individual variation linked with a previous detection that could be expected to positively or negatively influence the probability of being detected at subsequent occasions, depending on the state of knowledge in the territory or neighboring territories.

We accounted for spatio-temporal heterogeneity in detectability during *unstructured sampling* using:

- Spatio-temporal variation in unstructured sampling or carnivore observations (Figure A.1). For each detector grid cell and during each monitoring season (Oct 1 – Mar 31), we identified whether a) any carnivore sample had been registered in Rovbase (excluding successfully genotyped wolf samples already used in the OPSCR analysis) or b) any observation of carnivores had been registered in Skandobs. Roughly, the variable distinguishes areas with very low detection probability from those with higher probability that carnivore DNA samples, if present in a detector grid cell, could have been detected and submitted for genetic analysis.
- Spatio-temporal variation in snow cover.
- Spatial variation in accessibility measured as the average distance to the nearest road.
- Spatio-temporal variation between countries.
- Individual and temporal variation linked with the state of the individual (i.e., adult scent-marking individuals, other individuals).
- Individual and temporal variation linked with a previous detection that could be expected to positively influence the probability of being detected at subsequent occasions.

The different model components and data sources for covariates are described in detail in Bischof et al. (2019a), Bischof et al. (2019b), and Bischof et al. (2020).

Model fitting We fitted sex-specific Bayesian OPSCR models using Markov chain Monte Carlo (MCMC) simulation with NIMBLE version 0.12.2 (Turek et al., 2021; de Valpine et al., 2017; de Valpine et al., 2022) and nimbleSCR (Bischof et al., 2021) in R version 4.1.0 (R Core Team, 2021). We ran 4 chains each with 25 000 iterations, including a 5 000-iterations burn-in period. Due to the computing challenge associated with post-processing large amounts of data, we thinned chains by a factor of 2 from which abundance estimates were derived. We considered models as converged when the Gelman-Rubin diagnostics (Rhat, Gelman and Rubin, 1992) was ≤ 1.1 for all parameters and by visually inspecting the trace plots.

Abundance estimates To obtain an estimate of abundance for any given area, we summed the number of predicted AC locations (live individuals) that fell within that area for each iteration of the MCMC chains, thus generating a posterior distribution of abundance for that area. In this fashion, abundance estimates and the associated uncertainty can be extracted for any spatial unit, including countries or counties (Figure A.2). Individuals detected near a border can have their model-predicted AC placed on different sides of that border in different model iterations (even if detections are only made on one side of the border). As a result, the probability of designating such individuals to either side of the border can be integrated into jurisdiction-specific abundance estimates. This is especially relevant for wolves detected along the Swedish and Norwegian border ("cross-boundary wolves", Wabakken et al., 2022); individual wolves can be partially designated to both countries (Bischof et al., 2016).

To ensure that abundance estimates for spatial sub-units (jurisdictions) add up to the overall abundance estimate, we used the mean and associated 95% Bayesian credible interval limits to summarize posterior distributions of abundance. Combined (female/male) parameter estimates were obtained by merging posterior samples from the sex-specific models.

Density maps We used both the distribution of model-estimated AC positions and the scale parameter (σ) of the detection function to construct density maps based on individual utilization distributions. These maps are not only based on the position of the center of an individual’s home range, but also take into account the area over which that individual’s activity is spread, i.e., its space use (Bischof et al., 2020). To do so, we constructed raster maps (5 km resolution) of individual utilization distributions, scaled values in each raster to sum to one, and then summed rasters across individuals to create a single population-level raster map for each iteration. An overall density map was derived by calculating the mean across iterations in each cell (Bischof et al., 2020).

Other derived parameters The average proportion of individuals detected and the associated uncertainty (Table A.12) were obtained by dividing the number of individuals detected through NGS sampling by the abundance estimates and their associated Bayesian credible intervals, respectively.

We derived the proportion of females in the population and the associated uncertainty by dividing the posterior distribution representing the number of females by the combined abundance estimate for males and females. Yearly population growth rates were calculated as $\lambda_t = N_{t+1}/N_t$ (Table A.7).

Focus on uncertainty Although we reported median (or mean for abundance; see above) estimates for all parameters in the tables, we intentionally focused the main results of this report on the 95% Bayesian credible interval limits of the estimates. We did so with the aim of drawing the reader’s attention to the uncertainty around population size estimates, rather than on a single point estimate (Milleret et al., 2022a).

3 Results

3.1 Non-invasive genetic samples and dead recoveries

A total of 21 962 (9 622 female; 12 340 male) genotyped wolf genetic samples were included in the analysis, of which 77% originated from Sweden. These samples were associated with 1 948 (908 female; 1040 male) individuals. We did not include individuals with unknown sex in this analysis. Among all genotyped samples, 10 065 (4196 female; 5869 male) were assigned to the structured sampling and 11 897 (5426 female; 6471 male) to the unstructured sampling. We also used 680 (297 female; 383 male) dead recoveries of wolves in the OPSCR model, of which 573 (251 female; 322 male) were due to legal culling and 107 (46 female; 61 male) due to other causes of mortality. The majority of dead recoveries (66%) originated from Sweden. Annual total and country-specific tallies of NGS samples and associated individuals, as well as dead recoveries included in the analysis are provided in the Appendices (NGS samples: Table A.1 and Table A.2, NGS individuals: Table A.3, dead recoveries: Table A.4).

3.2 Density and abundance

Wolf abundance for the entire study area (322 400 km², excluding the buffer area) was likely (95% Bayesian credible interval) between 402 and 453 individuals in 2025/2026 (Table 1, Figure 1). Based on the model-predicted location of ACs, we estimated that in 2025/2026, between 350 to 394 individuals could be attributed to Sweden and 48 to 65 to Norway (Table 1). Estimates refer to the status of the population at the start of the annual sampling period (Oct. 1). The proportion of females in the Scandinavian wolf population was likely between 48% and 54 % in 2025/2026. See Table 1 for total and sex-specific abundance estimates for each country and management area. See Table A.5 and Table A.6 for annual wolf abundance estimates for all of Scandinavia, by region and by county between 2016/2017 and 2025/2026. In the last year, the wolf population has experienced a rebound after reaching 386-416 individuals in 2024/2025 and following a steady decrease from 496-529 individuals in 2021/2022 (Figure 2, Table A.5). Note that estimates for different years (Figure 2) shown here differ slightly from those provided in our previous report (Milleret et al., 2023b). This is due to the inclusion of additional years of data. The analysis yielded annual density maps, which illustrate changes in the distribution of wolves over time (Figure A.3).

Table 1: Wolf population size estimates in 2025/2026 within the study area by sex at several spatial scales: the entire study area, by country, by management unit (carnivore management regions in Norway and "Rovdjursförvaltningsområden" in Sweden), and counties ("Län" in Sweden); see also Figure A.2. Only counties and management units that are within or that intersect the study area are included in the table. Readers should focus on the 95% Bayesian credible interval provided in parentheses, as these - unlike mean values - convey the uncertainty inherent in abundance estimates. Numbers are based on estimated activity center locations of wolves. Combined female-male estimates were obtained by joining sex-specific posterior distributions. Rounding may result in small deviations between total estimates and the sum of the estimates for constituent regions. Note that the numbers reported here are predictions from a statistical model which always represents an oversimplification of reality and is based on available data (NGS and dead recoveries). As a consequence, especially at the local scale, the model-estimated number of wolves based on DNA sampling can deviate from the number of wolves inferred from ancillary observations (e.g., camera traps). Estimates for Norwegian counties are provided in Table A.6.

	Females	Males	Total
TOTAL	215.7 (198-238)	209.6 (196-225)	425.3 (402-453)
NORWAY	32.6 (26-41)	22.7 (20-27)	55.2 (48-65)
Region 2	0.3 (0-2)	0.3 (0-2)	0.6 (0-2)
Region 3	1.1 (0-4)	0.8 (0-3)	2 (0-5)
Region 4	8.8 (6-12)	7.8 (6-10)	16.6 (13-21)
Region 5	21 (16-27)	13.8 (11-17)	34.8 (29-41)
Region 6	1.3 (0-4)	0 (0-1)	1.4 (0-4)
SWEDEN	183.1 (168-202)	187 (174-202)	370.1 (350-394)
Norra	9.7 (4-17)	6.1 (3-11)	15.8 (9-24)
Jämtland	5.4 (1-11)	3.4 (1-7)	8.7 (3-15)
Västerbotten	0 (0-1)	0 (0-1)	0.1 (0-1)
Västernorrland	4.3 (2-8)	2.7 (1-6)	7 (4-12)
Mellersta	140.2 (128-155)	139.7 (129-152)	279.8 (263-298)
Dalarna	22.1 (15-30)	20.1 (14-27)	42.2 (33-53)
Gävleborg	11.5 (8-16)	13.7 (10-18)	25.2 (20-32)
Örebro	25.3 (21-30)	34.3 (30-38)	59.6 (54-66)
Stockholm	2 (1-4)	2.2 (1-4)	4.2 (2-7)
Uppsala	8.6 (6-11)	9.4 (7-12)	18 (15-22)
Värmland	26.1 (20-34)	17.5 (12-24)	43.6 (35-54)
Västmanland	16.4 (13-20)	20 (16-24)	36.5 (31-42)
Västra Götaland	28 (25-32)	22.4 (19-27)	50.4 (45-56)
Södra	33.3 (30-38)	41.2 (37-47)	74.5 (69-82)
Blekinge	0.1 (0-1)	0.2 (0-1)	0.3 (0-2)
Halland	1.1 (0-3)	1.2 (0-3)	2.3 (1-5)
Jönköping	8.5 (7-10)	7.1 (5-10)	15.5 (13-19)
Kalmar	1.3 (1-3)	0.5 (0-2)	1.9 (1-4)
Kronoberg	1.1 (0-3)	3.2 (1-6)	4.3 (1-8)
Östergötland	3.5 (2-6)	4.3 (3-7)	7.9 (5-11)
Skåne	5.7 (5-7)	8.6 (6-11)	14.2 (11-17)
Södermanland	12.1 (10-14)	16 (15-18)	28.1 (26-31)




Figure 1: Wolf density based on individual utilization distributions throughout the study area (white background) in Scandinavia in 2025/2026. This map is available as a geo-referenced raster file at <https://github.com/richbi/RovQuantPublic>




Figure 2: Annual wolf population size estimates within the study area by country (blue: Sweden, red: Norway), and total (black) between monitoring seasons 2016/2017 and 2025/2026. Darker and lighter bars show the 50% and 95% Bayesian credible intervals, respectively. Bayesian credible intervals indicate uncertainty in estimates given the model and data used to generate the estimates. Light grey bars in the background show the total number of individuals detected within the entire study area. Changes in the model and the data can result in different estimates and associated uncertainty compared with estimates provided in previous reports by RovQuant.

3.3 Vital rates

The OPSCR model produced annual estimates of legal hunting mortality, mortality associated with all other causes, and per capita recruitment rates (Figure 3, Table A.8, Figure 4). There was temporal variation in mortality rates, with the risk of mortality from causes other than culling generally higher than the risk of mortality from culling. Overall, scent-marking adult individuals had a lower mortality compared to other individuals, with no pronounced systematic difference between years or sexes (Figure A.4). Similar to the abundance estimates, recruitment experienced a slight rebound in 2025/26 after a steady decline since 2020/21 (Figure 4).




Figure 3: Mortality probabilities due to legal culling (light green) and all other causes (dark green) for female and male wolves. Darker and lighter bars show the 50% and 95% Bayesian credible intervals, respectively. Shown are overall estimates throughout the study area. Estimates refer to deaths occurring between the start of one sampling season and the start of the next.




Figure 4: Estimated annual number of recruits of female and male wolves. Recruitment represents the number of new individuals present in the population on Oct 1 (i.e., individuals that were born between the two consecutive monitoring seasons and survived to Oct 1 or that immigrated in the study area). Darker and lighter bars show the 50% and 95% Bayesian credible intervals, respectively.

3.4 Detection probability

The overall proportion of detected individuals in the study area was likely between 78% and 88% in 2025/2026 (Table A.12). The baseline detection probability during structured and unstructured sampling varied both in time and space (Figure A.5 and Figure A.6). More specifically, the length of recorded search tracks positively affected detection probability during structured sampling (2025/2026; males: $\beta_{2Structured} = 0.42$, CrI: 0.35 - 0.50; females: $\beta_{2Structured} = 0.44$, CrI: 0.37 - 0.51; Table A.10). Previous detection of individuals only had a significant effect on females ($\beta_{1Structured} = 0.43$, CrI: 0.12 - 0.74), while the average proportion of snow cover had no strong effect on detection probability for the structured sampling of both sexes (Table A.10). Unstructured search effort derived using the observation data in Skandobs and Rovbase had a strong positive effect on detection probability during unstructured sampling (2025/2026; males: $\beta_{4Unstructured} = 2.22$, CrI: 1.60 - 3.03; females: $\beta_{4Unstructured} = 2.46$, CrI: 1.82 - 3.18; Table A.11). The effect of previous detection also tended to have a positive effect on detection probability during unstructured sampling, but the pattern was not consistent across years (Table A.11).

4 Discussion

The size of the wolf population on the Scandinavian peninsula steadily declined from 496-529 individuals in 2021/22 to 386-416 individuals in 2024/25, but the latest analysis indicates a stabilization or slight increase relative to the previous monitoring season (Figure 2). The decline during the preceding years was pronounced both in Sweden, where population size decreased from 399-432 to 337-365 individuals between 2021/22 and 2024/25, and in Norway, where population size decreased from 87-107 individuals to 42-56 individuals during the same period (Figure 2, Table A.5). The reduction in the number of individuals during that period can at least in part be attributed to reduced recruitment of both females and males (Figure 4). The recent stabilization is consistent with an increase in recruitment (Figure 4), as we do not observe any corresponding reduction in overall mortality that could otherwise explain the change in trend (Figure 3).

From 2021/22 to 2024/25, the number of recorded packs dropped from 55 to 40 (Wabakken et al., 2022; Svensson et al., 2025). Because packs are the reproductive units of the wolf population, this reduction likely underlies the earlier decrease in recruitment (see also Chapron et al. (2016)). By contrast, the recent increase in recruitment may reflect improved reproductive output within the remaining packs rather than a recovery in pack numbers, given that the recorded numbers of packs (41), pairs (30) and documented reproductions (39) did not significantly change in the last year (Wabakken et al., 2026).

Currently, vital rates are estimated with the assumption that they are homogeneous across the population range. This assumption is likely to be violated and future analysis should consider spatially variable vital rates (Milleret et al., 2023a, 2025a, 2026), which may further help shed light on the drivers of the observed population trends.

5 Summary of improvements made

The analysis described in this report includes the following adjustments compared with previous analyses of wolf density in Scandinavia by RovQuant (Milleret et al., 2025b):

1. Shifted the time series of data analyzed by one year (2016/17 to 2025/26 instead of 2015/16 to 2024/25).
2. Fixed a temporal mismatch between the DNA data and the covariate used to account for spatio-temporal variation in unstructured sampling.

6 Suggestions for future improvements

RovQuant continues to work on improving the functionality and efficiency of OPSCR models. We recommend the following developments for future analyses of the Scandinavian wolf monitoring data:

1. Review and adjust spatial covariates on density. This may involve the addition of land cover and topographic variables.
2. Account for individual heterogeneity in detectability, for example by using a finite-mixture approach (Cubaynes et al., 2010).
3. Consider alternative detection models that do not assume a half-normal shape and/or circular home ranges (Sutherland et al., 2015).
4. Account for spatial variation in vital rates (i.e., survival, recruitment; Milleret et al. 2023a).

7 Acknowledgements

This work was made possible by the large carnivore monitoring programs and the extensive monitoring and observation data collected by Swedish (Länstyrelsen) and Norwegian (SNO) wildlife management authorities, as well as the public in both countries. Our analysis relied on genetic analyses conducted by the laboratory personnel at the DNA laboratories at the Swedish University of Agricultural Sciences, and the Norwegian Institute for Nature Research. We also thank Swedish and Norwegian wildlife managers for feedback provided during project RovQuant and the Research Council of Norway for partial funding (NFR 345279: project PopFlow). The computations/simulations were performed on resources provided by NMBU’s computing cluster “Orion”. J. Vermaat provided helpful comments on a draft of this report.

8 Data availability

Data, R code to reproduce the analysis, as well as figures, tables, and raster maps (Figure A.3) are available at <https://github.com/richbi/RovQuantPublic>.

References

- Åkesson, M., Danielsson, A., Hedmark, E., and Öhrn (2018). Teknisk rapport över genetiska analyser på varg i sverige år 2017. *Rapport från SLU*, 162 s.
- Åkesson, M., Liberg, O., Sand, H., Wabakken, P., Bensch, S., and Flagstad, O. (2016). Genetic rescue in a severely inbred wolf population. *Molecular Ecology*, 25(19):4745–4756.
- Bischof, R., Brøseth, H., and Gimenez, O. (2016). Wildlife in a politically divided world: Insularism inflates estimates of brown bear abundance. *Conservation Letters*, 9(2):122–130.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Åkesson, M., Brøseth, H., and Kindberg, J. (2019a). Estimating the size of the scandinavian wolf population with spatial capture recapture and conversion factors. *MINA fagrapport*, 57:80pp.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Brøseth, H., and Kindberg, J. (2019b). Estimating density, abundance and population dynamics of bears, wolverines, and wolves in scandinavia. *MINA fagrapport*, 63:79pp.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Tourani, M., Ordiz, A., de Valpine, P., Turek, D., Royle, J. A., Gimenez, O., Flagstad, Ø., Åkesson, M., Svensson, L., Brøseth, H., and Kindberg, J. (2020). Estimating and forecasting spatial population dynamics of apex predators using transnational genetic monitoring. *Proceedings of the National Academy of Sciences*, 117(48):30531–30538.
- Bischof, R., Turek, D., Milleret, C., Ergon, T., Dupont, P., Dey, S., and de Valpine, P. (2021). *nimbleSCR: Spatial Capture-Recapture (SCR) Methods Using 'nimble'*. R package version 0.1.2.
- Chandler, R. B., Hepinstall-Cymerman, J., Merker, S., Abernathy-Conners, H., and Cooper, R. J. (2018). Characterizing spatio-temporal variation in survival and recruitment with integrated population models. *The Auk*, 135(3):409–426.
- Chapron, G., Wikenros, C., Liberg, O., Wabakken, P., Flagstad, O., Milleret, C., Månsson, J., Svensson, L., Zimmermann, B., Åkesson, M., and Sand, H. (2016). Estimating wolf (*canis lupus*) population size from number of packs and an individual based model. *Ecological Modelling*, 339:33 – 44.
- Cubaynes, S., Pradel, R., Choquet, R., Duchamp, C., Gaillard, J. M., Lebreton, J. D., Marboutin, E., Miquel, C., Reboulet, A. M., Poillot, C., Taberlet, P., and Gimenez, O. (2010). Importance of accounting for detection heterogeneity when estimating abundance: the case of French wolves. *Conservation Biology*, 24(2):621–626.
- de Valpine, P., Paciorek, C., Turek, D., Michaud, N., Anderson-Bergman, C., Obermeyer, F., Wehrhahn Cortes, C., Rodríguez, A., Temple Lang, D., and Paganin, S. (2022). *NIMBLE User Manual*. R package manual version 0.12.2.
- de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., and Bodik, R. (2017). Programming with models: writing statistical algorithms for general model structures with nimble. *Journal of Computational and Graphical Statistics*, 26(2):403–413.
- Dupont, P., Milleret, C., Tourani, M., Brøseth, H., and Bischof, R. (2021). Integrating dead recoveries in open-population spatial capture–recapture models. *Ecosphere*, 12(7):e03571.
- Efford, M. (2004). Density estimation in live-trapping studies. *Oikos*, 106(3):598–610.
- Efford, M. G. (2011). Estimation of population density by spatially explicit capture-recapture analysis of data from area searches. *Ecology*, 92(12):2202–2207.
- Efford, M. G. and Schofield, M. R. (2022). A review of movement models in open population capture–recapture. *Methods in Ecology and Evolution*, 13(10):2106–2118.
- Ergon, T. and Gardner, B. (2014). Separating mortality and emigration: modelling space use, dispersal and survival with robust-design spatial capture–recapture data. *Methods in Ecology and Evolution*, 5(12):1327–1336.
- Gardner, B., Sollmann, R., Kumar, N. S., Jathanna, D., and Karanth, K. U. (2018). State space and movement specification in open population spatial capture–recapture models. *Ecology and Evolution*, 8(20):10336–10344.
- Gelman, A. and Rubin, D. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7:457–511. <http://www.stat.columbia.edu/~gelman/research/published/itsim.pdf>.
- Kéry, M. and Schaub, M. (2012). *Bayesian population analysis using WinBUGS: a hierarchical perspective*. Academic Press, Waltham, MA.
- Lebreton, J.-D. and Pradel, R. (2002). Multistate recapture models: modelling incomplete individual histories. *Journal of Applied Statistics*, 29:353–369.
- Liberg, O., Aronson, Å., Sand, H., Wabakken, P., Maartmann, E., Svensson, L., and Åkesson, M. (2012). Monitoring of wolves in scandinavia. *Hystrix, the Italian Journal of Mammalogy*, 23(1):29–34.
- Milleret, C., Dey, S., Dupont, P., Brøseth, H., Turek, D., de Valpine, P., and Bischof, R. (2023a). Estimating spatially variable and density-dependent survival using open-population spatial capture–recapture models. *Ecology*, 104(2):e3934.
- Milleret, C., Dupont, P., and Bischof, R. (2026). Spatial variation in wolverine mortality, 2019–2024. *MINA fagrapport*, 109:22pp.
- Milleret, C., Dupont, P., Brøseth, H., Flagstad, O., Kindberg, J., and Bischof, R. (2022a). Estimates of wolverine

- density, abundance, and population dynamics in scandinavia, 2013–2021. *MINA fagrapport*, 74:30pp.
- Milleret, C., Dupont, P., Brøseth, H., Øystein, F., Kindberg, Jonas Svensson, L., , and Bischof, R. (2023b). Estimates of wolf density, abundance, and population dynamics in Scandinavia, 2013–2023. *MINA fagrapport*, 85:34pp.
- Milleret, C., Dupont, P., Brøseth, H., Øystein, F., Kindberg, Jonas Svensson, L., , and Bischof, R. (2024). Estimates of wolf density, abundance, and population dynamics in Scandinavia, 2014–2024. *MINA fagrapport*, 97:34pp.
- Milleret, C., Dupont, P., Dey, S., Brøseth, H., Kindberg, J., Turek, D., de Valpine, P., Åkesson, M., Wabakken, P., Zimmermann, B., et al. (2025a). Map of death: spatially explicit mortality of the grey wolf. *Proceedings of the Royal Society B: Biological Sciences*, 292(2053).
- Milleret, C., Dupont, P., Semper-Pascual, A., Brøseth, H., Flagstad, Ø., Kindberg, J., Svensson, L., and Bischof, R. (2025b). Estimates of wolf density, abundance, and population dynamics in sweden and norway, 2015–2025. *MINA fagrapport*, 103:36pp.
- Milleret, C., Dupont, P., Åkesson, M., Brøseth, H., Kindberg, J., and Bischof, R. (2021). Estimates of wolf density, abundance, and population dynamics in scandinavia, 2012 - 2021. *MINA fagrapport*, 72:30pp.
- Milleret, C., Dupont, P., Åkesson, M., Brøseth, H., Kindberg, J., and Bischof, R. (2022b). Estimates of wolf density, abundance, and population dynamics in scandinavia, 2013-2022. *MINA fagrapport*, 77:35pp.
- R Core Team (2021). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Royle, J. A. and Dorazio, R. M. (2012). Parameter-expanded data augmentation for bayesian analysis of capture–recapture models. *Journal of Ornithology*, 152(2):521–537.
- Schaub, M. and Royle, J. A. (2014). Estimating true instead of apparent survival using spatial Cormack–Jolly–Seber models. *Methods in Ecology and Evolution*, 5(12):1316–1326.
- Sutherland, C., Fuller, A. K., and Royle, J. A. (2015). Modelling non-Euclidean movement and landscape connectivity in highly structured ecological networks. *Methods in Ecology and Evolution*, 6(2):169–177.
- Svensson, L., Wabakken, P., Maartmann, E., Nordli, K., Flagstad, Ø., Danielsson, A., Hensel, H., Pöchlacker, K., and Åkesson, M. (2023). Inventering av varg vintern 2022-2023. bestandsövervakning av ulv vintern 2022-2023. *Bestandsstatus for store rovdyr i Skandinavia. Bestandsstatus för stora rovdjur i Skandinavien 1-2023*, 1(65s).
- Svensson, L., Wabakken, P., Maartmann, E., Nordli, K., Jensen, M. Ø., Milleret, C., Dupont, P., Bischof, R., Åkesson, M., and Flagstad, Ø. (2025). Bestandsövervakning av ulv vintern 2024-2025. inventering av varg vintern 2024-2025. *Bestandsstatus for store rovdyr i Skandinavia. Bestandsstatus för stora rovdjur i Skandinavien 1-2025*, 1(62s).
- Taberlet, P., Griffin, S., Goossens, B., Questiau, S., Manceau, V., Escaravage, N., Waits, L. P., and Bouvet, J. (1996). Reliable genotyping of samples with very low dna quantities using pcr. *Nucleic acids research*, 24(16):3189–3194.
- Turek, D., Milleret, C., Ergon, T., Brøseth, H., Dupont, P., Bischof, R., and de Valpine, P. (2021). Efficient estimation of large-scale spatial capture–recapture models. *Ecosphere*, 12(2):e03385.
- Wabakken, P., Svensson, L., , Maartmann, E., Nordli, K., Flagstad, O., and Åkesson, M. (2022). Bestandsövervakning av ulv vintern 2021-2022. inventering av varg vintern 2021-2022. *Bestandsstatus for store rovdyr i Skandinavia. Bestandsstatus för stora rovdjur i Skandinavien 1-2022*, 1(59s).
- Wabakken, P., Svensson, L., Maartmann, E., Nordli, K., Dupont, P., Milleret, C., Bischof, R., Åkesson, M., and Flagstad, Ø. (2026). Bestandsövervakning av ulv vintern 2025-2026. inventering av varg vintern 2025-2026. *Bestandsstatus for store rovdyr i Skandinavia. Bestandsstatus för stora rovdjur i Skandinavien*, 1(63s).
- Wabakken, P., Svensson, L., Maartmann, E., Nordli, K., Flagstad, Ø., Danielsson, A., Cardoso Palacios, C., and Åkesson, M. (2024). Bestandsövervakning av ulv vintern 2023-2024. inventering av varg vintern 2023-2024. *Bestandsstatus for store rovdyr i Skandinavia. Bestandsstatus för stora rovdjur i Skandinavien 1-2024*, 1(61s).
- Åkesson, M., Svensson, L., Flagstad, Ø., Wabakken, P., and Frank, J. (2022). Wolf monitoring in scandinavia: evaluating counts of packs and reproduction events. *The Journal of Wildlife Management*, 86(4):e22206.

Appendices




Figure A.1: Covariate used to account for spatio-temporal variation in unstructured sampling in the study area. Green cells (10 x 10 km) represent areas with at least one carnivore record in Rovbase (rovbase.no, rovbase.se, excluding the ones used in the OPSCR model) or an observation record from Skandobs (skandobs.se, skandobs.no) during a given monitoring season (Oct 1 – Mar 31).




Figure A.2: Management jurisdictions in Norway and Sweden. Shaded polygons show carnivore management regions in Norway and counties in Sweden. The Norwegian county borders used to extract abundance estimates presented in Table A.6 are also displayed in white. Thick outlines delineate Swedish carnivore management regions ("Rovdjursförvaltningsområden") encompassing multiple counties.




Figure A.3: Wolf density based on individual utilization distributions throughout the study area (white background) in Scandinavia between 2016/2017 and 2025/2026. These maps are available as geo-referenced raster files at <https://github.com/richbi/RovQuantPublic>.




Figure A.4: Annual survival probabilities for scent-marking adults and other individuals and for female and male wolves. Darker and lighter bars show the 50% and 95% Bayesian credible intervals, respectively. Combined mortality probabilities (and therefore overall survival) are obtained by adding the mortality probabilities due to legal culling and all other causes (Figure 3). Shown are overall estimates for the entire study area between 2016/2017 and 2025/2026.




Figure A.5: Sex-specific baseline detection probabilities ($p_{0structured}$) of scent-marking adults and other individuals for the different Scandinavian jurisdictions during structured sampling as estimated by the open-population spatial capture-recapture model. Bars represent 95% Bayesian credible intervals. Results are separated into panels based on regions. Estimates are shown for the mean values of the detection covariates. Note that baseline detection probability (p_0) is a theoretical value of detection probability when a detector coincides with the location of an individual's activity center. It is not to be confused with detectability, i.e, the overall probability of detecting an individual.




Figure A.6: Sex- and country-specific baseline detection probabilities ($p_{0unstructured}$) of scent-marking adults and others individuals during unstructured sampling as estimated by the open-population spatial capture-recapture model. Bars represent 95% Bayesian credible intervals. Estimates are shown for the mean values of the detection covariates. Note that baseline detection probability (p_0) is a theoretical value of detection probability when a detector coincides with the location of an individual's activity center. It is not to be confused with detectability, i.e, the overall probability of detecting an individual.

Table A.1: Annual number of wolf non-invasive genetic samples included in the analysis. Numbers are reported by country, for females (F) and males (M), and for each type of sampling (structured and unstructured). We included only samples collected within the study area during the primary monitoring period (Oct 1 - Mar 31) between 2016/2017 and 2025/2026.

		2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		2021/2022		2022/2023		2023/2024		2024/2025		2025/2026	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
Norway	Structured	20	10	36	42	53	93	44	75	51	130	75	160	56	116	90	155	89	113	135	184
	Unstructured	245	188	232	238	163	224	188	231	260	316	126	223	102	96	100	143	86	89	77	89
Sweden	Structured	373	582	474	665	221	293	597	717	280	456	414	500	361	460	258	375	213	314	356	429
	Unstructured	463	536	603	830	129	162	835	831	256	340	278	385	430	490	257	340	308	359	288	361
Total	Structured	393	592	510	707	274	386	641	792	331	586	489	660	417	576	348	530	302	427	491	613
	Unstructured	708	724	835	1068	292	386	1023	1062	516	656	404	608	532	586	357	483	394	448	365	450

Table A.2: Annual number of wolf non-invasive genetic samples included in the analysis. Numbers are reported by country, for females (F) and males (M). We included only samples collected within the study area during the primary monitoring period (Oct 1 - March 31) between 2016/2017 and 2025/2026.

		2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		2021/2022		2022/2023		2023/2024		2024/2025		2025/2026	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
Norway		265	198	268	280	216	317	232	306	311	446	201	383	158	212	190	298	175	202	212	273
Sweden		836	1118	1077	1495	350	455	1432	1548	536	796	692	885	791	950	515	715	521	673	644	790
Total		1101	1316	1345	1775	566	772	1664	1854	847	1242	893	1268	949	1162	705	1013	696	875	856	1063

Table A.3: Annual number of individual wolves detected via non-invasive genetic sampling and included in the analysis. Numbers are reported by country, for females (F) and males (M), and for each type of sampling (structured and unstructured). We included only individuals associated with samples collected within the study area during the primary monitoring period (Oct 1 - Mar 31) between 2016/2017 and 2025/2026. Some individuals were detected in both countries during the same year, hence the sum of the national counts can exceed the total number of unique individuals detected in Scandinavia.

		2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		2021/2022		2022/2023		2023/2024		2024/2025		2025/2026	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
Norway	Structured	10	6	18	17	16	21	17	22	22	25	31	34	20	24	22	24	21	21	24	20
	Unstructured	46	36	47	44	31	43	36	44	38	52	40	48	27	20	23	31	23	22	20	21
Sweden	Structured	111	130	115	126	81	91	136	153	107	115	127	150	123	142	98	113	92	105	114	124
	Unstructured	104	123	120	144	69	69	146	158	98	115	121	131	134	144	96	115	94	119	99	102
Total	Structured	120	135	130	140	93	108	150	171	124	135	150	176	139	161	115	130	108	121	134	138
	Unstructured	142	154	161	178	96	107	178	194	135	162	152	170	158	162	119	143	114	139	117	120

Table A.4: Number of cause-specific dead recoveries of wolves in Scandinavia between 2016/17 and 2025/26 that were included in the OPSCR analysis. Numbers are reported by country, for females (F) and males (M). Note that dead recovery data from the final season (2025/2026) were not used in this analysis because survival is modeled as part of the transition from the start of one monitoring season to the start of the next one.

Country		2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		2021/2022		2022/2023		2023/2024		2024/2025		2025/2026	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
Other	Norway	0	0	1	3	1	1	1	1	0	2	1	2	0	1	2	1	0	0	0	0
	Sweden	4	5	8	2	4	5	5	7	8	5	1	6	3	2	5	6	1	8	1	4
Legal culling	Norway	6	6	18	16	10	8	6	12	11	16	14	22	10	9	11	11	3	7	0	1
	Sweden	17	23	12	31	4	4	6	12	18	30	22	18	39	40	20	27	24	25	0	4
Total	Total	27	34	39	52	19	18	18	32	37	53	38	48	52	52	38	45	28	40	1	9

Table A.5: Annual abundance estimates for wolf at several spatial scales: the entire study area, by country, by management unit (carnivore management regions in Norway and "Rovdjursförvaltningsområden" in Sweden), and counties ("Län" in Sweden); see also Figure A.2. Only counties and management units that are within or that intersect the study area are included in the table. Estimates are based on model-estimated activity center locations. Bayesian credible intervals (95%) are shown in parentheses. Small deviations between the total estimate and the sum of abundance estimates from the constituent subregions may arise due to rounding. Note that the numbers reported here are predictions from a statistical model which always represents an oversimplification of reality and is based on available data (NGS and dead recoveries). As a consequence, especially at the local scale, the model-estimated number of wolves based on DNA sampling can deviate from the number of wolves inferred from ancillary observations (e.g., camera traps).

	2016/2017	2017/2018	2018/2019	2019/2020	2020/2021	2021/2022	2022/2023	2023/2024	2024/2025	2025/2026
TOTAL	425.4 (412-441)	415.7 (406-426)	372.3 (356-391)	440.3 (432-450)	452.8 (439-468)	511.5 (496-529)	487.7 (473-504)	444.3 (428-463)	399.3 (386-416)	425.3 (402-453)
NORWAY	90.9 (82-101)	96.6 (89-105)	81 (74-88)	87.4 (79-96)	91.2 (83-100)	96.8 (87-107)	66.3 (58-76)	62.4 (55-71)	48.8 (42-56)	55.2 (48-65)
Region 2	1.8 (0-5)	0.7 (0-3)	1.2 (0-3)	1.3 (0-4)	0.4 (0-2)	0.8 (0-3)	0.7 (0-3)	0.7 (0-3)	0.6 (0-2)	0.6 (0-2)
Region 3	2.8 (0-7)	1.7 (0-4)	1.1 (0-3)	2 (0-5)	1.8 (0-5)	2.5 (0-6)	1.8 (0-5)	1.2 (0-4)	1.1 (0-3)	2 (0-5)
Region 4	18.2 (14-23)	22.4 (18-27)	24.6 (20-29)	28 (23-33)	22.9 (18-28)	27.3 (22-32)	24.3 (21-28)	22.6 (19-27)	14.6 (11-19)	16.6 (13-21)
Region 5	65.3 (59-72)	69.7 (64-76)	52.2 (47-58)	53.5 (48-60)	64.8 (58-72)	63.6 (56-72)	38.2 (32-46)	37.4 (32-44)	31.9 (27-38)	34.8 (29-41)
Region 6	2.9 (0-7)	2 (0-5)	1.9 (1-4)	2.6 (1-6)	1.4 (0-4)	2.6 (1-5)	1.3 (0-4)	0.6 (0-3)	0.7 (0-3)	1.4 (0-4)
SWEDEN	334.5 (322-349)	319.1 (310-329)	291.3 (275-310)	352.9 (346-361)	361.7 (348-377)	414.7 (399-432)	421.4 (409-436)	381.9 (366-398)	350.5 (337-365)	370.1 (350-394)
Norra	19.7 (14-26)	8.8 (5-14)	17.4 (11-24)	17.1 (13-21)	18.2 (13-24)	12.1 (7-18)	22 (17-28)	15.2 (10-21)	12.5 (7-18)	15.8 (9-24)
Jämtland	16.1 (11-22)	7.7 (4-12)	13.8 (9-20)	15.4 (11-20)	15.7 (11-21)	9.5 (5-15)	14.7 (10-20)	9.9 (6-15)	6.5 (3-11)	8.7 (3-15)
Västerbotten	0.1 (0-1)	0 (0-1)	0 (0-1)	0 (0-1)	0 (0-1)	0 (0-1)	0 (0-1)	0 (0-1)	0 (0-1)	0.1 (0-1)
Västernorrland	3.5 (1-7)	1.1 (0-3)	3.6 (0-8)	1.7 (0-4)	2.4 (0-6)	2.5 (0-6)	7.3 (5-11)	5.3 (2-9)	6 (4-9)	7 (4-12)
Mellersta	298.9 (288-310)	289.5 (282-298)	260.8 (248-276)	314.6 (308-323)	309.8 (297-324)	354.1 (341-368)	324.5 (313-338)	288.9 (276-303)	261.5 (250-275)	279.8 (263-298)
Dalarna	80.8 (74-88)	63.9 (58-70)	56.6 (48-66)	70.8 (65-77)	69.9 (62-79)	79 (71-88)	61.5 (54-70)	53.6 (46-62)	38.8 (32-46)	42.2 (33-53)
Gävleborg	50 (44-56)	54.2 (50-59)	46.2 (39-54)	69.3 (65-74)	60.9 (56-67)	51.2 (46-57)	34.1 (29-39)	21.6 (17-27)	21.5 (17-26)	25.2 (20-32)
Örebro	38.4 (34-44)	34.8 (31-39)	35.4 (29-42)	48.7 (43-54)	46.8 (40-54)	55.8 (49-63)	52.7 (47-59)	56.5 (51-63)	55.8 (50-62)	59.6 (54-66)
Stockholm	3.8 (2-7)	5 (3-7)	2.2 (0-5)	3.8 (2-6)	8.2 (5-11)	7.2 (4-11)	7.6 (4-11)	3.8 (2-6)	3.6 (2-6)	4.2 (2-7)
Uppsala	7.9 (4-12)	9.1 (6-13)	6.6 (4-10)	5.2 (3-7)	8.2 (5-11)	12.8 (10-16)	7.6 (5-11)	10.1 (8-13)	10.6 (9-13)	18 (15-22)
Värmland	85.6 (79-93)	87.7 (82-94)	83.7 (76-92)	83.6 (78-89)	78.5 (70-88)	104 (94-114)	101 (94-109)	72.9 (65-81)	48.7 (41-57)	43.6 (35-54)
Västmanland	25.7 (22-30)	26.2 (23-30)	16.3 (12-21)	27.3 (23-32)	19.1 (14-25)	24.5 (19-30)	27.8 (24-32)	24.1 (20-28)	36.3 (32-41)	36.5 (31-42)
Västra Götaland	6.6 (2-12)	8.7 (5-13)	13.7 (9-19)	6 (3-9)	18.3 (13-24)	19.6 (15-25)	32.4 (27-38)	46.4 (42-52)	46.2 (41-52)	50.4 (45-56)
Södra	16 (10-24)	20.8 (17-25)	13.1 (8-19)	21.2 (18-25)	33.7 (28-41)	48.5 (42-56)	74.8 (69-81)	77.8 (72-84)	76.6 (71-83)	74.5 (69-82)
Blekinge	1.4 (1-3)	0.2 (0-1)	0.2 (0-1)	0.1 (0-1)	0.2 (0-1)	0.4 (0-2)	0.1 (0-1)	0.4 (0-2)	0.4 (0-2)	0.3 (0-2)
Halland	0.5 (0-2)	0.3 (0-2)	0.4 (0-2)	0.2 (0-1)	1.4 (0-3)	0.7 (0-3)	1.2 (0-3)	2.5 (1-5)	3.5 (1-6)	2.3 (1-5)
Jönköping	1.7 (0-5)	0.5 (0-2)	0.9 (0-3)	1.2 (0-3)	2.8 (1-6)	7.6 (5-11)	14.1 (11-18)	10.7 (7-14)	11.9 (9-16)	15.5 (13-19)
Kalmar	1 (0-3)	1 (0-2)	1.5 (1-3)	0.7 (0-2)	1.7 (1-4)	1.3 (0-4)	0.8 (0-3)	1.4 (0-4)	1.8 (0-5)	1.9 (1-4)
Kronoberg	2.2 (1-5)	0.7 (0-3)	0.7 (0-3)	0.6 (0-2)	1.8 (1-4)	2 (1-4)	3.4 (2-5)	5.7 (3-8)	5.8 (3-9)	4.3 (1-8)
Östergötland	3.1 (1-6)	3.6 (3-6)	3.9 (2-7)	9.8 (9-12)	8.3 (6-12)	12 (9-16)	10.7 (9-14)	10.9 (8-14)	10.5 (8-14)	7.9 (5-11)
Skåne	2.7 (1-6)	2.3 (1-4)	0.6 (0-3)	0.4 (0-2)	2.7 (2-5)	5.2 (4-8)	12.5 (12-14)	18.3 (16-21)	15.4 (13-18)	14.2 (11-17)
Södermanland	3.5 (2-6)	12.2 (11-14)	5 (3-8)	8 (6-10)	14.8 (12-19)	19.4 (16-24)	32 (29-36)	27.8 (26-31)	27.3 (25-31)	28.1 (26-31)

Table A.6: Annual abundance estimates for wolf at the county level in Norway; see also Figure A.2. Only counties that are within or that intersect the study area are included in the table. Estimates are based on model-estimated activity center locations. Bayesian credible intervals (95%) are shown in parentheses. Small deviations between the total estimate and the sum of abundance estimates from the constituent subregions may arise due to rounding. Note that the numbers reported here are predictions from a statistical model which always represents an oversimplification of reality and is based on available data (NGS and dead recoveries). As a consequence, especially at the local scale, the model-estimated number of wolves based on DNA sampling can deviate from the number of wolves inferred from ancillary observations (e.g., camera traps).

	2016/2017	2017/2018	2018/2019	2019/2020	2020/2021	2021/2022	2022/2023	2023/2024	2024/2025	2025/2026
NORWAY	90.9 (82-101)	96.6 (89-105)	81 (74-88)	87.4 (79-96)	91.2 (83-100)	96.8 (87-107)	66.3 (58-76)	62.4 (55-71)	48.8 (42-56)	55.2 (48-65)
Akershus	12.5 (9-16)	13.3 (10-17)	16.5 (12-21)	16.9 (12-22)	15.8 (13-19)	20.3 (16-25)	22.2 (19-26)	19.5 (16-23)	13.3 (10-17)	13.5 (11-16)
Buskerud	1.1 (0-3)	0.5 (0-2)	1.1 (0-3)	1 (0-3)	0.2 (0-1)	0.6 (0-3)	0.5 (0-2)	0.5 (0-2)	0.4 (0-2)	0.4 (0-2)
Innlandet	68.1 (61-76)	71.4 (65-78)	53.3 (48-59)	55.5 (49-63)	66.6 (60-74)	66.1 (58-75)	40 (33-48)	38.5 (33-45)	32.9 (27-39)	36.7 (31-44)
Oslo	0.2 (0-1)	0.1 (0-1)	0.8 (0-2)	0.7 (0-2)	0 (0-0)	0 (0-1)	0.1 (0-1)	0.6 (0-2)	0.3 (0-1)	0.1 (0-1)
Østfold	5.5 (3-9)	9 (6-13)	7.3 (5-11)	10.3 (7-14)	7 (4-11)	6.9 (4-10)	2 (0-5)	2.5 (1-5)	1.1 (0-3)	3 (1-6)
Trøndelag	2.9 (0-7)	2 (0-5)	1.9 (1-4)	2.6 (1-6)	1.3 (0-4)	2.6 (1-5)	1.3 (0-4)	0.6 (0-3)	0.7 (0-3)	1.4 (0-4)
Vestfold	0.7 (0-3)	0.3 (0-2)	0.1 (0-1)	0.3 (0-2)	0.1 (0-1)	0.1 (0-1)	0.2 (0-1)	0.2 (0-1)	0.2 (0-1)	0.2 (0-1)

Table A.7: Annual population growth rate estimates for the wolf population in Scandinavia ("Total") and separately for Norway and Sweden. Estimates were derived using the posterior distributions of annual abundance estimates (Table 1) and indicate the change in estimated abundance at the start of one monitoring season to the start of the next one. Bayesian credible intervals (95%) are shown in parentheses.

	2016-2017	2017-2018	2018-2019	2019-2020	2020-2021	2021-2022	2022-2023	2023-2024	2024-2025
Norway	1.07 (0.93-1.21)	0.84 (0.74-0.95)	1.08 (0.95-1.22)	1.05 (0.91-1.19)	1.06 (0.93-1.21)	0.69 (0.58-0.81)	0.95 (0.79-1.12)	0.78 (0.65-0.94)	1.14 (0.93-1.38)
Sweden	0.95 (0.91-1.00)	0.91 (0.86-0.97)	1.21 (1.14-1.29)	1.02 (0.98-1.07)	1.15 (1.09-1.21)	1.02 (0.97-1.07)	0.91 (0.86-0.95)	0.92 (0.87-0.97)	1.06 (0.99-1.13)
Total	0.98 (0.94-1.02)	0.90 (0.85-0.95)	1.18 (1.12-1.24)	1.03 (0.99-1.07)	1.13 (1.08-1.18)	0.95 (0.91-1.00)	0.91 (0.87-0.96)	0.90 (0.85-0.95)	1.07 (1.00-1.14)

Table A.8: Estimates of the demographic parameters obtained from the sex-specific wolf OPSCR models. Parameters represent transition rates from Oct 1 to Sep 30 in the following year. State 2 and 3 represent other and scent-marking individuals, respectively. Median estimates and 95% Bayesian credible intervals (in parentheses) for per capita recruitment rate (ρ), survival (ϕ), transition probability from other to scent-marking adult state (ψ), mortality due to legal culling (h), recovered mortality due to other causes (rw), and mortality due to other causes not recovered (w) are presented for males (M) and females (F).

State		2016-2017		2017-2018		2018-2019		2019-2020		2020-2021	
		M	F	M	F	M	F	M	F	M	F
ρ	-	0.60 (0.51-0.68)	0.69 (0.59-0.79)	0.64 (0.54-0.76)	0.45 (0.36-0.55)	0.83 (0.69-0.97)	0.91 (0.75-1.05)	0.74 (0.66-0.84)	0.62 (0.55-0.72)	0.82 (0.70-0.95)	0.79 (0.67-0.91)
	2	0.45 (0.37-0.54)	0.57 (0.48-0.66)	0.47 (0.38-0.56)	0.50 (0.41-0.59)	0.60 (0.50-0.69)	0.56 (0.46-0.66)	0.46 (0.38-0.53)	0.52 (0.43-0.61)	0.51 (0.43-0.60)	0.56 (0.47-0.66)
ϕ	3	0.54 (0.42-0.65)	0.55 (0.44-0.66)	0.56 (0.45-0.67)	0.60 (0.49-0.71)	0.60 (0.48-0.71)	0.63 (0.51-0.74)	0.69 (0.58-0.79)	0.65 (0.55-0.76)	0.63 (0.52-0.73)	0.72 (0.61-0.81)
ψ	-	0.42 (0.31-0.54)	0.42 (0.31-0.53)	0.33 (0.22-0.46)	0.27 (0.17-0.38)	0.35 (0.25-0.46)	0.49 (0.36-0.62)	0.38 (0.28-0.50)	0.38 (0.27-0.50)	0.36 (0.26-0.47)	0.34 (0.24-0.46)
h	2	0.13 (0.09-0.19)	0.11 (0.07-0.17)	0.24 (0.17-0.31)	0.14 (0.09-0.21)	0.06 (0.03-0.11)	0.09 (0.05-0.15)	0.12 (0.08-0.17)	0.05 (0.02-0.10)	0.18 (0.13-0.24)	0.17 (0.11-0.24)
	3	0.10 (0.05-0.19)	0.11 (0.05-0.19)	0.18 (0.11-0.28)	0.13 (0.07-0.22)	0.07 (0.02-0.15)	0.07 (0.02-0.14)	0.07 (0.02-0.14)	0.07 (0.03-0.14)	0.18 (0.11-0.27)	0.09 (0.04-0.17)
rw	2	0.03 (0.01-0.06)	0.02 (0.00-0.05)	0.04 (0.02-0.08)	0.06 (0.03-0.11)	0.04 (0.02-0.08)	0.05 (0.02-0.10)	0.05 (0.02-0.09)	0.03 (0.01-0.07)	0.03 (0.01-0.06)	0.06 (0.03-0.11)
	3	0.02 (0.00-0.07)	0.03 (0.01-0.09)	0.01 (0.00-0.05)	0.02 (0.00-0.07)	0.03 (0.00-0.08)	0.01 (0.00-0.05)	0.01 (0.00-0.05)	0.03 (0.01-0.09)	0.04 (0.01-0.10)	0.01 (0.00-0.05)
w	2	0.38 (0.30-0.47)	0.30 (0.21-0.39)	0.25 (0.17-0.33)	0.29 (0.21-0.38)	0.30 (0.21-0.40)	0.30 (0.21-0.40)	0.37 (0.29-0.45)	0.39 (0.31-0.48)	0.28 (0.20-0.36)	0.20 (0.13-0.29)
	3	0.33 (0.22-0.45)	0.30 (0.20-0.41)	0.24 (0.15-0.34)	0.24 (0.15-0.34)	0.29 (0.19-0.41)	0.28 (0.18-0.39)	0.23 (0.14-0.33)	0.23 (0.15-0.33)	0.14 (0.07-0.22)	0.17 (0.10-0.27)
State		2021-2022		2022-2023		2023-2024		2024-2025			
		M	F	M	F	M	F	M	F		
ρ	-	0.59 (0.50-0.68)	0.81 (0.72-0.92)	0.57 (0.49-0.66)	0.66 (0.55-0.79)	0.61 (0.52-0.70)	0.60 (0.49-0.73)	0.77 (0.65-0.91)	0.87 (0.69-1.09)		
	2	0.42 (0.34-0.49)	0.43 (0.35-0.52)	0.45 (0.37-0.54)	0.43 (0.35-0.52)	0.48 (0.39-0.56)	0.39 (0.31-0.49)	0.52 (0.42-0.63)	0.53 (0.42-0.63)		
ϕ	3	0.58 (0.47-0.68)	0.59 (0.48-0.69)	0.59 (0.48-0.69)	0.62 (0.51-0.72)	0.61 (0.50-0.71)	0.61 (0.50-0.71)	0.54 (0.42-0.65)	0.65 (0.54-0.76)		
ψ	-	0.40 (0.29-0.52)	0.53 (0.41-0.65)	0.37 (0.26-0.48)	0.36 (0.25-0.48)	0.37 (0.26-0.49)	0.49 (0.35-0.63)	0.44 (0.32-0.58)	0.45 (0.32-0.59)		
h	2	0.16 (0.11-0.21)	0.16 (0.11-0.22)	0.22 (0.16-0.29)	0.24 (0.18-0.31)	0.20 (0.14-0.27)	0.15 (0.10-0.21)	0.17 (0.12-0.24)	0.15 (0.09-0.22)		
	3	0.11 (0.06-0.19)	0.13 (0.07-0.21)	0.15 (0.08-0.23)	0.13 (0.07-0.21)	0.12 (0.06-0.20)	0.12 (0.06-0.20)	0.12 (0.06-0.21)	0.11 (0.05-0.18)		
rw	2	0.04 (0.02-0.07)	0.02 (0.00-0.04)	0.02 (0.01-0.05)	0.01 (0.00-0.03)	0.04 (0.02-0.08)	0.04 (0.02-0.09)	0.06 (0.03-0.11)	0.01 (0.00-0.04)		
	3	0.02 (0.00-0.06)	0.01 (0.00-0.04)	0.01 (0.00-0.04)	0.03 (0.01-0.08)	0.03 (0.01-0.09)	0.02 (0.00-0.06)	0.01 (0.00-0.05)	0.01 (0.00-0.04)		
w	2	0.39 (0.31-0.47)	0.38 (0.30-0.47)	0.30 (0.23-0.39)	0.31 (0.23-0.40)	0.28 (0.20-0.37)	0.41 (0.32-0.51)	0.24 (0.15-0.34)	0.30 (0.20-0.41)		
	3	0.28 (0.19-0.39)	0.26 (0.17-0.36)	0.25 (0.16-0.35)	0.21 (0.13-0.31)	0.23 (0.15-0.33)	0.24 (0.16-0.35)	0.32 (0.22-0.43)	0.22 (0.14-0.33)		

Table A.9: Estimates of the spatial and movement process parameters obtained from the sex-specific wolf OPSCR models. β_{dens} represents the effect of the number of known wolf packs on AC locations (Bischof et al., 2020). The scale parameter σ of the detection function is expressed in kilometers and estimated separately for other (σ_1) and scent-marking (σ_2) individuals. τ represents the mean of the exponential movement parameter, describing individual movement distances between years, estimated separately for other (τ_1) and scent-marking (τ_2) individuals. Bayesian credible intervals (95%) are shown in parentheses. Parameters that were assumed to be constant across years are marked with *.

2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		
	M	F	M	F	M	F	M	F	M	F
β_{dens}^*	0.31 (0.29-0.33)	0.35 (0.32-0.37)	0.31 (0.29-0.33)	0.35 (0.32-0.37)	0.31 (0.29-0.33)	0.35 (0.32-0.37)	0.31 (0.29-0.33)	0.35 (0.32-0.37)	0.31 (0.29-0.33)	0.35 (0.32-0.37)
σ_1	7.61 (7.19-8.04)	7.92 (7.48-8.40)	7.76 (7.38-8.17)	8.09 (7.69-8.52)	7.17 (6.61-7.80)	6.77 (6.17-7.43)	7.86 (7.55-8.19)	7.63 (7.31-7.96)	7.76 (7.28-8.28)	7.35 (6.82-7.93)
σ_2	7.89 (7.54-8.27)	7.96 (7.58-8.36)	7.98 (7.67-8.30)	7.40 (7.07-7.76)	8.55 (8.07-9.05)	8.24 (7.70-8.82)	8.15 (7.81-8.51)	7.76 (7.44-8.10)	7.68 (7.32-8.07)	7.25 (6.86-7.67)
τ_1^*	69.72 (67.41-72.30)	59.17 (56.92-61.72)	69.72 (67.41-72.30)	59.17 (56.92-61.72)	69.72 (67.41-72.30)	59.17 (56.92-61.72)	69.72 (67.41-72.30)	59.17 (56.92-61.72)	69.72 (67.41-72.30)	59.17 (56.92-61.72)
τ_2^*	7.60 (7.06-8.18)	4.20 (3.85-4.58)	7.60 (7.06-8.18)	4.20 (3.85-4.58)	7.60 (7.06-8.18)	4.20 (3.85-4.58)	7.60 (7.06-8.18)	4.20 (3.85-4.58)	7.60 (7.06-8.18)	4.20 (3.85-4.58)
2021/2022		2022/2023		2023/2024		2024/2025		2025/2026		
	M	F	M	F	M	F	M	F	M	F
β_{dens}^*	0.31 (0.29-0.33)	0.35 (0.32-0.37)	0.31 (0.29-0.33)	0.35 (0.32-0.37)	0.31 (0.29-0.33)	0.35 (0.32-0.37)	0.31 (0.29-0.33)	0.35 (0.32-0.37)	0.31 (0.29-0.33)	0.35 (0.32-0.37)
σ_1	6.80 (6.38-7.26)	8.02 (7.51-8.58)	6.31 (5.92-6.74)	7.47 (6.97-8.02)	6.76 (6.27-7.31)	7.46 (6.85-8.15)	7.30 (6.78-7.87)	7.81 (7.20-8.49)	6.96 (6.45-7.54)	7.87 (7.28-8.53)
σ_2	7.66 (7.32-8.02)	7.83 (7.41-8.29)	7.42 (7.07-7.79)	6.60 (6.28-6.94)	8.32 (7.90-8.76)	7.04 (6.65-7.47)	7.56 (7.16-7.99)	7.27 (6.84-7.74)	7.85 (7.45-8.27)	7.07 (6.67-7.49)
τ_1^*	69.72 (67.41-72.30)	59.17 (56.92-61.72)	69.72 (67.41-72.30)	59.17 (56.92-61.72)	69.72 (67.41-72.30)	59.17 (56.92-61.72)	69.72 (67.41-72.30)	59.17 (56.92-61.72)	69.72 (67.41-72.30)	59.17 (56.92-61.72)
τ_2^*	7.60 (7.06-8.18)	4.20 (3.85-4.58)	7.60 (7.06-8.18)	4.20 (3.85-4.58)	7.60 (7.06-8.18)	4.20 (3.85-4.58)	7.60 (7.06-8.18)	4.20 (3.85-4.58)	7.60 (7.06-8.18)	4.20 (3.85-4.58)

Table A.10: Estimates of the detection process parameters for the structured sampling $\beta_{1Structured}$ corresponds to the effect of previous detection of an individual on detection probability, $\beta_{2Structured}$ to the effect of search-effort (track length) on detection probability, and $\beta_{3Structured}$ to the effect of average snow cover during the monitoring period on detection probability ($p_{0Structured}$). Coefficients are associated with scaled covariates. Bayesian credible intervals (95%) are shown in parentheses.

2016/2017			2017/2018			2018/2019			2019/2020			2020/2021	
	M	F	M	F	M	F	M	F	M	F	M	F	
$\beta_{1Structured}$	0.41 (0.11-0.71)	0.36 (0.06-0.66)	-0.11 (-0.37-0.17)	0.33 (0.06-0.61)	-0.36 (-0.86-0.08)	-0.02 (-0.39-0.38)	0.07 (-0.17-0.30)	0.04 (-0.19-0.27)	0.70 (0.37-1.00)	0.31 (-0.07-0.69)			
$\beta_{2Structured}$	0.29 (0.22-0.37)	0.32 (0.25-0.39)	0.34 (0.28-0.40)	0.32 (0.25-0.38)	0.33 (0.24-0.42)	0.36 (0.27-0.45)	0.42 (0.35-0.48)	0.36 (0.28-0.43)	0.25 (0.19-0.31)	0.26 (0.19-0.33)			
$\beta_{3Structured}$	-0.23 (-0.49-0.03)	0.02 (-0.31- 0.33)	0.05 (-0.27-0.39)	0.34 (-0.05- 0.73)	0.16 (-0.17-0.50)	0.02 (-0.39- 0.46)	0.03 (-0.16-0.23)	0.09 (-0.12- 0.29)	0.34 (0.15-0.54)	0.44 (0.20- 0.70)			
2021/2022			2022/2023			2023/2024			2024/2025			2025/2026	
	M	F	M	F	M	F	M	F	M	F	M	F	
$\beta_{1Structured}$	0.45 (0.16-0.73)	-0.06 (-0.33-0.22)	0.20 (-0.08-0.47)	0.11 (-0.19-0.41)	-0.04 (-0.37-0.29)	-0.08 (-0.45-0.29)	-0.08 (-0.44-0.28)	0.02 (-0.32-0.37)	-0.02 (-0.34-0.30)	0.43 (0.12-0.74)			
$\beta_{2Structured}$	0.28 (0.22-0.35)	0.28 (0.22-0.35)	0.35 (0.28-0.43)	0.34 (0.27-0.41)	0.16 (0.11-0.22)	0.26 (0.18-0.35)	0.43 (0.35-0.52)	0.42 (0.33-0.50)	0.42 (0.35-0.50)	0.44 (0.37-0.51)			
$\beta_{3Structured}$	0.41 (0.20-0.63)	0.17 (-0.07- 0.41)	0.08 (-0.09-0.26)	0.14 (-0.06- 0.35)	0.24 (0.05-0.44)	-0.14 (-0.34- 0.07)	-0.08 (-0.33-0.17)	-0.04 (-0.30- 0.22)	-0.04 (-0.21-0.14)	-0.21 (-0.37-0.05)			

Table A.11: Estimates of the detection process parameters for the unstructured sampling $\beta_{1Unstructured}$ corresponds to the effect of previous detection of an individual on baseline detection probability, $\beta_{2Unstructured}$ to the effect of distance to the nearest roads on detection probability, $\beta_{3Unstructured}$ to the effect of average snow cover during the monitoring period on detection probability, and $\beta_{4Unstructured}$ to the effect of spatio-temporal heterogeneity in unstructured sampling derived using the observation data in Skandobs and Rovbase. Coefficients are associated with scaled covariates. Bayesian credible intervals (95%) are shown in parentheses.

	2016/2017		2017/2018		2018/2019		2019/2020		2020/2021	
	M	F	M	F	M	F	M	F	M	F
$\beta_{1Unstructured}$	0.24 (0.02-0.48)	0.48 (0.25-0.71)	0.29 (0.10-0.49)	0.73 (0.53-0.94)	0.01 (-0.34-0.36)	0.83 (0.44-1.24)	0.32 (0.13-0.52)	0.53 (0.34-0.72)	0.82 (0.54-1.11)	0.59 (0.26-0.91)
$\beta_{2Unstructured}$	-0.25 (-0.80- 0.18)	-0.11 (-0.63- 0.34)	-1.26 (-2.07-0.60)	-0.83 (-1.40-0.36)	0.07 (-0.54- 0.53)	0.50 (-0.15- 1.10)	-0.47 (-1.20- 0.08)	-0.24 (-1.11- 0.45)	-0.58 (-1.44- 0.06)	0.68 (-0.71- 2.53)
$\beta_{3Unstructured}$	0.01 (-0.17- 0.20)	0.07 (-0.13- 0.27)	-0.11 (-0.33- 0.10)	-0.16 (-0.42- 0.10)	-0.45 (-0.72-0.17)	-0.75 (-1.05-0.44)	0.02 (-0.09- 0.13)	0.11 (-0.01- 0.23)	-0.16 (-0.32- 0.00)	-0.21 (-0.39-0.03)
$\beta_{4Unstructured}$	2.39 (1.85-2.96)	2.35 (1.84-2.94)	1.70 (1.39-2.05)	1.82 (1.48-2.18)	1.88 (1.36-2.55)	1.82 (1.30-2.46)	2.32 (1.91-2.79)	1.82 (1.45-2.23)	1.65 (1.23-2.13)	2.11 (1.51-2.72)
	2021/2022		2022/2023		2023/2024		2024/2025		2025/2026	
	M	F	M	F	M	F	M	F	M	F
$\beta_{1Unstructured}$	0.15 (-0.14-0.43)	0.41 (0.13-0.69)	0.49 (0.24-0.74)	0.49 (0.21-0.75)	0.14 (-0.17-0.46)	0.75 (0.41-1.10)	0.42 (0.10-0.74)	0.40 (0.10-0.71)	0.46 (0.12-0.80)	0.58 (0.26-0.92)
$\beta_{2Unstructured}$	0.06 (-0.42- 0.44)	-1.28 (-3.28- 0.12)	0.26 (-0.43- 0.93)	-0.36 (-1.70- 0.70)	0.11 (-0.60- 0.63)	0.82 (0.03- 2.72)	1.16 (0.09- 3.00)	0.09 (-1.48- 2.34)	4.52 (3.33- 4.98)	0.26 (-0.56- 1.17)
$\beta_{3Unstructured}$	-0.11 (-0.27- 0.04)	-0.24 (-0.40-0.07)	-0.46 (-0.58-0.34)	-0.31 (-0.43-0.19)	-0.19 (-0.34-0.04)	-0.34 (-0.49-0.18)	-0.24 (-0.43-0.06)	-0.22 (-0.41-0.02)	-0.14 (-0.30- 0.02)	-0.37 (-0.52-0.21)
$\beta_{4Unstructured}$	2.38 (1.80-3.06)	1.94 (1.41-2.51)	1.77 (1.30-2.36)	1.38 (0.96-1.82)	2.05 (1.54-2.71)	2.17 (1.63-2.82)	1.76 (1.19-2.40)	2.57 (1.88-3.51)	2.22 (1.60-3.03)	2.46 (1.82-3.18)

Table A.12: Average proportion of individuals detected via non-invasive genetic sampling (NGS) in Scandinavia. Values were calculated as the number of individuals detected with NGS (Table A.3) divided by the total and sex-specific abundance estimates obtained from the OPSCR models (??). Bayesian credible intervals (95%) are shown in parentheses.

	2016/2017	2017/2018	2018/2019	2019/2020	2020/2021	2021/2022	2022/2023	2023/2024	2024/2025	2025/2026
M	0.85 (0.81-0.89)	0.92 (0.89-0.95)	0.78 (0.72-0.83)	0.94 (0.90-0.96)	0.83 (0.79-0.87)	0.85 (0.80-0.89)	0.87 (0.83-0.91)	0.88 (0.84-0.92)	0.87 (0.84-0.91)	0.85 (0.80-0.91)
F	0.86 (0.81-0.89)	0.90 (0.87-0.93)	0.77 (0.72-0.81)	0.93 (0.90-0.96)	0.86 (0.82-0.89)	0.87 (0.83-0.90)	0.85 (0.81-0.89)	0.77 (0.72-0.82)	0.80 (0.75-0.84)	0.81 (0.73-0.88)
Total	0.85 (0.82-0.88)	0.91 (0.89-0.93)	0.77 (0.74-0.81)	0.93 (0.91-0.95)	0.84 (0.82-0.87)	0.86 (0.83-0.88)	0.86 (0.84-0.89)	0.83 (0.79-0.86)	0.84 (0.80-0.87)	0.83 (0.78-0.88)