Norwegian University of Life Sciences
Faculty of Environmental Sciences and Natural Resource Management

2026
ISSN 2535-2806

MINA fagrapport 115

Estimates of brown bear density, abundance, and population dynamics in Norway 2016 - 2025

Pierre Dupont
Nina Rosita Hansen
Cyril Milleret
Asunción Semper-Pascual
Henrik Brøseth
Alexander Kopatz
Jonas Kindberg
Richard Bischof



Dupont, P., Hansen, N.R., Milleret, C., Semper-Pascual, A., Brøseth, H., Kopatz, A., Kindberg, J. and Bischof, R. 2026. **Estimates of brown bear density, abundance, and population dynamics in Norway 2016 - 2025** - MINA fagrapport 115. 35pp.

Ås, September 2026

ISSN: 2535-2806

COPYRIGHT
© Norwegian University of Life Sciences (NMBU)
The publication may be freely cited where the source is acknowledged

AVAILABILITY
Open

PUBLICATION TYPE
Digital document (pdf)

QUALITY CONTROLLED BY
The Research committee (FU), MINA, NMBU

PRINCIPAL
Miljødirektoratet, Ref: 23047064, Contact person: Terje Bø

COVER PICTURE
Female brown bear with cub, Staffan Widstrand Photography.

NØKKELORD
Ursus arctos, brunbjørn, tetthet, populasjonsdynamikk, deteksjonssannsynlighet, ikke-invasiv genetisk prøveinnsamling, åpen populasjon romlig fangst-gjenfangst, rovdyrforvaltning

KEY WORDS
Ursus arctos, brown bear, population density, population dynamics, detection probability, non-invasive genetic sampling, open-population spatial capture-recapture, carnivore management

Pierre Dupont, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Nina Rosita Hansen, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Cyril Milleret, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Asunción Semper-Pascual, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Henrik Brøseth, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway.

Alexander Kopatz, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway.

Jonas Kindberg, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway.


Richard Bischof (richard.bischof@nmbu.no), Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Summary

Background The Scandinavian brown bear (*Ursus arctos*) population is monitored annually in Norway using non-invasive genetic sampling (NGS) and recovery of dead individuals. DNA extracted from faeces, urine, hair, saliva or tissue is used to identify the species, sex and individual from which each sample originated. These data are compiled annually in the Scandinavian large carnivore database Rovbase 3.0 (rovbase.no, rovbase.se).

Approach Using the Bayesian open-population spatial capture-recapture (OPSCR) model developed by RovQuant, we estimated the population dynamics of the Norwegian portion of the Scandinavian brown bear population between 2016 and 2025. We provide annual density maps, as well as estimates of jurisdiction-specific population sizes, cause-specific survival, recruitment, and detection probabilities. Associated uncertainties are reported with all estimates.

Results We estimated that, within the study area (228 100 km^2), the Norwegian brown bear population was likely (95% credible interval) made up of between 150 and 186 individuals in 2025. Each year, a large proportion of bears detected in Norway can be attributed to neighbouring countries. Specifically, in 2025, between 41 and 59 (95% credible interval) of the individuals detected in Norway were attributed to neighbouring countries (Sweden = 16 to 29, Finland = 8 to 19, Russia = 9 to 19). The OPSCR results also highlight the ongoing recovery of the brown bear population in Norway, with an overall increase in population size, driven primarily by increases in abundance in carnivore regions 5 (Hedmark) and 8 (Troms and Finnmark).




Density map and estimated brown bear abundance range in 2025

Sammendrag

Bakgrunn Den skandinaviske bestanden av brunbjørn (*Ursus arctos*) blir overvåket årlig ved bruk av ikke-invasiv genetisk prøveinnsamling (NGS) og gjenfunn av døde individer. DNA isolert fra skit, urin, hår, spytt og vev brukes til å identifisere art, kjønn og individ for hver enkelt prøve. Denne informasjonen samles og ivaretas i Rovbase 3.0 (rovbase.se, rovbase.no).

Tilnærming Ved bruk av en Bayesiansk åpen romlig fangst-gjenfangst populasjonsmodell (OPSCR) utviklet av RovQuant, estimerte vi bestandsdynamikken til den norske delen av den skandinaviske bjørnebestanden. Metoden gjør det mulig å produsere årlige tettshetskart, estimater på bestandsstørrelse, samt estimater for overlevelses-, rekrutterings- og deteksjonssannsynlighet med tilhørende usikkerhet for brunbjørn i Norge mellom 2016 og 2025.

Resultater Basert på OPSCR-modellen var den norske brunbjørnbestanden i 2025 sannsynligvis (95% kredibelt intervall) på mellom 150 og 186 individer i hovedutbredelsesområdet (228 100 km^2). Årlig kan en stor andel av bjørnene påvist i Norge ha sitt aktivitetssentrum i nabolandene. I 2025 ble mellom 41 og 59 (95% kredibelt intervall) bjørner påvist i Norge estimert å ha sitt aktivitetssentrum i Sverige (16-29), Finland (8-19) eller Russland (9-19). OPSCR-resultatene viser en pågående vekst av brunbjørnbestanden i Norge. Bestandsøkningen av bjørn er hovedsakelig drevet av en sterkere økning i rovviltregioner 5 (Hedmark) og 8 (Troms og Finnmark) de siste 10 årene.



Kart som viser tetthet av brunbjørn med intervall for estimert antall brunbjørn i 2025

Contents

1 Introduction **7**

2 Methods **8**
2.1 Data 8
2.2 Open-population spatial capture-recapture model 10

3 Results **14**
3.1 Non-invasive genetic samples and dead recoveries 14
3.2 Density and abundance 14
3.3 Vital rates 18
3.4 Detection probability 18

4 Discussion **19**

5 Summary of improvements made **19**

6 Suggestions for future improvements **21**

7 Acknowledgements **21**

8 Data availability **21**

References **23**

Appendices **25**

1 Introduction

Non-invasive genetic sampling (NGS) and dead recoveries are a centerpiece of national and regional large carnivore monitoring in Norway and Sweden. Over nearly two decades, both countries have accumulated one of the largest and most extensive individual-based data sets on large carnivores in the world, and plan to continue such transnational monitoring in the future. Each year, Rovdata (rovdata.no) compiles, analyzes, and reports monitoring data on the Eurasian lynx (*Lynx lynx*), wolverine (*Gulo gulo*), brown bear (*Ursus arctos*), wolf (*Canis lupus*) and golden eagle (*Aquila chrysaetos*) in Norway. NGS data and other sources of information are made publicly available and stored in the Scandinavian large carnivore database Rovbase 3.0 (rovbase.se, rovbase.no).

Since 2017, project RovQuant (nmbu.no/en/research/projects/rovquant) has been developing statistical methods to exploit the potential of this database and allow a comprehensive assessment of the population status and dynamics of three large carnivore species, namely the brown bear, wolf, and wolverine. At the core of this analytical framework (Bischof et al., 2019b, 2020a) are Bayesian spatial capture-recapture (SCR) and open-population spatial capture-recapture (OPSCR) models (Ergon and Gardner, 2014; Bischof et al., 2016; Chandler et al., 2018; Dupont et al., 2021). SCR models use the spatial information contained in the repeated detection of individuals to estimate their location and population density. In addition, OPSCR models use the temporal information contained in data collected over multiple years to also estimate recruitment and survival probabilities, as well as inter-annual movements. Importantly, this approach accounts for imperfect detection, i.e., the probability that an individual present in the population remains undetected. The method ultimately generates density maps and vital rate estimates, from which abundance estimates can be derived for any spatial subunit (e.g., county or management region). Another crucial feature of this method is that all estimates are accompanied by their associated uncertainty, based on the model and data used (Bischof et al., 2019b, 2020b).

Since 2019, RovQuant has published annual reports for wolves and wolverines based on the Rovbase data and OPSCR models (last reports for wolves: Dupont et al. 2026, and wolverines: Milleret et al. 2026). In addition, since 2023, RovQuant has reported results about the status and population dynamics of the brown bear population in Norway (Dupont et al., 2023a,b, 2024b, 2025). Although RovQuant aims to provide comprehensive analyses of large carnivore populations throughout their entire range in Norway and Sweden, we only analyzed and present results for the Norwegian brown bear monitoring data in this report. Previous analyses revealed multiple difficulties related to the Swedish brown bear non-invasive genetic monitoring data (Dupont et al., 2024a; Milleret et al., 2024), leading among other things to inconsistent results across model permutations when the Scandinavian brown bear monitoring data were analyzed jointly for Norway and Sweden (Dupont et al., 2023a).

In the current report, we thus present an updated analysis of a 10-year time series (2016-2025) of the Norwegian portion of the Scandinavian brown bear population using the latest available bear monitoring data from Norway and the most recent version of the OPSCR model. We provide the following information:

- annual and sex-specific estimates of the number of bears (with credible intervals) for large carnivore regions, (*"forvaltningsregioner for rovvilt"*), counties (*"fylker"*), and for the entire study area within the country of Norway.
- estimated proportion of individuals detected through NGS,
- bear density maps throughout Norway,
- annual estimates of mortality, recruitment, and population growth rate, and
- uncertainty estimates associated with all quantitative results.

Box 1: Terms and acronyms used

AC: Activity center. Model-based equivalent of the center of an individual's home range during the monitoring period. "AC location" refers to the spatial coordinates of an individual AC in a given year and "AC movement" to the movement of an individual AC between consecutive years.
CrI: 95% credible interval associated with a posterior sample distribution.
Detectors: Potential detection locations in the spatial capture-recapture framework. These can refer to fixed locations (e.g., camera-trap locations) or in this report to areas searched (e.g., habitat grid cells where searches for genetic samples were conducted). The searched area was defined as a 50 km buffer around all NGS data collected during the period considered.
Habitat buffer: Buffer surrounding the searched area that is considered potentially suitable habitat but was not searched.
Länsstyrelserna: Swedish County Administrative Boards, in charge of the monitoring of large carnivores at the county level.
Legal culling: Lethal removal of individuals by legal means, including licensed recreational hunting, management removals, and defense of life and property.
LLCC: Local Large Carnivore Contact (Rovviltkontakt). A person in the local community that is employed by the Norwegian Nature Inspectorate on a limited basis. The role as a LLCC is primarily to carry out investigations, including necropsies of domestic animals in the field, to determine if a large carnivore is the primary cause of death or injury. LLCC are also employed to monitor large carnivores during winter.
MCMC: Markov chain Monte Carlo.
Naturvårdsverket: The Swedish Environmental Protection Agency, in charge of conducting and coordinating Sweden's environmental work.
NGS: Non-invasive genetic sampling.
OPSCR: Open-population spatial capture-recapture.
p_0**:** Baseline detection probability; probability of detecting an individual at a given detector, if the individual's AC is located exactly at the detector location.
$\boldsymbol{\sigma}$: Scale parameter of the detection function; related to the size of the circular home-range.
RCLC: Regional Coordinator for Large Carnivore Monitoring and Documentation (Regionalt fagansvarlig rovviltdokumentasjon). A person in full employment in the Norwegian Nature Inspectorate with a coordinating function in the large carnivore monitoring at the regional level.
SCR: Spatial capture-recapture.
SNO: Statens naturoppsyn (Norwegian Nature Inspectorate) is the operative field branch of the Norwegian Environment Directorate (Miljødirektoratet).
Statsforvalteren: Norwegian state's representative in the county, responsible for following up decisions, goals, and guidelines from the legislature and the government.
RovQuant: Research project at the Norwegian University of Life Sciences (Ås, Norway) that develops and applies OPSCR models.

2 Methods

2.1 Data

For this analysis, we relied on data from multiple sources, the primary one being the Scandinavian large carnivore database Rovbase 3.0 (rovbase.se and rovbase.no; last extraction: 2026-04-18). This database is used jointly by Norway and Sweden to record detailed information associated with large carnivore monitoring, including, but not limited to, non-invasive genetic sampling (NGS) data and dead recoveries. In the following sections, we describe the various types of data used in the analysis.

Non-invasive genetic sampling The Norwegian Nature Inspectorate (Statens Naturoppsyn,

SNO) has primary responsibility for the collection of brown bear non-invasive genetic samples in Norway. Non-invasive genetic sampling consists of extracting DNA from samples left behind by the target species, without directly handling, capturing, or disturbing the animal itself, such as faeces and hair. This is accomplished by the regional coordinators for large carnivore monitoring and documentation (RCLC) and a network of local large carnivore contacts (LLC) through targeted collection of samples via snow tracking, visits to recently used dens and in connection with depredation investigations and observation reports from the public. In addition, samples are collected by hikers, landowners and others throughout the year, as well as by hunters in the fall. For this analysis, we included samples collected in Norway between April 1 and November 30.

All samples were analysed with a 96-marker SNP (single nucleotide polymorphism) assay: 85 autosomal SNPs, four mitochondrial and seven sex-specific SNPs (four Y-chromosomal and three X-chromosomal SNPs) for sex and individual identification (Norman and Spong, 2015; Norman et al., 2013). Due to potentially low DNA quality in non-invasively collected material, all samples, except tissue samples, were analysed in two independent SNP-genotyping runs. The results for each marker from the two independent runs had to be identical to allow for a consensus DNA- or SNP-profile. To assign a new individual, a SNP-profile had to consist of at least 75 autosomal SNPs, sex, and seven microsatellite markers (see below). For already known and previously identified individuals, a SNP-profile of at least 66 SNPs was considered sufficient for individual identification. For SNP-profiles with less than 66 SNPs and at least three mitochondrial SNPs, no identification was assigned but the sample was categorised as brown bear. Samples, or profiles, with less than 66 SNPs were not categorised as brown bear. Additionally, one sample from each newly identified individual was analysed using the eight microsatellite markers previously used for DNA-based monitoring in Norway and Sweden: G10L, Mu05, Mu09, Mu10, Mu23, Mu50, Mu51 og Mu59; (Paetkau and Strobeck, 1998; Taberlet et al., 1996), as well as three sex-specific microsatellite markers (Bidon et al., 2013) to check against DNA-profiles of previously identified individuals in Norway and Sweden. All confirmed new individuals were then analysed with an additional four microsatellite markers (Andreassen et al., 2012), to keep the initial microsatellite monitoring database up-to-date (Brøseth et al., 2026).

Dead recoveries In Scandinavia, all brown bears killed legally (e.g., legal hunting, management kills, defense of life and property) have to be reported to the management authorities (Statsforvalteren or SNO in Norway and Länsstyrelserna or the police in Sweden). Although some mortalities due to other causes (e.g., natural deaths, vehicle and train collisions, illegal hunting) are also reported, an unknown proportion remains undetected. Tissue is collected from all reported dead bears for DNA extraction and analysis. DNA from dead bears, if genotyped, can be linked with NGS data via individual IDs and provide definite information about the fate of individuals. Dead recoveries associated with geographic coordinates also provide additional information for the estimation of individual locations, especially valuable for individuals with few or no NGS detections (Dupont et al., 2021).

Observation reports in Skandobs We used all observation records in the Skandobs database that were recorded during the bear monitoring season since 2016 (skandobs.se, skandobs.no; last extraction: 2026-04-18). Skandobs is a web application developed in collaboration between Rovdata in Norway and Naturvårdsverket in Sweden, that allows anyone to register observations (e.g., visual, tracks, faeces, etc.) of bears, lynx, wolves, and wolverines in Scandinavia. This data set currently consists of around 120 000 records of possible large carnivore observations. Although most observations are not verified, they offer the best available proxy for spatio-temporal variation in opportunistic effort (see below).

2.2 Open-population spatial capture-recapture model

We analysed the data collected between 2016 and 2025 using a Bayesian open-population spatial capture-recapture (OPSCR) model (Bischof et al., 2020b; Dupont et al., 2024a, 2023b). OPSCR models allow the simultaneous analysis of NGS data from multiple years and provide estimates of vital rates and individual movements between monitoring seasons in addition to annual densities. OPSCR models can be extended to incorporate dead recovery data to improve inferences (Dupont et al., 2021). The Bayesian OPSCR model (Bischof et al., 2019b) we developed and used addresses three challenges associated with population-level wildlife inventories:

1. Detection is imperfect and sampling effort heterogeneous in space and time: not all individuals present in the study area are detected (Kéry and Schaub, 2011).
2. Individuals that reside primarily outside the surveyed area may be detected within it. This is especially relevant for the Norwegian bear population given the comparatively higher population densities on the Swedish side of the Norwegian-Swedish border (Bischof et al., 2016), and the proximity with the bear populations in Finland and Russia. Without this explicit link between population size and geographic area characteristic of the SCR framework, density cannot be estimated and population size is ill-defined (Efford, 2004).
3. Non-spatial population dynamic models (such as CR models) usually estimate "apparent" survival and recruitment, as these parameters include the probability of permanent emigration and immigration, respectively. By explicitly modelling movement of individuals between years, the OPSCR model can help return unbiased estimates of demographic parameters (Ergon and Gardner, 2014; Schaub and Royle, 2014; Gardner et al., 2018; Dupont et al., 2021; Efford and Schofield, 2022).

The OPSCR model (Ergon and Gardner, 2014; Bischof et al., 2016; Chandler et al., 2018) is composed of three sub-models:

1. A model for the spatial distribution of individuals in the population and their movement between years (i.e. density and movement).
2. A model for population dynamics (i.e. recruitment and mortality) and population size.
3. A model for detection of individuals during DNA searches and dead recoveries.

Density and movement sub-model We used a Bernoulli point process to model the distribution of individual activity centers (ACs, Zhang et al., 2022). In the first year, individuals were located according to an intensity surface, which was a function of two spatial covariates: the locations of all bears recovered dead throughout the 2016-2025 period (Figure A.2; see Bischof et al., 2019b and Bischof et al., 2020b for more details), and the presence/absence of all bear observations registered in SkandObs during the 2016-2025 period (Figure A.2). For all subsequent years ($t > 1$), the location of an individual's activity center was a function of the distance from its previous activity center (at time $t - 1$) as well as the two spatial covariates mentioned above. As in previous analyses (Dupont et al., 2023a,b, 2024b, 2025), we used a half-normal model to describe the movement propensity of individuals between years.

Population dynamics and population size sub-model We used a multi-state formulation (Lebreton and Pradel, 2002) where each individual life history is represented by a succession of up to 5 discrete states $z_{i,t}$: 1) "unborn" if the individual has not yet been recruited into the population (state "unborn" is required for the data augmentation procedure, see below); 2) "alive" if it is alive; 3) "recovered legal" if it was legally killed in Norway between the start of the current and the start of the next monitoring period; 4) "recovered other" if it died from other causes of mortality (e.g., disease, car collision) and was reported to the authorities; and 5) "dead" if it died due to another mortality cause but was not recovered, or if it was already dead in the previous monitoring season. We then modelled the transitions from one state to

another between consecutive monitoring seasons (t to $t+1$) to estimate vital rates (recruitment and cause-specific mortality). In order to account for the fact that some individuals might never be detected, we used data augmentation (Royle and Dorazio, 2012), whereby additional, undetected individuals are available for inclusion in the population at each time step.

Detection sub-model OPSCR models account for the spatial variation in detectability by modelling individual detection probability as a function of the distance between the coordinates of a detection (i.e. a detector) and the individual's AC location (estimated by the density sub-model). A half-normal function is generally used to express the declining probability of detection with increasing distance between the AC and the detector. We accounted for additional sources of spatial and temporal variation in individual detection probability by using several spatio-temporal covariates:

- Spatio-temporal variation in unstructured sampling (Figure A.3). For each detector grid cell and during each monitoring season (Apr 1 - Nov 30), we identified whether a) any carnivore sample had been registered in Rovbase and b) any observation of carnivores had been registered in Skandobs. Roughly, this binary variable distinguishes areas with very low detection probability from those with a higher probability that carnivore DNA samples could have been detected and submitted for genetic analysis.
- Spatial variation in accessibility measured as the average distance to the nearest road (Figure A.4).
- Spatio-temporal variation between three regions defined based on the aggregation of neighboring counties (Figure A.6).

More details about the various model components can be found in Bischof et al. (2019a,b, 2020b).

Model fitting We fitted sex-specific Bayesian OPSCR models using Markov chain Monte Carlo (MCMC) simulation with NIMBLE version 1.2.0 (Turek et al., 2021; de Valpine et al., 2017; NIMBLE Development Team, 2019) and nimbleSCR version 0.2.1 (Bischof et al., 2021) in R version 4.3.3 (R Core Team, 2022). We ran 4 chains of 100 000 iterations each, including a 25 000-iterations burn-in period. Due to the computing challenge of post-processing large amounts of data, we thinned chains by a factor of 10 from which abundance estimates were derived. We considered models as converged when the Gelman-Rubin diagnostics (Rhat, Gelman and Rubin, 1992) was $\leq$1.1 for all parameters and by visually inspecting trace plots.

Abundance estimates and density maps Here, we report abundance based on AC locations (Box 2). To obtain an estimate of abundance for any given area, we summed the number of predicted AC locations of live individuals that fell within that area for each iteration of the MCMC chains, thus generating a posterior distribution of abundance for that area. In this fashion, abundance estimates and the associated uncertainty can be extracted for any spatial unit, including countries, counties or management regions (Figure A.1). Individuals detected near a border can have their model-predicted AC placed on different sides of that border in different model iterations, even if detections are only available on one side of the border. As a result, the probability of designating such individuals to either side of the border is integrated into jurisdiction-specific abundance estimates. This is especially relevant for bears detected along the Swedish and Norwegian border ("trans-boundary bears", Box 2) as individual bears can be partially designated to both countries (Bischof et al., 2016). To ensure that abundance estimates for spatial sub-units add up to overall abundance estimate, we used the mean and associated 95% credible interval to summarize posterior distributions of abundance. Combined (female/male) parameter estimates were obtained by merging posterior samples from the sex-specific models.

We also used both the posterior distribution of model-estimated AC positions and the scale parameter (σ) of the detection function to construct density maps based on individual space-use (Box 2). These maps are not only based on the position of the center of an individual's home range but also take into account the area (defined by σ, Royle et al. 2014) over which that individual's activity is spread, i.e., its space use (Bischof et al., 2020b). To do so, we constructed individual space-use raster maps (5 km resolution), scaled values in each raster to sum to one, and then summed across all individual rasters (cell-by-cell) to create a single population-level raster map. By repeating this procedure for each MCMC iteration, we obtained a posterior distribution of population space use. The overall density map was then derived by calculating the mean abundance across iterations in each cell (Bischof et al., 2020b). Abundance for any given region can also be derived from these maps (Box 2), but note that this approach relies on the assumption of circular home ranges of identical size for all individuals of a given sex. In other words, it does not take into account individual variation in home-range size and shape.

Other derived parameters The proportion of the Norwegian bear population detected each year, together with the associated uncertainty, was obtained by dividing the number of individuals *with their estimated AC located in Norway* that were detected through NGS sampling in a given year (Table A.1) by the abundance estimates from Norway for each MCMC iteration in the same year (Table A.4). This generated posterior distributions of the proportion of the Norwegian bear population that had been detected, from which the annual means and associated 95% credible intervals could be extracted. This procedure was used to calculate female, male and overall proportions of individuals detected.

The number of bears detected in Norway that could be attributed to other countries was obtained by summing the number of individuals detected in Norway that had their AC located in Sweden, Finland or Russia for each MCMC sample. With this procedure, we were able to generate posterior distributions of the number of individuals detected in Norway that could be attributed to neighboring countries, from which the annual means and associated 95% credible intervals could be extracted. The proportion of the individuals detected in Norway that could be attributed to each country was then obtained by dividing these numbers by the total number of individuals detected through NGS sampling.

We derived the proportion of females in the population and the associated uncertainty by dividing the posterior distribution of the number of females by the combined abundance estimate for males and females (Table A.4). Yearly population growth rates were calculated as $\lambda_t = N_{t+1}/N_t$.

Focus on uncertainty Although in the tables we report mean estimates for abundance and median estimates for all other parameters, we intentionally focused the main results of this report on the 95% credible interval limits of the estimates. We did so with the aim of drawing the reader's attention to the uncertainty around population size estimates, rather than a single point estimate (Milleret et al., 2022).




Figure 1: Annual distribution of brown bear non-invasive genetic samples (NGS, yellow crosses) and dead recoveries (purple crosses) collected in Norway between 2016 and 2025 and included in the OPSCR analyses. We included only samples collected within Norway during the primary monitoring period (Apr 1 - Nov 30).

3 Results

3.1 Non-invasive genetic samples and dead recoveries

A total of 6554 (2804 female; 3750 male) genotyped bear genetic samples collected in Norway between 2016 and 2025 were included in the analysis (Figure 1). These samples were associated with 634 (222 female; 412 male) individuals. We also included 120 bears recovered dead in Norway in the OPSCR model, of which 105 (16 female; 89 male) were due to legal culling and 15 (7 female; 8 male) due to other causes of mortality. Annual tallies of NGS samples and associated individuals, as well as dead recoveries included in the analysis for the period 2016-2025 are provided in the Appendices (NGS samples and individuals: Table A.1, dead recoveries: Table A.2).

3.2 Density and abundance

Brown bear abundance for the entire study area in Norway (228 100 km^2, excluding the buffer area) was estimated (95% credible interval) to be between 150 and 186 in 2025, based on the estimated locations of individual ACs (Table 1, Figure 2). Estimates refer to the status of the population at the start of the annual sampling period (April 1). The Norwegian bear population size increased over the 10-year study period with yearly population growth rates λ_t varying between 0.80-1.03 (95% credible interval) in 2019/2020 and 1.11-1.59 in 2016/2017 (Table A.7). The proportion of females in the Norwegian bear population also increased over the years (Table A.3) and was likely between 40% and 51% in 2025 compared with 32%-47% in 2016. However, the population sex-ratio is still biased towards males in 2025.

This overall pattern hides variable population trajectories in the different large carnivore regions of Norway (Figure 5). Bear abundance appears to be relatively stable and low in large carnivore regions 3 (Oppland) and 7 (Nordland), and to fluctuate around 30 bears in large carnivore region 6 (Trøndelag, Møre og Romsdal). By contrast, the number of brown bears increased in region 8 (Troms and Finnmark), from 24-41 in 2016 to 56-78 in 2025 (Figure 5), and numbers almost tripled in region 5 (Hedmark) during the past 10 years, from 16-26 individuals in 2016 to 50-68 in 2025. The evolution of the population sex-ratio also varied spatially, with the strongest estimated increase in the proportion of females in region 5 (from 0.20-0.42 in 2016 to 0.33-0.47 in 2025), while the highest proportion of females was found in region 8 with 0.45-0.62 in 2025.

We provide annual abundance estimates for the entire study area, by large carnivore region (Table A.4), and by county (Table A.5) between 2016 and 2025. The analysis also yielded annual density maps that reveal changes in the distribution of bears over time. It especially shows the expansion and increase in density in region 5 (Figure 5, Figure A.5).

Table 1: Bear population size estimates in 2025 by sex and large carnivore management region within the main Norwegian population range. Only management units that are within or that intersect the study area are included in the table. The percentage of the total area of each unit included in the analysis is provided in the last column. Readers should focus on the 95% credible interval provided in parentheses as these - unlike mean values - convey uncertainty inherent in abundance estimates. Numbers are based on estimated AC locations of bears. Combined female-male estimates were obtained by joining sex-specific posterior distributions. Rounding may result in small deviations between total estimates and the sum of the estimates for constituent regions.

	Females	Males	Total	% Area
Region 1	0.1 (0-1)	0.1 (0-1)	0.3 (0-2)	6
Region 2	0.8 (0-3)	1.1 (0-4)	1.9 (0-5)	36
Region 3	0.7 (0-3)	1 (0-3)	1.8 (0-5)	100
Region 4	0.1 (0-1)	0.5 (0-2)	0.5 (0-2)	35
Region 5	23.7 (18-30)	35.5 (29-43)	59.2 (50-68)	100
Region 6	13.8 (11-17)	19 (15-24)	32.8 (28-39)	91
Region 7	0.6 (0-3)	3.6 (1-7)	4.2 (2-8)	100
Region 8	35.7 (28-44)	30.8 (23-39)	66.6 (56-78)	100
TOTAL	75.6 (64-88)	91.7 (79-105)	167.2 (150-186)	73




Figure 2: Total (black) and sex-specific (blue: males, red: females) annual bear population size estimates within the main Norwegian population range between 2016 and 2025 derived from the open-population spatial capture-recapture model. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Yellow bars show the number of individuals detected in Norway each year. The number of individuals detected in Norway may exceed the estimated population size, as individuals with activity centers in neighboring countries, near the Norwegian border, can be detected inside Norway.

Box 2: On the definition of population size

Population size can be defined in different ways and the interpretation of the estimates will depend on the chosen definition. Bear population size in Norway can for example be defined as the total number of bears that spent any time within Norwegian boundaries, i.e., that can potentially be detected in Norway. However, this definition is problematic for different reasons. Most importantly, trans-boundary bears, i.e., bears that spend time on both sides of a border, will potentially be double-counted, which leads to overestimation of the total population size when adding-up estimates from multiple regions (Bischof et al., 2016). In the SCR framework, population size is usually defined as the number of individuals that have their AC within the borders of the region of interest (*AC-based abundance*). Because the AC of an individual cannot be located on both sides of a border, this resolves the issue of double counting. Note that while the AC of an individual cannot be located on both sides of the border, the uncertainty about its exact location and therefore the probability that an individual has its AC located in a given region, are directly integrated in the credible interval (see the **Abundance estimates** section). However, large carnivores have large home ranges and their influence extends beyond their AC location. Because SCR models also estimate average individual space-use, it is possible to derive the proportion of time each individual spends in a given region. From this, population size can be defined as the sum of the individual space use distributions within a given region (*space-use-based abundance* see the **Density maps** section). This definition also avoids double-counting. Contrary to *AC-based abundance*, abundance based on space use does not assign an individual to one of several regions but instead attributes the presence of a single individual to all regions proportionally to its use of that region.

Table 2: Brown bear population size estimates by sex in Norway in 2025 based on individual activity center locations.

	Females	Males	Total
Region 1	0.1 (0-1)	0.1 (0-1)	0.3 (0-2)
Region 2	0.8 (0-3)	1.1 (0-4)	1.9 (0-5)
Region 3	0.7 (0-3)	1 (0-3)	1.8 (0-5)
Region 4	0.1 (0-1)	0.5 (0-2)	0.5 (0-2)
Region 5	23.7 (18-30)	35.5 (29-43)	59.2 (50-68)
Region 6	13.8 (11-17)	19 (15-24)	32.8 (28-39)
Region 7	0.6 (0-3)	3.6 (1-7)	4.2 (2-8)
Region 8	35.7 (28-44)	30.8 (23-39)	66.6 (56-78)
TOTAL	75.6 (64-88)	91.7 (79-105)	167.2 (150-186)

Table 3: Brown bear population size estimates by sex in Norway in 2025 based on individual space use distributions.

	Females	Males	Total
Region 1	0.3 (0-1)	0.2 (0-1)	0.4 (0-2)
Region 2	0.9 (0-2)	1.3 (0-2)	2.1 (0-4)
Region 3	0.5 (0-1)	1.5 (0-4)	2 (0-5)
Region 4	0 (0-0)	0.5 (0-1)	0.5 (0-1)
Region 5	23.9 (20-30)	36.3 (31-42)	60.2 (51-69)
Region 6	14 (13-15)	20.8 (18-23)	34.8 (33-37)
Region 7	0.7 (0-2)	3.7 (2-7)	4.4 (3-8)
Region 8	35.1 (31-40)	31.5 (27-38)	66.6 (61-78)
TOTAL	75.4 (67-83)	95.7 (84-106)	171.1 (158-183)

In the case of the Norwegian bear population, the higher population densities in Sweden imply that many individuals will have their AC estimated to be located in Sweden while their space use distribution will extend into Norway. Consequently, population size estimates for Norway based on individual space-use tend to be slightly higher than estimates based on activity center locations, and especially for males, who use larger areas compared to females (Table A.8).




Figure 3: Brown bear density based on individual space use distributions throughout the study area (white background) in Norway in 2025. Density was estimated with an open-population spatial-capture recapture model fitted to NGS and dead recovery data collected between 2016 and 2025. See the 'Methods' section for further detail.

3.3 Vital rates

The OPSCR model produced annual estimates of legal hunting mortality and mortality associated with all other causes (Figure 4), as well as estimates of per capita recruitment rates over the 10-year period (Table A.6).




Figure 4: Mortality probabilities due to legal culling and all other causes for female and male brown bears. Shown are overall estimates throughout the study area in Norway, based on the sex-specific OPSCR models. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Estimates refer to deaths occurring between the start of one sampling season and the start of the next one. Consequently, estimates for mortality between 2025 and 2026 are not yet available for this analysis.

3.4 Detection probability

Every year, more bears were detected through NGS in Norway than the estimated population size. This is due to the detection of individuals with estimated activity centers located in neighboring countries (Figure 2, see also explanation in Bischof et al. 2016). Specifically, the OPSCR model estimated that between 41 and 59 individuals detected through NGS in Norway in 2025 had their activity center located in either Sweden (16 to 29 individuals), Finland (8 to 19) or Russia (9 to 19), which corresponds to between 23% and 33% of the individuals detected in Norway that year.

When it comes to detectability in the Norwegian bear population, in 2025, the model estimated that between 66% and 80% of the Norwegian bear population were detected through NGS conducted in Norway. Detectability was higher for males compared to females and varied between years, from 0.54-0.74 for females in 2018 to 0.79-0.93 for males in 2021 (Table A.9).

Baseline detection probability (p_0) varied between jurisdictions and was always higher for females compared to males (Figure A.6). Detection probability decreased with the distance to the nearest road (Figure A.4, Table A.8) and strongly increased with the proxy for search effort derived from the observation data in Skandobs and Rovbase (Figure A.3, Table A.8).

4 Discussion

Our analysis revealed a positive population trajectory for the bear population in Norway during the last 10 years (Figure 2). In previous analyses of the Norwegian portion of the Scandinavian brown bear population, we highlighted that the female portion of the population was increasing faster than its male's counterparts, leading to a more balanced population sex-ratio in recent years (Dupont et al., 2023a,b, 2024b, 2025). This pattern, consistent with a bear population in a recolonization phase, was mostly visible between 2012 and 2018 and is less visible in this year's report. Male bears, who move more, disperse more frequently and further than females (Zedrosser et al., 2007; Støen et al., 2006), can recolonize areas at a higher pace, leading to a higher proportion of males on the outskirts of the population (Kindberg et al., 2011). Being philopatric, recolonization of females is slower (Zedrosser et al., 2007; Kindberg et al., 2011; Støen et al., 2006), but their proportion should slowly catch up as population density increases, due in part to the higher estimated survival rate of females compared to males (Figure 4, Table A.6).

The spatially-explicit nature of our analysis also allowed us to highlight regional differences in the brown bear population dynamics in Norway (Figure 5). The overall population increase in the Norwegian brown bear population seems mainly driven by increasing bear numbers in carnivore management region 5 (Hedmark) and 8 (Troms and Finnmark), whereas it remained stable in regions 3 (Oppland), 6 (Møre og Romsdal, Trøndelag), and 7 (Nordland). Note that in this year's report we also provide abundance estimates for regions 1, 2 and 4 (Table A.4) because of the larger study area considered (see below), but mean estimates for these peripheral regions were below 1 across the 10-year study period. The increase in the proportion of females also varied between management regions and was most visible in region 5 and 8. Region 8 is also the only high-density region where the estimated number of females reaches the same levels or even surpasses the estimated number of males in recent years. This may reflect the fact that this region is more advanced in the recolonization process as it is closer to areas of potentially high bear population density in Finland and Russia, compared to regions at the edge of the brown bear range like Oppland (region 3) where the bear population is still small and dominated by males (Table A.3).

5 Summary of improvements made

The analysis described in this report includes the following adjustments compared with the previous analyses of the brown bear monitoring data in Norway by RovQuant (Dupont et al., 2023a,b, 2024b, 2025):

1. Addition of the data from the 2025 monitoring season, and dropping of the data from the 2015 monitoring season.
2. Adjustment of the data filtering process to include NGS and dead recoveries registered as being part of the Norwegian monitoring program only (i.e. exclusion of data from the Swedish monitoring programs, even if samples fell within detector grid cells attributed to Norway).
3. Use of a larger, fixed study area that is not based solely on the buffered spatial distribution of DNA samples. This avoids variation in abundance estimates between annual reports due to changes in the definition of the study area.




Figure 5: Total (black) and sex-specific (blue: males, red: females) annual bear population size estimates in the different carnivore management regions in Norway (Figure A.1) between 2016 and 2025. Abundance estimates were derived from the locations of individual activity centers estimated by sex-specific open-population spatial capture-recapture models. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Here, we only present carnivore regions with a mean estimated population size >= 3 in at least one year during the study period.

6 Suggestions for future improvements

As RovQuant continues to work on improving the functionality and efficiency of OPSCR models, we intend to test and potentially implement the following developments in future analyses of the Scandinavian bear monitoring data:

1. Further explore possibilities to use the NGS data collected in Sweden to derive reliable population size estimates jointly for Sweden and Norway.
2. Review and adjust spatial covariates on density to better reflect habitat selection by bears and therefore population density. This may involve the addition of land cover and topographic variables.
3. Distinguish between dispersing and non-dispersing individuals, for example by using a finite-mixture approach.
4. Consider alternative detection models that do not assume a half-normal shape and/or circular home ranges (Sutherland et al., 2015; Dey et al., 2022a).
5. Account for unknown sources of spatial variation in detectability and density through the use of spatially autocorrelated random effects (Dey et al., 2022b).
6. Account for spatial variation in survival and other vital rates (Milleret et al., 2023).

7 Acknowledgements

This work was made possible by the large carnivore monitoring programs and the extensive monitoring data collected by Norwegian (SNO) and Swedish (Länsstyrelserna) wildlife management authorities, as well as the public. Our analyses relied on genetic analyses conducted by the laboratory personnel at the DNA laboratories at NIBIO Svanhovd, the Swedish Museum of Natural History (NRM), and the Norwegian Institute for Nature Research (NINA). We also thank Swedish and Norwegian wildlife managers for feedback provided during project RovQuant and the Research Council of Norway for partial funding (NFR 345279; project PopFlow). The computations/simulations were performed on resources provided by NMBU's computing cluster "Orion", administered by the Centre for Integrative Genetics. J. Vermaat provided helpful comments on drafts of this report.

8 Data availability

Data, R code to reproduce the analysis, as well as figures, tables, and raster maps (Figure A.5) are available on GitHub (https://github.com/richbi/RovQuantPublic)

References

Andreassen, R., Schregel, J., Kopatz, A., Tobiassen, C., Knappskog, P., Hagen, S., Kleven, O., Schneider, M., Kojola, I., Aspi, J., et al. (2012). A forensic dna profiling system for northern european brown bears (ursus arctos). *Forensic Science International: Genetics*, 6(6):798–809.

Bidon, T., Frosch, C., Eiken, H. G., Kutschera, V. E., Hagen, S. B., Aarnes, S. G., Fain, S. R., Janke, A., and Hailer, F. (2013). A sensitive and specific multiplex pcr approach for sex identification of ursine and tremarctine bears suitable for non-invasive samples. *Molecular Ecology Resources*, 13(3):362–368.

Bischof, R., Brøseth, H., and Gimenez, O. (2016). Wildlife in a politically divided world: Insularism inflates estimates of brown bear abundance. *Conservation Letters*, 9(2):122–130.

Bischof, R., Dupont, P., Milleret, C., Chipperfield, J., and Royle, J. A. (2020a). Consequences of ignoring group association in spatial capture–recapture analysis. *Wildlife Biology*, 2020(1).

Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Åkesson, M., Brøseth, H., and Kindberg, J. (2019a). Estimating the size of the scandinavian wolf population with spatial capture recapture and conversion factors. *MINA fagrapport 57*, 80pp.

Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Brøseth, H., and Kindberg, J. (2019b). Estimating density, abundance and population dynamics of bears, wolverines, and wolvesin scandinavia. *MINA fagrapport 63*, 79pp.

Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Tourani, M., Ordiz, A., de Valpine, P., Turek, D., Royle, J. A., Gimenez, O., Flagstad, Ø., Åkesson, M., Svensson, L., Brøseth, H., and Kindberg, J. (2020b). Estimating and forecasting spatial population dynamics of apex predators using transnational genetic monitoring. *Proceedings of the National Academy of Sciences*, 117(48):30531–30538.

Bischof, R., Turek, D., Milleret, C., Ergon, T., Dupont, P., Dey, S., and de Valpine, P. (2021). *nimbleSCR: Spatial Capture-Recapture (SCR) Methods Using 'nimble'*. R package version 0.1.2.

Brøseth, H., Kleven, O., Jensen, M. Ø., and Kopatz, A. (2026). Dna-basert overvåking av brunbjørn i norge i 2025. *NINA Rapport 2760*.

Chandler, R. B., Hepinstall-Cymerman, J., Merker, S., Abernathy-Conners, H., and Cooper, R. J. (2018). Characterizing spatio-temporal variation in survival and recruitment with integrated population models. *The Auk: Ornithological Advances*, 135(3):409–426.

de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., and Bodik, R. (2017). Programming with models: writing statistical algorithms for general model structures with nimble. *Journal of Computational and Graphical Statistics*, 26(2):403–413.

Dey, S., Bischof, R., Dupont, P. P. A., and Milleret, C. (2022a). Does the punishment fit the crime? consequences and diagnosis of misspecified detection functions in bayesian spatial capture–recapture modeling. *Ecology and Evolution*, 12(2):e8600.

Dey, S., Moqanaki, E. M., Milleret, C., Dupont, P., Tourani, M., and Bischof, R. (2022b). Modelling spatially autocorrelated detection probabilities in spatial capture-recapture using random effects. *bioRxiv*, pages 2022–05.

Dupont, P., Milleret, C., Brøseth, H., Kindberg, J., and Bischof, R. (2023a). Estimates of brown bear density, abundance, and population dynamics in Norway 2012-2021. *MINA fagrapport 82*, 32pp.

Dupont, P., Milleret, C., Brøseth, H., Kindberg, J., and Bischof, R. (2023b). Estimates of brown bear density, abundance, and population dynamics in Norway 2012-2022. *MINA fagrapport 86*, 33pp.

Dupont, P., Milleret, C., Brøseth, H., Kindberg, J., and Bischof, R. (2024a). Challenges in estimating range-wide brown bear density and abundance in sweden (2012-2021). *MINA fagrapport 95*, 47pp.

Dupont, P., Milleret, C., Brøseth, H., Kindberg, J., and Bischof, R. (2024b). Estimates of brown bear density, abundance, and population dynamics in Norway 2014-2023. *MINA fagrapport 98*, 34pp.

Dupont, P., Milleret, C., Semper-Pascual, A., Brøseth, H., Flagstad, Ø., Kindberg, J., Svensson, L., and Bischof, R. (2026). Estimates of wolf density, abundance, and population dynamics in Sweden and Norway, 2016/17–2025/26. *MINA fagrapport 113*, 37pp.

Dupont, P., Milleret, C., Semper-Pascual, A., Brøseth, H., Kopatz, A., Kindberg, J., and Bischof, R. (2025). Estimates of brown bear density, abundance, and population dynamics in Norway 2015-2024. *MINA fagrapport 105*, 35pp.

Dupont, P., Milleret, C., Tourani, M., Brøseth, H., and Bischof, R. (2021). Integrating dead recoveries in open-population spatial capture–recapture models. *Ecosphere*, 12(7):e03571.

Efford, M. (2004). Density estimation in live-trapping studies. *Oikos*, 106(3):598–610.

Efford, M. G. and Schofield, M. R. (2022). A review of movement models in open population capture–recapture. *Methods in Ecology and Evolution*, 13(10):2106–2118.

Ergon, T. and Gardner, B. (2014). Separating mortality and emigration: modelling space use, dispersal and survival with robust-design spatial capture–recapture data. *Methods in Ecology and Evolution*, 5(12):1327–1336.

Gardner, B., Sollmann, R., Kumar, N. S., Jathanna, D., and Karanth, K. U. (2018). State space and movement specification in open population spatial capture-recapture models. *Ecology and Evolution*, 8(20):10336–10344.

Gelman, A. and Rubin, D. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7:457–511.

Kéry, M. and Schaub, M. (2011). *Bayesian population analysis using WinBUGS: a hierarchical perspective*. Academic Press.

Kindberg, J., Swenson, J. E., Ericsson, G., Bellemain, E., Miquel, C., and Taberlet, P. (2011). Estimating population size and trends of the swedish brown bear ursus arctos population. *Wildlife Biology*, 17(2):114–123.

Lebreton, J.-D. and Pradel, R. (2002). Multistate recapture models: modelling incomplete individual histories. *Journal of Applied Statistics*, 29(1-4):353–369.

Milleret, C., Dey, S., Dupont, P., Brøseth, H., Turek, D., de Valpine, P., and Bischof, R. (2023). Estimating spatially variable and density-dependent survival using open-population spatial capture–recapture models. *Ecology*, 104(2):e3934.

Milleret, C., Dupont, Pierre Semper-Pascual, A., Hansen, N. R., Brøseth, H., Flagstad, Ø., Hedmark, E., Kleven, O., Königsson, H., Spong, G., Kindberg, J., and Bischof, R. (2026). Estimates of wolverine density, abundance, and population dynamics in Scandinavia, 2015/16–2024/25. *MINA fagrapport 112*, 34pp.

Milleret, C., Dupont, P., Åsbrink, J., and Bischof, R. (2024). Simultaneous range-wide genetic sampling of brown bears in Sweden: a pilot study. *MINA fagrapport 96*, 23pp.

Milleret, C., Dupont, P., Moqanaki, E., Brøseth, H., Flagstad, Ø., Kleven, O., Kindberg, J., and Bischof, R. (2022). Estimates of wolverine density, abundance, and population dynamics in Scandinavia, 2014–2022. *MINA fagrapport 79*, 35pp.

NIMBLE Development Team (2019). *NIMBLE: MCMC, Particle Filtering, and Programmable Hierarchical Modeling*. https://cran.r-project.org/package=nimble.

Norman, A. J. and Spong, G. (2015). Single nucleotide polymorphism-based dispersal estimates using noninvasive sampling. *Ecology and Evolution*, 5(15):3056–3065.

Norman, A. J., Street, N. R., and Spong, G. (2013). De novo snp discovery in the scandinavian brown bear (ursus arctos). *PLoS One*, 8(11):e81012.

Paetkau, D. and Strobeck, C. (1998). Ecological genetic studies of bears using microsatellite analysis. *Ursus*, pages 299–306.

R Core Team (2022). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Royle, J. A., Chandler, R. B., Sollmann, R., and Gardner, B. (2014). *Spatial Capture-Recapture*. Academic Press.

Royle, J. A. and Dorazio, R. M. (2012). Parameter-expanded data augmentation for bayesian analysis of capture–recapture models. *Journal of Ornithology*, 152(2):521–537.

Schaub, M. and Royle, J. A. (2014). Estimating true instead of apparent survival using spatial cormack-jolly-seber models. *Methods in Ecology and Evolution*, 5(12):1316–1326.

Støen, O.-G., Zedrosser, A., Sæbø, S., and Swenson, J. E. (2006). Inversely density-dependent natal dispersal in brown bears ursus arctos. *Oecologia*, 148:356–364.

Sutherland, C., Fuller, A. K., and Royle, J. A. (2015). Modelling non-euclidean movement and landscape connectivity in highly structured ecological networks. *Methods in Ecology and Evolution*, 6(2):169–177.

Taberlet, P., Griffin, S., Goossens, B., Questiau, S., Manceau, V., Escaravage, N., Waits, L. P., and Bouvet, J. (1996). Reliable genotyping of samples with very low dna quantities using pcr. *Nucleic acids research*, 24(16):3189–3194.

Turek, D., Milleret, C., Ergon, T., Brøseth, H., Dupont, P., Bischof, R., and De Valpine, P. (2021). Efficient estimation of large-scale spatial capture–recapture models. *Ecosphere*, 12(2):e03385.

Zedrosser, A., Støen, O.-G., Sæbø, S., and Swenson, J. E. (2007). Should i stay or should i go? natal dispersal in the brown bear. *Animal Behaviour*, 74(3):369–376.

Zhang, W., Chipperfield, J. D., Illian, J. B., Dupont, P., Milleret, C., de Valpine, P., and Bischof, R. (2022). A flexible and efficient bayesian implementation of point process models for spatial capture-recapture data. *Ecology*, page e3887.

Appendices




Figure A.1: Large carnivore management regions (*"forvaltningsregioner for rovvilt"*; in shades of red, from 1 to 8) and counties (*"fylker"*; black outlines and labels) in Norway. Note that Innlandet county is now composed of former counties Oppland (region 3) and Hedmark (region 5).




Figure A.2: Spatial covariates denoting the scaled density of brown bear dead recoveries (left) and brown bear observations recorded in Skandobs and Rovabse (right) in Norway between 2016 and 2025. These covariates were used in the density and movement sub-models of the open-population spatial capture-recapture model.




Figure A.3: Spatial covariate denoting the presence (green cells) or absence (white cells) of at least one carnivore observation between 2016 and 2025 within each 5 x 5km detector grid cell. This covariate was used in the detection sub-model of the open-population spatial capture-recapture model.




Figure A.4: Spatial covariate denoting the average distance to the nearest road in Norway for each 5 x 5km detector grid cell. This covariate was used in the detection sub-model of the open-population spatial capture-recapture model.




Figure A.5: Annual brown bear density based on individual space-use throughout Norway between 2016 and 2025. Density was estimated using the open-population spatial capture-recapture model.




Figure A.6: Sex-specific baseline detection probability (p_0) estimated by the open-population spatial capture recapture models (females in red and males in blue). Results are separated into panels based on regions (each combining multiple counties). Estimates are shown for the mean values of the detection covariates. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Note that baseline detection probability (p_0) is a theoretical value of detection probability when a detector coincides with the location of an individual's activity center; it is not to be confused with detectability, i.e, the overall probability of detecting an individual.

Table A.1: Annual number of brown bear non-invasive genetic samples collected and individuals detected in Norway and included in the OPSCR analyses for females (F) and males (M). In the analysis, we included samples collected within the study area during the primary monitoring period (Apr 1 - Nov 30) between 2016 and 2025.

	2016		2017		2018		2019		2020		2021		2022		2023		2024		2025	
	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
number of NGS samples	205	291	186	273	215	213	240	360	268	445	289	471	365	400	444	452	285	389	307	456
number of NGS individuals	51	74	54	67	61	69	55	86	72	84	67	90	80	97	78	98	86	102	77	102

Table A.2: Number of cause-specific dead recoveries of bears included in the OPSCR analysis between 2016 and 2025. Numbers are reported by country, for females (F) and males (M). Note that, i) contrary to previous reports (Dupont et al., 2025), bears recovered dead in Sweden inside the 50 km habitat buffer were *not* included in this year's analysis (see the *Summary of improvements made* section), and ii) dead recovery data from the final season (2025) were not used in this analysis because survival is modeled as part of the transition from one season to the next.

	Country	2016		2017		2018		2019		2020		2021		2022		2023		2024		2025	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
Other	Norway	0	0	0	1	1	1	2	0	1	4	0	0	2	1	0	0	0	1	1	0
	Sweden	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Legal culling	Norway	0	8	0	8	2	8	2	11	1	4	2	10	1	10	1	7	3	9	4	14
	Sweden	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL	Total	0	8	0	9	3	9	4	11	2	8	2	10	3	11	1	7	3	10	5	14

Table A.3: Annual proportion of females in the brown bear population of Norway. Estimates were derived using the posterior distributions of sex-specific annual abundance estimates. Grey font is used for large carnivore regions with a mean estimated population size <= 5 in at least one year, to emphasize the small sample size (number of individuals) associated with these sex ratio estimates. Credible intervals (95%) are shown in parentheses.

	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025
Region 1	0.35 (0.00-1.00)	0.32 (0.00-1.00)	0.39 (0.00-1.00)	0.42 (0.00-1.00)	0.43 (0.00-1.00)	0.41 (0.00-1.00)	0.33 (0.00-1.00)	0.35 (0.00-1.00)	0.46 (0.00-1.00)	0.50 (0.00-1.00)
Region 2	0.34 (0.00-1.00)	0.29 (0.00-1.00)	0.34 (0.00-1.00)	0.37 (0.00-1.00)	0.37 (0.00-1.00)	0.33 (0.00-1.00)	0.26 (0.00-1.00)	0.22 (0.00-1.00)	0.42 (0.00-1.00)	0.41 (0.00-1.00)
Region 3	0.02 (0.00-0.33)	0.05 (0.00-0.50)	0.14 (0.00-1.00)	0.08 (0.00-0.50)	0.10 (0.00-1.00)	0.09 (0.00-1.00)	0.02 (0.00-0.50)	0.04 (0.00-0.50)	0.09 (0.00-0.50)	0.42 (0.00-1.00)
Region 4	0.08 (0.00-1.00)	0.08 (0.00-1.00)	0.20 (0.00-1.00)	0.20 (0.00-1.00)	0.25 (0.00-1.00)	0.03 (0.00-1.00)	0.04 (0.00-1.00)	0.01 (0.00-0.00)	0.12 (0.00-1.00)	0.09 (0.00-1.00)
Region 5	0.30 (0.20-0.42)	0.36 (0.26-0.45)	0.45 (0.33-0.56)	0.42 (0.32-0.52)	0.40 (0.32-0.48)	0.45 (0.37-0.53)	0.40 (0.32-0.48)	0.43 (0.37-0.49)	0.44 (0.39-0.50)	0.40 (0.33-0.47)
Region 6	0.38 (0.27-0.48)	0.34 (0.25-0.44)	0.43 (0.34-0.53)	0.47 (0.40-0.55)	0.55 (0.47-0.62)	0.37 (0.31-0.44)	0.35 (0.28-0.42)	0.35 (0.27-0.44)	0.37 (0.30-0.44)	0.42 (0.35-0.50)
Region 7	0.22 (0.00-0.67)	0.41 (0.17-1.00)	0.36 (0.14-0.67)	0.20 (0.00-0.40)	0.03 (0.00-0.25)	0.02 (0.00-0.33)	0.02 (0.00-0.25)	0.03 (0.00-0.33)	0.05 (0.00-0.33)	0.13 (0.00-0.50)
Region 8	0.51 (0.38-0.64)	0.46 (0.35-0.58)	0.45 (0.36-0.54)	0.49 (0.42-0.57)	0.49 (0.40-0.58)	0.50 (0.41-0.58)	0.52 (0.42-0.62)	0.49 (0.40-0.58)	0.58 (0.50-0.67)	0.54 (0.45-0.62)
TOTAL	0.39 (0.32-0.47)	0.38 (0.32-0.45)	0.44 (0.38-0.50)	0.45 (0.40-0.50)	0.45 (0.40-0.50)	0.44 (0.39-0.49)	0.41 (0.36-0.47)	0.42 (0.37-0.47)	0.47 (0.42-0.51)	0.45 (0.40-0.51)

Table A.4: Annual abundance estimates for Norway and by large carnivore management regions (Figure A.1). Estimates are based on activity center locations estimated by the open-population spatial capture-recapture (OPSCR) model. Credible intervals (95%) are shown in parentheses. Small deviations between the total estimate and the sum of abundance estimates from the constituent subregions may arise due to rounding. See Methods section for further detail.

	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025
Region 1	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.3 (0-2)
Region 2	0.5 (0-2)	0.8 (0-3)	0.8 (0-3)	0.8 (0-3)	0.5 (0-2)	0.4 (0-2)	0.4 (0-2)	0.6 (0-2)	0.8 (0-3)	1.9 (0-5)
Region 3	3 (1-6)	2.5 (0-5)	1.4 (0-4)	2.1 (0-5)	0.9 (0-3)	0.5 (0-2)	1.7 (1-4)	1.4 (0-3)	2 (1-4)	1.8 (0-5)
Region 4	0 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0 (0-1)	0.1 (0-1)	0 (0-1)	0.1 (0-1)	0.1 (0-1)	0.5 (0-2)
Region 5	20.5 (16-26)	32.3 (26-39)	30.8 (24-38)	36.7 (29-44)	43.1 (37-50)	40.4 (34-47)	45.8 (39-53)	55.2 (48-62)	60.1 (53-67)	59.2 (50-68)
Region 6	24.6 (19-30)	31 (25-38)	33.5 (28-40)	31.7 (27-37)	28 (24-32)	25.5 (22-29)	27.7 (24-32)	27.2 (21-34)	31.3 (26-37)	32.8 (28-39)
Region 7	2.9 (1-6)	3.4 (1-6)	3.8 (2-6)	5.8 (4-8)	3.3 (3-5)	2.3 (1-4)	3.5 (2-6)	3.8 (1-7)	4.2 (3-7)	4.2 (2-8)
Region 8	31.9 (24-41)	40.2 (31-50)	46.1 (38-55)	51.8 (44-60)	41.5 (34-49)	41.7 (35-49)	42.9 (35-52)	50.6 (42-60)	61.6 (52-72)	66.6 (56-78)
TOTAL	83.5 (71-96)	110.3 (97-125)	116.6 (103-131)	129 (117-142)	117.4 (106-129)	111 (100-122)	122 (109-135)	139 (125-154)	160.3 (146-175)	167.2 (150-186)

Table A.5: Annual abundance estimates for Norway and by county (Figure A.1). Innlandet county encompasses the former counties Oppland (region 3) and Hedmark (region 5). Estimates are based on activity center locations estimated by the open-population spatial capture-recapture (OPSCR) model. Credible intervals (95%) are shown in parentheses. Small deviations between the total estimate and the sum of abundance estimates from the constituent subregions may arise due to rounding. See Methods section for further detail.

	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025
Akershus	0 (0-1)	0 (0-1)	0.1 (0-1)	0.1 (0-1)	0 (0-1)	0.1 (0-1)	0 (0-1)	0.1 (0-1)	0.1 (0-1)	0.5 (0-2)
Buskerud	0.3 (0-2)	0.5 (0-2)	0.5 (0-2)	0.6 (0-2)	0.4 (0-2)	0.3 (0-2)	0.3 (0-2)	0.5 (0-2)	0.6 (0-3)	1.6 (0-5)
Finnmark	21.3 (14-29)	29.7 (22-38)	33.8 (26-42)	43.9 (37-51)	34 (27-42)	33.7 (27-41)	39.1 (31-47)	47.2 (39-56)	54.9 (45-65)	56.6 (46-68)
Innlandet	23.5 (19-29)	34.8 (29-41)	32.2 (25-40)	38.7 (31-47)	44 (37-51)	40.9 (34-48)	47.5 (40-55)	56.6 (50-64)	62.1 (55-70)	61 (52-71)
Møre og Romsdal	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0 (0-1)	0 (0-1)	0.1 (0-1)	0.1 (0-1)	0.4 (0-2)
Nordland	2.9 (1-6)	3.4 (1-6)	3.8 (2-6)	5.8 (4-8)	3.3 (3-5)	2.3 (1-4)	3.5 (2-6)	3.8 (1-7)	4.2 (3-7)	4.2 (2-8)
Oslo	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Telemark	0.2 (0-1)	0.2 (0-1)	0.2 (0-1)	0.2 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.2 (0-1)	0.3 (0-2)
Troms	10.6 (8-13)	10.5 (8-14)	12.3 (11-15)	7.9 (6-10)	7.5 (7-9)	8 (7-9)	3.8 (3-6)	3.4 (2-6)	6.7 (5-9)	10 (8-13)
Trøndelag	24.5 (19-30)	30.9 (25-38)	33.4 (28-40)	31.6 (27-37)	27.9 (24-32)	25.4 (22-29)	27.6 (23-32)	27.1 (21-34)	31.1 (26-37)	32.4 (28-38)
Vestland	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)	0.3 (0-2)
TOTAL	83.5 (71-96)	110.3 (97-125)	116.6 (103-131)	129 (117-142)	117.4 (106-129)	111 (100-122)	122 (109-135)	139 (125-154)	160.3 (146-175)	167.2 (150-186)

Table A.6: Estimates of the demographic parameters obtained from the bear OPSCR model and data collected in Norway between 2016 and 2025. Median estimates and 95% credible intervals (in parentheses) for per capita recruitment rate (ρ), survival (ϕ), mortality due to legal culling (h), mortality due to other causes (w), and proportion of individuals dead to other causes recovered (r), are presented for males (M) and females (F). Note that survival and mortality do not always sum to 1 because the posterior median is presented (not the mean).

	2016 to 2017		2017 to 2018		2018 to 2019		2019 to 2020		2020 to 2021	
	F	M	F	M	F	M	F	M	F	M
ρ	0.15 (0.03-0.37)	0.33 (0.14-0.56)	0.19 (0.07-0.35)	0.20 (0.06-0.38)	0.19 (0.08-0.34)	0.20 (0.08-0.34)	0.08 (0.01-0.20)	0.19 (0.08-0.31)	0.10 (0.03-0.20)	0.18 (0.09-0.28)
ϕ	0.97 (0.88-1.00)	0.86 (0.75-0.94)	0.87 (0.77-0.95)	0.75 (0.65-0.84)	0.85 (0.75-0.92)	0.79 (0.69-0.86)	0.86 (0.76-0.93)	0.73 (0.63-0.81)	0.86 (0.77-0.93)	0.76 (0.66-0.84)
h	0.01 (0.00-0.03)	0.06 (0.03-0.11)	0.00 (0.00-0.02)	0.05 (0.02-0.09)	0.02 (0.00-0.04)	0.05 (0.02-0.09)	0.02 (0.00-0.04)	0.07 (0.04-0.11)	0.01 (0.00-0.03)	0.03 (0.01-0.07)
w	0.02 (0.00-0.11)	0.08 (0.01-0.18)	0.12 (0.04-0.23)	0.20 (0.11-0.30)	0.13 (0.06-0.23)	0.16 (0.09-0.25)	0.12 (0.05-0.22)	0.20 (0.12-0.29)	0.13 (0.06-0.22)	0.21 (0.13-0.30)
r	0.21 (0.01-0.92)	0.06 (0.00-0.40)	0.04 (0.00-0.21)	0.05 (0.01-0.16)	0.08 (0.01-0.26)	0.06 (0.01-0.20)	0.13 (0.03-0.36)	0.02 (0.00-0.10)	0.08 (0.01-0.26)	0.14 (0.05-0.30)

	2021 to 2022		2022 to 2023		2023 to 2024		2024 to 2025	
	F	M	F	M	F	M	F	M
ρ	0.05 (0.00-0.14)	0.39 (0.26-0.53)	0.16 (0.09-0.24)	0.28 (0.17-0.43)	0.16 (0.08-0.27)	0.24 (0.13-0.38)	0.18 (0.09-0.30)	0.31 (0.18-0.46)
ϕ	0.95 (0.88-0.98)	0.75 (0.65-0.83)	0.87 (0.79-0.94)	0.81 (0.71-0.89)	0.94 (0.86-0.98)	0.72 (0.62-0.80)	0.77 (0.64-0.88)	0.77 (0.66-0.86)
h	0.02 (0.00-0.05)	0.07 (0.04-0.12)	0.01 (0.00-0.03)	0.06 (0.03-0.11)	0.01 (0.00-0.03)	0.04 (0.02-0.08)	0.02 (0.01-0.05)	0.05 (0.03-0.10)
w	0.03 (0.00-0.10)	0.18 (0.10-0.27)	0.11 (0.05-0.20)	0.12 (0.05-0.22)	0.05 (0.01-0.13)	0.24 (0.16-0.34)	0.21 (0.10-0.33)	0.17 (0.09-0.28)
r	0.12 (0.00-0.68)	0.03 (0.00-0.14)	0.15 (0.03-0.39)	0.08 (0.01-0.27)	0.08 (0.00-0.44)	0.02 (0.00-0.08)	0.02 (0.00-0.11)	0.05 (0.01-0.19)

Table A.7: Annual population growth rate estimates for the bear population in Norway. Estimates were derived using the posterior distributions of annual abundance estimates (Table A.4). Credible intervals (95%) are shown in parentheses.

	2016 to 2017	2017 to 2018	2018 to 2019	2019 to 2020	2020 to 2021	2021 to 2022	2022 to 2023	2023 to 2024	2024 to 2025
λ	1.33 (1.11-1.59)	1.06 (0.90-1.25)	1.11 (0.96-1.28)	0.91 (0.80-1.03)	0.95 (0.83-1.07)	1.10 (0.96-1.26)	1.14 (0.99-1.31)	1.16 (1.01-1.31)	1.05 (0.92-1.19)

Table A.8: Parameter estimates for the spatial and detection processes of the OPSCR models for male (M) and female (F) bears in Norway. The inter-annual movement parameter (τ) and the scale parameter of the detection function (σ) are expressed in kilometers. β_{dead_1} and β_{dead_2} correspond to the effect of the locations of all bears recovered dead throughout the 2016-2025 period on the initial and subsequent AC locations, respectively. $\beta_{skandobs_1}$ and $\beta_{skandobs_2}$ correspond to the effect of the presence of bear observations registered in SkandObs on the initial and subsequent AC locations, respectively (Bischof et al., 2020b). β_{roads} corresponds to the effect of the average distance to the nearest road and β_{obs} corresponds to the effect of the presence of any large carnivore observation registered in Rovbase or SkandObs on the baseline detection probability. All coefficients are associated with scaled covariates. Credible intervals (95%) are shown in parentheses.

	Parameters	**F**	**M**
Spatial process	τ	9.36 (8.54-10.25)	33.24 (31.28-35.41)
	β_{dead_1}	0.48 (0.37-0.61)	0.42 (0.29-0.60)
	$\beta_{skandobs_1}$	0.00 (-0.17-0.14)	0.10 (-0.10-0.28)
	β_{dead_2}	-0.10 (-0.42-0.19)	0.18 (0.10-0.27)
	$\beta_{skandobs_2}$	1.44 (0.99-1.93)	0.78 (0.63-0.94)
Detection process	σ	8.55 (8.26-8.87)	11.80 (11.44-12.18)
	β_{roads}	-0.78 (-0.89-(-)0.67)	-0.77 (-0.87-(-)0.67)
	β_{obs}	1.68 (1.44-1.94)	1.91 (1.72-2.10)

Table A.9: Average proportion of the Norwegian bear population detected via non-invasive genetic sampling (NGS) in Norway. Values were calculated as the proportion of individuals with their activity center estimated in Norway by the OPSCR models (Table A.4) that were detected with NGS (Table A.1). Credible intervals (95%) are shown in parentheses.

	2016	2017	2018	2019	2020
F	0.67 (0.53-0.81)	0.65 (0.54-0.76)	0.64 (0.54-0.74)	0.64 (0.55-0.73)	0.78 (0.69-0.88)
M	0.77 (0.67-0.88)	0.69 (0.59-0.79)	0.69 (0.60-0.79)	0.80 (0.71-0.88)	0.85 (0.77-0.93)
Total	0.73 (0.64-0.81)	0.67 (0.60-0.75)	0.67 (0.60-0.74)	0.72 (0.66-0.79)	0.82 (0.76-0.88)

	2021	2022	2023	2024	2025
F	0.75 (0.66-0.84)	0.74 (0.65-0.83)	0.74 (0.66-0.83)	0.75 (0.66-0.83)	0.68 (0.58-0.78)
M	0.87 (0.79-0.93)	0.83 (0.76-0.91)	0.76 (0.67-0.85)	0.80 (0.72-0.87)	0.77 (0.69-0.86)
Total	0.81 (0.75-0.87)	0.79 (0.73-0.85)	0.75 (0.69-0.81)	0.77 (0.72-0.83)	0.73 (0.66-0.80)