Norwegian University of Life Sciences
Faculty of Environmental Sciences and Natural Resource Management

2026
ISSN 2535-2806

MINA fagrapport 117

Preliminary estimates of wolverine density and abundance in Sweden, 2025/2026

Asunción Semper-Pascual
Pierre Dupont
Cyril Milleret
Eva Hedmark
Carlos Cardoso Palacios
Göran Spong
Helena Königsson
Richard Bischof

Semper-Pascual, A., Dupont, P., Milleret, C., Hedmark, E., Cardoso Palacios, C., Spong, G., Königsson, H., and Bischof, R., 2026. **Preliminary estimates of wolverine density and abundance in Sweden, 2025/2026** - MINA fagrapport 117. 18 pp.

Ås, October 2026

ISSN: 2535-2806

COPYRIGHT
© Norwegian University of Life Sciences (NMBU)
The publication may be freely cited where the source is acknowledged

AVAILABILITY
Open

PUBLICATION TYPE
Digital document (pdf)

QUALITY CONTROLLED BY
The Research committee (FU), MINA, NMBU

PRINCIPAL
Naturvårdsverket, Ref: NV-02444-23 , Contact person: Robert Ekblom

COVER PICTURE

NYCKELORD
Gulo gulo, järv, populationstäthet, detekteringssannolikhet, icke-invasiv genetisk provtagning, rumslig fångst–återfångst, rovdjursförvaltning

KEY WORDS
Gulo gulo, wolverine, population density, detection probability, non-invasive genetic sampling, spatial capture-recapture, carnivore management

Asunción Semper-Pascual, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway

Pierre Dupont, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway

Cyril Milleret, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway

Eva Hedmark, Department of Ecology, Grimsö Wildlife Research Station, Swedish University of Agricultural Sciences, Sweden

Carlos Cardoso Palacios, Department of Ecology, Grimsö Wildlife Research Station, Swedish University of Agricultural Sciences, Sweden

Göran Spong, Department of Wildlife, Fish, and Environmental Studies, Swedish University of Agricultural Sciences, Sweden

Helena Königsson, Department of Wildlife, Fish, and Environmental Studies, Swedish University of Agricultural Sciences, Sweden


Richard Bischof (richard.bischof@nmbu.no), Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway

Summary

Background Project RovQuant has produced density maps and abundance estimates for large carnivores (wolf, wolverine, and brown bear) throughout Scandinavia since 2019 based on non-invasive genetic sampling (NGS) and dead recovery data collected annually by Swedish and Norwegian authorities. In 2026, the Swedish Environmental Protection Agency (Naturvårdsverket) has requested preliminary wolverine abundance estimates for the Swedish portion of the wolverine population during winter 2025/2026, prior to the combined estimates provided annually for Sweden and Norway.

Approach Using NGS data and a single-season spatial capture-recapture (SCR) model, we estimated and mapped the density of wolverines in Sweden during the 2025/2026 winter. The method produced a total wolverine estimate for Sweden and separate estimates for each Swedish county in the current range of the species.

Results and Discussion The total number of wolverines in Sweden during winter 2025/2026 was likely (95% credible interval) between 740 and 802 individuals. We estimated that the proportion of females in the population was likely between 56% and 59%. The highest wolverine abundance was estimated in northern counties: Jämtland (202-232 individuals), Norrrbotten (193-225), and Västerbotten (136-162).




Density map and estimated wolverine abundance range in 2025/2026

Sammanfattning

Bakgrund Projekt RovQuant har sedan 2019 tagit fram täthetskartor och populationsuppskattningar för stora rovdjur (varg, järv och brunbjörn) i Skandinavien, baserat på icke-invasiv genetisk provtagning (NGS) och data från döda djur som samlas in årligen av svenska och norska myndigheter. År 2026 har Naturvårdsverket begärt preliminära populationsuppskattningar för järv inom den svenska delen av järvpopulationen under vintern 2025/2026, innan de samlade uppskattningarna för Sverige och Norge presenteras.

Metod Med hjälp av NGS-data och en single-season rumslig fångst–återfångstmodell (SCR) uppskattade och kartlade vi tätheten av järv i Sverige under vintern 2025/2026. Metoden gav uppskattningar av järvstammens storlek för hela Sverige samt för de svenska län som ingår i artens aktuella utbredningsområde.

Resultat och Diskussion Det totala antalet järvar i Sverige under vintern 2025/2026 uppskattades (95% konfidensintervall) till mellan 740 och 802 individer. Vi beräknade att andelen honor i populationen sannolikt låg mellan 56% och 59%. Den högsta tätheten av järv uppskattades i de nordliga länen Jämtland (202–232 individer), Norrbotten (193–225) och Västerbotten (136–162).



Täthetskarta och beräknad järvtäthet i Sverige under vintern 2025/26

Contents

1 Introduction **6**

2 Methods **8**
2.1 Data 8
2.2 Spatial capture-recapture model 9

3 Results **14**
3.1 Non-invasive genetic samples and dead recoveries 14
3.2 Density and abundance 14

4 Discussion **16**

5 Acknowledgments **16**

6 Data availability **16**

References **18**

1 Introduction

Sweden and Norway monitor large carnivores using non-invasive genetic sampling (NGS) and dead recoveries. Both countries have collected an extensive individual-based data set for the wolverine (*Gulo gulo*), which is stored in the Scandinavian large carnivore database Rovbase (www.rovbase.se, www.rovbase.no). Since 2017, project RovQuant (nmbu.no) has been developing statistical methods that allow a comprehensive assessment of the status and dynamics of large carnivore populations using NGS data and other sources of information stored in Rovbase (Bischof et al., 2019b, 2020). The analytical framework developed by RovQuant is based on spatial capture-recapture (SCR) models (Efford, 2004; Royle et al., 2009). These models use the spatial information contained in the repeated genetic detections of individuals to estimate various population parameters, including spatially-explicit abundance (i.e., density). Importantly, the approach accounts for imperfect detection during sampling (i.e., the fact that some individuals are not detected at all) and animal movement (i.e., the fact that individuals may use and be detected in multiple counties or countries). The SCR method brings along several advantages, including the ability to map density, derive county-specific abundance, and yield tractable measures of uncertainty (Bischof et al., 2019a, 2020).

RovQuant publishes abundance estimates for wolves, wolverines, and bears for different geographic units annually. Joint estimates for wolves and wolverines in Norway and Sweden have been available since 2019 (last reports: Milleret et al. 2024a, 2025b,a; Dupont et al. 2026b); estimates for bears in Norway since 2023 (last reports: Dupont et al. 2025, 2026a); and in 2024 estimates for wolverines in administrative entities associated with the Sámi reindeer herding areas in Norrbotten County (Sweden) were reported (Milleret et al., 2024b). Since 2025, the Swedish Environmental Protection Agency (SEPA, Naturvårdsverket) has requested preliminary wolverine abundance estimates for the Swedish portion of the wolverine population (Semper-Pascual et al., 2025), prior to the combined estimates provided annually for Sweden and Norway.

Here, as requested by the SEPA, we provide preliminary wolverine abundance estimates for Sweden during the winter of 2025/2026 using a single-season SCR model. We provide the following information:

- Sex-specific estimates of the number of wolverines in 2025/2026 in Sweden.
- Sex-specific estimates of the number of wolverines in 2025/2026 for each Swedish county intersecting the species range.
- Map of wolverine density for Sweden.
- Estimated proportion of individuals detected through non-invasive genetic sampling in Sweden.

All estimates are accompanied by their 95% Bayesian credible intervals as a measure of uncertainty. All estimates presented in this report are preliminary and will be replaced by those generated from the final analysis scheduled for completion at the end of 2026, following the completion of NGS data processing and quality control in Norway.

Box 1: Terms and acronyms used

AC: Activity center. Model-based equivalent of the center of an individual's home range during the monitoring period. "AC location" refers to the spatial coordinates of an individual AC in a given year.
Detectors: Potential detection locations in the spatial capture-recapture framework. These can refer to fixed locations (e.g., camera-trap locations) or, in this report, to areas searched (e.g., habitat grid cells where searches for genetic samples were conducted).
Habitat buffer: Buffer surrounding the searched area that is considered potentially suitable habitat but was not searched (60km in this report).
Länsstyrelserna: Swedish County Administrative Boards, in charge of the monitoring of large carnivores at the county level.
MCMC: Markov chain Monte Carlo.
NGS: Non-invasive genetic sampling.
p_0**:** Baseline detection probability; probability of detecting an individual at a given detector, if the individual's AC is located exactly at the detector location.
σ: Scale parameter of the detection function; related to the size of the circular home-range.
RovQuant: Project at the Norwegian University of Life Sciences (Ås, Norway) that develops and applies SCR models.
SCR: Spatial capture-recapture.
Skandobs: Web application (skandobs.se, skandobs.no) that allows anyone to register observations (e.g., visual, tracks, faeces, etc.) of wolverines, bears, wolves and lynx in Sweden and Norway.
SNO: Statens naturoppsyn (Norwegian Nature Inspectorate) is the operative field branch of the Norwegian Environment Directorate (Miljødirektoratet).

2 Methods

2.1 Data

We used data from the Scandinavian large carnivore database Rovbase 3.0 (rovbase.se and rovbase.no; last extraction: 2026-09-29). This database is used jointly by Norway and Sweden to record detailed information associated with large carnivore monitoring, including, but not limited to, NGS data, dead recoveries, and GPS search tracks. In the following sections, we describe the various types of data used in the analysis. We used data collected in Sweden and Norway during the 2025/2026 winter, however, here we report estimates only for Sweden, as NGS data processing and quality control in Norway was not yet complete at the time of this analysis.

Non-invasive genetic sampling In Sweden, the collection of scat and hair is managed by the County Administrative Boards (Länsstyrelserna) at the regional level and carried out by their field officers, with additional samples collected through public participation, particularly by members of Sámi reindeer-herding communities (samebyar). In Norway, the collection of wolverine scat, urine, glandular secretion, and hair is managed at the level of counties by SNO. Sample collection is conducted by SNO field officers, wardens at Statskog Fjelltjenesten (statskog.no), wardens at Fjellstyrene (fjellstyrene.no), local predator contacts, hunters and other members of the public. Rovdata (rovdata.no), a unit within the Norwegian Institute for Nature Research, has responsibility for the Norwegian large carnivore monitoring program. Although samples may be collected throughout the year, the official survey period in Sweden for the 2024/2025 season was Oct 1 – May 15, while in Norway was Dec 1 - June 30. For this analysis, we used data from Oct 1- June 30 to ensure that all NGS samples collected at the beginning of the sampling season in Sweden were included. In addition, samples suspected to be from cubs were not included in the analysis meaning that we only retained samples from individuals that were one year or older. DNA was isolated with an extraction robot (Maxwell 16, KingFisher or QIAsymphony instrument) and samples were genotyped using 96 SNPs (Single Nucleotide Polymorphism) on a microfluidic-based platform (Biomark X9 instrument) for sex determination and individual identification. Genetic analyses were performed by the Swedish University of Agricultural Sciences (SLU, Umeå, Sweden) and the Norwegian Institute for Nature Research (NINA, Trondheim, Norway). For further details on the DNA analysis procedure see Flagstad et al. (2004), Flagstad et al. (2021), and Kleven et al. (2026).

GPS search tracks Government employees involved in structured searches for wolverine DNA (via snowmobiles, skis, snowshoes, etc.) document their effort with GPS track logs, which are registered in Rovbase 3.0. GPS search tracks were included in the SCR model to account for spatial variation in search effort during NGS.

Observation reports in Skandobs We used all observation records in the Skandobs database that were recorded during the wolverine monitoring season (skandobs.se, skandobs.no; last extraction: 2026-09-28). Skandobs is a web application that allows anyone to anonymously register observations (visual, tracks, feces, etc.) of bears (*Ursus arctos*), lynx (*Lynx lynx*), wolves (*Canis lupus*), and wolverines in Scandinavia. This data currently consists of more than 100 000 records of possible large carnivore observations since 2012. Although most observations are not verified, they offer the best available proxy for spatio-temporal variation in opportunistic effort at this time.

2.2 Spatial capture-recapture model

We analysed the data collected in Sweden and Norway during the 2025/2026 monitoring season using a Bayesian single-season spatial capture-recapture (SCR) model (Bischof et al., 2019b), which addresses two challenges associated with population-level wildlife inventories:

1. Detection is imperfect and sampling effort is heterogeneous in space: not all individuals present in the study area are detected (Kéry and Schaub, 2012).
2. Individuals that reside primarily outside the surveyed area may be detected within it. Without an explicit link between the population size parameter and the geographic area the population occupies, density cannot be estimated and population size is ill-defined (Efford, 2004).

The SCR model is composed of three sub-models:

1. A model for population size.
2. A model for density.
3. A model for detections during DNA searches.

Population size sub-model We used data augmentation to estimate population size (Royle et al., 2014). Individual state z_i follows a Bernoulli distribution with probability ψ. Individual state z_i takes the value 1 if the individual is considered as part of the population and 0 otherwise.

Density sub-model We used a Bernoulli point process to model the distribution of individual ACs (Zhang et al., 2022). Individual ACs were located according to an intensity surface, which was a function of the locations of all known dens recorded between 2009 and 2025 (see Bischof et al., 2019b and Bischof et al., 2020 for more details).

Detection sub-model SCR models take into account the spatial variation in individual detection probability based on the distance between AC locations (estimated by the density sub-model) and a given detector. A half-normal function was used to express the declining probability of detection with increasing distance between the AC and the detector (Royle et al., 2014).

DNA material was collected following two main processes: 1) Authorities collected genetic samples and recorded the corresponding search effort during official searches ("structured sampling" hereafter); 2) DNA material was collected by members of the public or by the authorities in a more or less opportunistic manner, which means that search effort is not directly available ("unstructured sampling" hereafter). In the past, it was not possible to unambiguously distinguish between samples collected by the authorities during the structured or unstructured sampling in Rovbase. We therefore assigned each sample to structured or unstructured sampling based on whether it matched in time and space with recorded search tracks. A sample was assigned to the "structured" sampling if it was collected by the authorities (marked as collected by "Statsforvalteren", "Länsstyrelsen", "SNO" or "Fylkesmannen" in Rovbase) and located within 500 m of a GPS search track recorded the same day. All remaining samples were assigned to the unstructured sampling. In addition, 37 DNA samples in Sweden were obtained using hair traps, these samples were considered as being part of the unstructured sampling.

We assumed that both sampling processes could in theory occur within the study area and therefore used the same 10×10 km detector grid for both observation processes. Samples were then assigned to the closest detector (see details in Bischof et al., 2019b, and Bischof et al., 2020). However, spatial variation in the probability to detect a sample during structured or unstructured sampling were assumed to be driven by different processes.

We accounted for spatial and individual heterogeneity in detectability during ***structured sampling*** using:

- Spatial variation in search effort represented by the length of GPS search tracks in each 10×10 km detector grid cell.
- Spatial variation in snow cover during the monitoring period calculated as the average percentage of snow cover in each detector grid cell (MODIS at 0.1 degrees resolution, https://cmr.earthdata.nasa.gov).
- Spatial variation in monitoring regimes between jurisdictions (groups of counties in Sweden, carnivore management regions in Norway)
- Individual variation linked with a detection during the previous occasion (2024/25 monitoring season) that could be expected to influence the probability of being detected at the current occasion.

We accounted for spatial and individual heterogeneity in detectability during ***unstructured sampling*** using:

- Spatial distribution of ancillary samples and samples not successfully genotyped. For each detector grid cell and during the monitoring season (Oct 1 - Jun 30), we identified whether a) any carnivore sample had been registered in Rovbase (excluding successfully genotyped wolverine samples already used in the SCR analysis), b) any observation of carnivores had been registered in Skandobs. Roughly, this binary variable distinguishes areas with very low detection probability from those with a higher probability that carnivore DNA samples, if present in a detector grid cell, could have been detected and submitted for genetic analysis.
- Spatial variation in snow cover during the monitoring period calculated as the average percentage of snow cover in each detector grid cell (MODIS at 0.1 degrees resolution, https://cmr.earthdata.nasa.gov).
- Spatial variation in accessibility measured as the average distance to the nearest road.
- Spatio-temporal variation between countries.
- Individual variation linked with a detection during the previous occasion (2024/2025 monitoring season) that could be expected to influence the probability of being detected at the current occasion.

The different model components and data sources for covariates are described in detail in Bischof et al. (2019a), Bischof et al. (2019b), and Bischof et al. (2020).

Model fitting We fitted sex-specific Bayesian SCR models using MCMC simulation with NIMBLE version 1.4.2 (de Valpine et al., 2017; Turek et al., 2021; de Valpine et al., 2022) and R package nimbleSCR version 0.2.1 (Bischof et al., 2021) in R version 4.6.0 (R Core Team, 2021). We ran 4 chains each with 120 000 iterations, including a 20 000-iterations burn-in period. Due to the computing challenge associated with post-processing large amounts of data, we thinned chains by a factor of 5 before deriving abundance estimates. We considered models as converged when the Gelman-Rubin diagnostics (Rhat, Gelman and Rubin, 1992) was $\leq$1.1 for all parameters and when mixing between chains was satisfactory based on visual inspection of trace plots.

Abundance estimates To obtain an estimate of abundance for any given area, we summed the number of predicted AC locations of live individuals that fell within that area for each iteration of the MCMC chains. This produced a posterior distribution of abundance for that area. From such posteriors, abundance estimates and the associated uncertainty can be extracted for any spatial unit, including countries or counties (Figure 1). Individuals detected near a border (e.g., a county border) can have their model-predicted AC placed on different sides of that border in different model iterations (even if detections are only made on one side of the border). As a result, the probability of designating such individuals to either side of the border can be integrated into county-specific abundance estimates. This is especially relevant for wolverines detected along the Swedish and Norwegian border as individual wolverines can be partially designated to both countries (Bischof, 2015; Bischof et al., 2016).

Note again that, although we fit the SCR model to NGS data from both Sweden and Norway, results are only extracted and reported for Sweden, as entry and quality control of the Norwegian NGS data was not yet complete at the time of this analysis.

To ensure that abundance estimates for spatial sub-units (counties in Sweden, Figure 1) add up to the overall abundance estimate, we used the mean and associated 95% credible interval limits to summarize posterior distributions of abundance. Combined (female and male) parameter estimates were obtained by merging posterior samples from the sex-specific models.



Figure 1: Counties in Sweden.

Density maps We used both the distribution of model-estimated AC positions and the scale parameter (σ) of the detection function to construct density maps based on individual utilization distributions. These maps are not only based on the position of the AC of an individual, but also take into account the area over which that individual's activity is spread, i.e., its space use (Bischof et al., 2020). To do so, we constructed raster maps (5 km resolution) of individual utilization distributions, scaled values in each raster to sum to one, and then summed rasters across individuals to create a single population-level raster map for each iteration. An overall density map was derived by calculating the mean across iterations in each cell (Bischof et al., 2020). Note that this approach assumes circular home ranges of average size for all individuals of a given sex and does not take into account individual variation in home-range size and shape.

Other derived parameters The average proportion of individuals detected, and the associated uncertainty, was obtained by dividing the number of individuals *with their AC estimated in Sweden* that were detected through NGS sampling by the corresponding mean abundance estimates for Sweden for each MCMC iteration. This generated a posterior distribution of the

proportion of the Swedish wolverine population that had been detected, from which the mean and associated 95% credible interval could be extracted.

Focus on uncertainty Although we reported mean estimates in the abundance estimates table, we intentionally focused the main results of our report on the 95% credible interval limits of the estimates. We did so with the aim of drawing the reader's attention to the uncertainty around population size estimates, rather than to a single point estimate (Milleret et al., 2022).

3 Results

3.1 Non-invasive genetic samples and dead recoveries

A total of 2 190 (748 female; 1 442 male) genotyped wolverine genetic samples were collected in Sweden in the winter 2025/2026. These samples were associated with 662 (333 female; 329 male) individuals. Among all genotyped samples, 1 774 (609 female; 1165 male) were assigned to structured sampling and 416 (139 female; 277 male) to unstructured sampling.

3.2 Density and abundance

In the winter 2025/2026, we estimated that wolverine abundance in Sweden was likely (95% credible interval) between 740 and 802 (Figure 2) individuals (Females: 414-472; Males: 319-338). Estimates refer to the status of the population at the start of the annual sampling period in Sweden (October 1). Abundance estimates of wolverines extracted for the different counties are provided in Table 1. Between 79 and 86% of the wolverines estimated to have their AC in Sweden were detected during this study.

The results provided in this report are based on the extended sampling season that includes an earlier starting date in Sweden (Oct 1 – Jun 30). We additionally ran a separate analysis limited to NGS data collected during the Scandinavia-wide sampling season (Dec 1 - Jun 30) used during the join annual analysis (last reports: Milleret et al. (2024a, 2025b)). The wolverine abundance estimates for Sweden during the Dec 1 - Jun 30 sampling season (733-796 individuals) did not differ substantially from the estimates reported for the Oct 1 – Jun 30 sampling season (740-802 individuals).




Figure 2: Wolverine density based on individual utilization distributions in Sweden during winter 2025/2026 based on a single-season spatial capture-recapture model. This map can be accessed as a geo-referenced raster file at https://github.com/richbi/RovQuantPublic

Table 1: Wolverine abundance estimates for the different counties in Sweden in 2025/2026. Numbers are based on estimated AC locations of wolverines. Combined female-male estimates were obtained by joining sex-specific posterior estimates. Credible intervals (95%) are shown in parentheses. These estimates are provided in https://github.com/richbi/RovQuantPublic.

	Females	Males	Total
SWEDEN	441.7 (414-472)	328.3 (319-338)	770 (740-802)
Norra	371.8 (347-399)	265.7 (257-275)	637.4 (611-666)
Jämtland	124.1 (111-138)	92 (86-98)	216.1 (202-232)
Norrbotten	132.5 (118-148)	75.9 (71-82)	208.4 (193-225)
Västerbotten	85.4 (74-98)	62.8 (58-69)	148.2 (136-162)
Västernorrland	29.7 (24-36)	35 (31-39)	64.7 (57-72)
Mellersta	68.5 (59-79)	62.4 (57-68)	130.9 (120-143)
Dalarna	25.4 (20-31)	21.4 (18-25)	46.8 (41-54)
Gävleborg	28.1 (24-33)	24.3 (21-28)	52.3 (47-58)
Örebro	3.4 (2-6)	4.2 (2-6)	7.6 (5-11)
Stockholm	1.5 (0-4)	0.4 (0-2)	1.9 (0-5)
Uppsala	1.6 (0-5)	0.6 (0-2)	2.3 (0-6)
Värmland	6.6 (3-11)	11.1 (8-14)	17.7 (13-23)
Västmanland	1.1 (0-4)	0.2 (0-1)	1.3 (0-4)
Västra Götaland	0.8 (0-3)	0.2 (0-1)	1 (0-3)
Södra	1.4 (0-4)	0.3 (0-2)	1.7 (0-5)
Östergötland	0.1 (0-1)	0 (0-1)	0.2 (0-1)
Södermanland	1.3 (0-4)	0.3 (0-2)	1.5 (0-4)

4 Discussion

We estimated that the Swedish wolverine population during the 2025/2026 winter was between 740 and 802 individuals. These estimates are higher than those reported during the two previous seasons (642-690 individuals in 2023/2024 and 675-731 individuals in 2024/2025, Milleret et al., 2024a, 2025b). This increase in the estimated population size coincides with an increase of 63 (54 males and 9 females) wolverines detected in Sweden during the current monitoring season compared to the previous monitoring season. The higher population size can also be attributed to the larger search area considered in the analysis this year. Even though the new areas incorporated are areas of low population density (e.g., Stockholm county), a larger area potentially translates into a larger abundance estimate. In previous analyses, the search area was defined based on detections recorded during the preceding 10 years, meaning that its extent could vary slightly between years. In the present analysis, we used a larger, fixed search area that will also be applied in future analyses to ensure consistency across years.

The proportion of females in the Swedish population was likely between 56% and 59%. We also provided county-specific estimates for counties in which NGS samples were collected during the 2025/2026 season (i.e., the searched area), which represents the current range of the species. Abundance was estimated to be highest in Northern counties such as Jämtland (202-232), Norrbotten (193-225) and Västerbotten (136-162).

It is important to note that the estimates presented in this report should be considered preliminary. They will be superseded by the final estimates from the annual joint RovQuant wolverine report for Norway and Sweden, scheduled for the end of 2026. The preliminary numbers reported here and the final numbers may differ for two main reasons. First, while this report uses NGS data from Sweden and Norway, only the Swedish data have undergone final processing and quality control. As a result, corrections to Norwegian NGS data collected close to the Swedish border may lead to minor changes in the Swedish estimates. Second, in this report we used data from October 1 to June 30. This differs from the monitoring season in Norway, which extends from December 1 to June 30. The joint RovQuant reports use data from December 1 to June 30 (as it includes Norway and Sweden), meaning that fewer samples will be included in the analysis. However, preliminary analysis (see Results section) suggests that abundance estimates do not differ substantially when the sampling period is shortened to December 1-June 30.

5 Acknowledgments

This work was made possible by the large carnivore monitoring programs and the extensive monitoring data collected by Swedish (Länstyrelsena) and Norwegian (SNO) wildlife management authorities, as well as the public in both countries. Our analyses relied on genetic analyses conducted by the laboratory personnel of the Molecular Ecology Group at the Swedish University of Agricultural Sciences, and NINAGEN group at the Norwegian Institute for Nature Research. We also thank Swedish and Norwegian wildlife managers for feedback provided during project RovQuant. This work was funded by Naturvårdsverket (NV-02444-23), Miljødirektoratet (23047064), and the Research Council of Norway (NFR 286886; project WildMap, NFR 345279: project PopFlow). Computation was performed on resources provided by NMBU's computing cluster Orion. J. Vermaat provided helpful comments on a draft of this report.

6 Data availability

Data, R code to reproduce the analysis, as well as figures, tables, and the raster map (Figure 2) are available at https://github.com/richbi/RovQuantPublic.

References

Bischof, R. (2015). Approaches for assessing illegal hunting of brown bears and other large carnivores in Sweden. *Report 2015:5 from the Scandinavian Brown Bear Research Project to the Swedish Environmental Protection Agency. Norwegian University of Life Sciences, Ås, Norway.*

Bischof, R., Brøseth, H., and Gimenez, O. (2016). Wildlife in a politically divided world: Insularism inflates estimates of brown bear abundance. *Conservation Letters*, 9(2):122–130.

Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Åkesson, M., Brøseth, H., and Kindberg, J. (2019a). Estimating the size of the Scandinavian wolf population with spatial capture recapture and conversion factors. *MINAfagrapport 57*, 80pp.

Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Brøseth, H., and Kindberg, J. (2019b). Estimating density, abundance and population dynamics of bears, wolverines, and wolves in Scandinavia. *MINAfagrapport 63*, 79pp.

Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Tourani, M., Ordiz, A., de Valpine, P., Turek, D., Royle, J. A., Gimenez, O., Flagstad, O., Åkesson, M., Svensson, L., Brøseth, H., and Kindberg, J. (2020). Estimating and forecasting spatial population dynamics of apex predators using transnational genetic monitoring. *Proceedings of the National Academy of Sciences*, 117(48):30531–30538.

Bischof, R., Turek, D., Milleret, C., Ergon, T., Dupont, P., Dey, S., and de Valpine, P. (2021). *nimbleSCR: Spatial Capture-Recapture (SCR) Methods Using 'nimble'*. R package version 0.1.2.

de Valpine, P., Paciorek, C., Turek, D., Michaud, N., Anderson-Bergman, C., Obermeyer, F., Wehrhahn Cortes, C., Rodrìguez, A., Temple Lang, D., and Paganin, S. (2022). *NIMBLE User Manual*. R package manual version 0.12.2.

de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., and Bodik, R. (2017). Programming with models: writing statistical algorithms for general model structures with NIMBLE. *Journal of Computational and Graphical Statistics*, 26(2):403–413.

Dupont, P., Hansen, N. R., Milleret, C., Semper-Pascual, A., Brøseth, H., Kopatz, A., Kindberg, J., and Bischof, R. (2026a). Estimates of brown bear density, abundance, and population dynamics in Norway 2016-2025. *MINA fagrapport 115*, 35pp.

Dupont, P., Milleret, C., Semper-Pascual, A., Brøseth, H., Kopatz, A., Kindberg, J., and Bischof, R. (2025). Estimates of brown bear density, abundance, and population dynamics in Norway 2015-2024. *MINA fagrapport 105*, 35pp.

Dupont, P., Milleret, C., Semper-Pascual, A., Brøseth, H., Flagstad, , Kindberg, J., Svensson, L., , and Bischof, R. (2026b). Estimates of wolf density, abundance, and population dynamics in Sweden and Norway, 2016/17–2025/26. *MINAfagrapport 113*, 37pp.

Efford, M. (2004). Density estimation in live-trapping studies. *Oikos*, 106(3):598–610.

Flagstad, Ø., Hedmark, E., Landa, A., Brøseth, H., Persson, J., Andersen, R., SegerstrOm, P., and Ellegren, H. (2004). Colonization history and noninvasive monitoring of a reestablished wolverine population. *Conservation Biology*, 18(3):676–688.

Flagstad, Ø., Kleven, O., Brandsegg, H., Spets, M., Eriksen, L., Andersskog, I., Johansson, M., Ekblom, R., Ellegren, H., and Brøseth, H. (2021). DNA-basert overvåking av den skandinaviske jervebestanden 2020. *Norsk institutt for naturforskning(NINA), Trondheim*, NINA Rapport nr 1956.

Gelman, A. and Rubin, D. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7:457–511. `http://www.stat.columbia.edu/~gelman/research/published/itsim.pdf`.

Kéry, M. and Schaub, M. (2012). *Bayesian population analysis using WinBUGS: a hierarchical perspective*. Academic Press, Waltham, MA.

Kleven, O., Bergen, S., Bischof, R., Dupont, P., Hedmark, E., Königsson, H., Milleret, C., Spets, M., Spong, G., and Brøseth, H. (2026). DNA-basert overvåking av den skandinaviske jervebestanden 2025. *Norsk institutt for naturforskning(NINA)*, NINA Rapport nr 2731.

Milleret, C., Dupont, P., Brøseth, H., Flagstad, O., Kindberg, J., and Bischof, R. (2022). Estimates of wolverine density, abundance, and population dynamics in Scandinavia, 2013–2021. *MINAfagrapport 74*, 30pp.

Milleret, C., Dupont, P., Brøseth, H., Flagstad, O., Kleven, O., Königsson, H., Spong, G., Kindberg, J., and Bischof, R. (2024a). Estimates of wolverine density, abundance, and population dynamics in Scandinavia, 2015-2024. *MINAfagrapport 101*, 34pp.

Milleret, C., Dupont, P., Semper-Pascual, A., Brøseth, H., Flagstad, , Kindberg, J., Svensson, L., , and Bischof, R. (2025a). Estimates of wolf density, abundance, and population dynamics in Scandinavia, 2015–2025. *MINAfagrapport 85*, 36pp.

Milleret, C., Dupont, P., Semper-Pascual, A., Hansen, N. R., Brøseth, H., Flagstad, O., Hedmark, E., Kleven, O., Königsson, H., Spong, G., Kindberg, J., and Bischof, R. (2025b). Estimates of wolverine density, abundance, and population dynamics in Scandinavia, 2015/16-2024/25. *MINAfagrapport 112*, 34pp.

Milleret, C., Dupont, P., Winiger, A., Spong, G., Königsson, H., and Bischof, R. (2024b). Estimates of wolverine

density and abundance in Norrbotten county in Sweden and associated reindeer herding areas, 2023/2024. *MINAfagrapport 100*, 20pp.

R Core Team (2021). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Royle, J. A., Chandler, R. B., Sollmann, R., and Gardner, B. (2014). *Spatial Capture-Recapture*. Academic Press.

Royle, J. A., Karanth, K. U., Gopalaswamy, A. M., and Kumar, N. S. (2009). Bayesian inference in camera trapping studies for a class of spatial capture-recapture models. *Ecology*, 90(11):3233–3244.

Semper-Pascual, A., Milleret, C., Dupont, P., Hedmark, E., Spong, G., Königsson, H., and Bischof, R. (2025). Preliminary estimates of wolverine density and abundance in Sweden, 2024/2025. *MINAfagrapport 106*, 17pp.

Turek, D., Milleret, C., Ergon, T., Brøseth, H., Dupont, P., Bischof, R., and de Valpine, P. (2021). Efficient estimation of large-scale spatial capture–recapture models. *Ecosphere*, 12(2):e03385.

Zhang, W., Chipperfield, J. D., Illian, J. B., Dupont, P., Milleret, C., de Valpine, P., and Bischof, R. (2022). A flexible and efficient bayesian implementation of point process models for spatial capture-recapture data. *Ecology*, page e3887. e3887.