

**Norwegian University of Life Sciences**

Faculty of Environmental Sciences and Natural Resource Management

**2022**

ISSN 2535-2806

MINA fagrapport 79

## **Estimates of wolverine density, abundance, and population dynamics in Scandinavia, 2014–2022**

Cyril Milleret  
Pierre Dupont  
Ehsan Moqanaki  
Henrik Brøseth  
Øystein Flagstad  
Oddmund Kleven  
Jonas Kindberg  
Richard Bischof



Milleret, C., Dupont, P., Moqanaki, E., Brøseth, H., Flagstad, Ø, Kleven, O., Kindberg, J., and Bischof, R., 2022. **Estimates of wolverine density, abundance, and population dynamics in Scandinavia, 2014–2022** - MINA fagrapport 79. 35 pp.

Ås, December 2022

ISSN: 2535-2806

COPYRIGHT

© Norwegian University of Life Sciences (NMBU)

The publication may be freely cited where the source is acknowledged

AVAILABILITY

Open

PUBLICATION TYPE

Digital document (pdf)

QUALITY CONTROLLED BY

The Research committee (FU), MINA, NMBU

PRINCIPAL

Naturvårdsverket, Ref: NV-04078-22, Contact person: Robert Ekblom

Miljødirektoratet, Ref: 22047026, Contact person: Terje Bø

COVER PICTURE

Wolverine, J. Percy/Shutterstock.

NØKKEORD

*Gulo gulo*, jerv, tetthet, populasjonsdynamikk, deteksjonssannsynlighet, ikke-invaserende innsamling av genetisk materiale, åpen populasjon romlig fangst-gjenfangst, rovdyrforvaltning

KEY WORDS

*Gulo gulo*, wolverine, population density, population dynamics, detection probability, non-invasive genetic sampling, open-population spatial capture-recapture, carnivore management

Cyril Milleret, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway

Pierre Dupont, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway

Ehsan Moqanaki, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway

Henrik Brøseth, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway

Øystein Flagstad, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway

Oddmund Kleven, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway

Jonas Kindberg, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway


Richard Bischof ([richard.bischof@nmbu.no](mailto:richard.bischof@nmbu.no)), Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway

## Summary

**Background** The Scandinavian wolverine (*Gulo gulo*) population is being monitored annually using non-invasive genetic sampling (NGS) and recovery of dead individuals. DNA extracted from feces, urine, hair, secretion, and tissue is used to identify the species, sex, and individual from which each sample originated. These data have been compiled in the Scandinavian large carnivore database Rovbase 3.0. ([www.rovbase.se](http://www.rovbase.se), [www.rovbase.no](http://www.rovbase.no)).

**Approach** Using the Bayesian open-population spatial capture-recapture (OPSCR) model developed by RovQuant, we estimated annual density and vital rates of the Scandinavian wolverine population for nine consecutive seasons from 2014 to 2022.

**Results** We generated annual density maps and estimated total and jurisdiction-specific population sizes for the wolverine during 2014 to 2022. Based on the OPSCR model, the size of the Scandinavian wolverine population was likely (95% credible interval) between 980 and 1088 individuals in 2022, with 625 to 709 individuals attributed to Sweden and 349 to 391 to Norway. In addition to annual density and jurisdiction-specific abundance estimates, we report, for each sex, annual estimates of cause-specific mortality, recruitment, and detection probability.




Density map and ranges of abundance estimated for wolverines in 2022

## Sammendrag

**Bakgrunn** Den skandinaviske bestanden av jerv (*Gulo gulo*) blir overvåket årlig ved bruk av ikke-invasiv genetisk prøveinnsamling (NGS) og gjenfunn av døde individer. DNA ekstrahert fra skit, urin, hår og vev brukes til å identifisere art, kjønn og individ for hver enkelt prøve. Denne informasjonen samles og ivaretas i den skandinaviske databasen for store rovdyr; Rovbase 3.0 ([www.rovbase.se](http://www.rovbase.se), [www.rovbase.no](http://www.rovbase.no)).

**Tilnærming** Ved bruk av en Bayesiansk åpen romlig fangst-gjenfangst populasjons modell (OPSCR), utviklet av RovQuant, estimerte vi årlige tettheter og demografiske rater hos den skandinaviske jervebestanden i ni sesonger fra 2014 til 2022.

**Resultater** Vi laget årlige kart med tetthet av jerv fra 2014 til 2022, hvor bestandsstørrelsen både totalt og innenfor ulike administrative enheter kunne avledes. Basert på OPSCR modellen var den skandinaviske bestanden av jerv mellom 980 og 1088 individer i 2022 (95% kredibelt intervall), med 625 til 709 individer i Sverige og 349 til 391 individer i Norge. I tillegg til årlige tettheter og områdespesifikke bestandsestimater, gir rapporten estimater på dødlighetsfaktorer, rekruttering og oppdagbarhet.



Kart som viser tetthet av jerv i 2022 sammen med intervaller for estimert antall jerv

# Contents

<b>1</b>	<b>Introduction</b>	<b>6</b>
<b>2</b>	<b>Methods</b>	<b>8</b>
2.1	Data . . . . .	8
2.2	Open-population spatial capture-recapture model . . . . .	9
<b>3</b>	<b>Results</b>	<b>13</b>
3.1	Non-invasive genetic samples and dead recoveries . . . . .	13
3.2	Density and abundance . . . . .	13
3.3	Vital rates . . . . .	15
3.4	Detection probability . . . . .	17
<b>4</b>	<b>Summary of improvements made</b>	<b>18</b>
<b>5</b>	<b>Suggestions for future improvements</b>	<b>18</b>
<b>6</b>	<b>Other recommendations</b>	<b>18</b>
<b>7</b>	<b>Acknowledgements</b>	<b>19</b>
<b>8</b>	<b>Data availability</b>	<b>19</b>
	<b>References</b>	<b>21</b>
	<b>Appendices</b>	<b>22</b>

# 1 Introduction

Sweden and Norway monitor large carnivores using non-invasive genetic sampling (NGS) and dead recoveries. Both countries have collected an extensive individual-based data set for the wolverine (*Gulo gulo*), which is stored in the Scandinavian large carnivore database Rovbase ([www.rovbase.se](http://www.rovbase.se), [www.rovbase.no](http://www.rovbase.no)).

Since 2017, project RovQuant has been developing statistical methods that allow a comprehensive assessment of the status and dynamics of large carnivore populations using NGS data and other sources of information stored in Rovbase (Bischof et al., 2019b, 2020b). The analytical framework developed by RovQuant is based on Bayesian open-population spatial capture-recapture (OPSCR) models (Ergon and Gardner, 2014; Bischof et al., 2016; Chandler et al., 2018). These models use the spatial and temporal information contained in the repeated genetic detections of individuals to estimate various population parameters, including spatially-explicit abundance (i.e., density) and vital rates (e.g., recruitment and survival). Importantly, the approach accounts for imperfect detection during sampling (i.e., the fact that some individuals are not detected at all) and animal movement (i.e., the fact that individuals may use and be detected in multiple management units or countries). The OPSCR method brings along several advantages, including the ability to map density, derive jurisdiction-specific abundance, estimate survival and recruitment (which are needed for making population projections), and yield tractable measures of uncertainty (Bischof et al., 2019a, 2020b).

RovQuant reported abundance estimates for wolverines and wolves (*Canis lupus*) on an annual basis (Bischof et al., 2019a,b, 2020b; Milleret et al., 2021b, 2022b,c; Flagstad et al., 2021). During these and other analyses (Milleret et al., 2018, 2019; Bischof et al., 2020a; Dupont et al., 2021; Turek et al., 2021; Dey et al., 2022), RovQuant has continuously improved the performance of the OPSCR models. In the present report, we summarize the analysis of a 9-year time series (2014–2022) using the latest available wolverine monitoring data (Kleven et al., 2022b) and the most recent version of the OPSCR model. We provide the following information:

- Annual and sex-specific estimates of the number of wolverines for Sweden, Norway, and both countries combined, as well as estimates by county in Sweden and by carnivore management region in both countries.
- Annual maps of wolverine density throughout the species' range in Scandinavia.
- Annual and sex-specific estimates of survival, cause-specific mortality, recruitment, and population growth rate.
- Estimated proportion of individuals detected through non-invasive genetic sampling.

All estimates are accompanied by credible intervals.

## Box 1: Terms and acronyms used

**AC:** Activity center. Model-based equivalent of the center of an individual's home range during the monitoring period. "AC location" refers to the spatial coordinates of an individual AC in a given year and "AC movement" to the movement of an individual AC between consecutive years.

**CrI:** 95% credible interval associated with a posterior sample distribution.

**Detectors:** Potential detection locations in the spatial capture-recapture framework. These can refer to fixed locations (e.g., camera-trap locations) or in this report to areas searched (e.g., habitat grid cells where searches for genetic samples were conducted). The searched area was defined as a 90 km buffer around all NGS data collected during the period considered.

**Statsforvalteren:** Norwegian state's representative in the county, responsible for following up decisions, goals, and guidelines from the legislature and the government.

**Habitat buffer:** Buffer surrounding the searched area that is considered potentially suitable habitat but was not searched (60km in this report).

**Legal culling:** Lethal removal of individuals by legal means, including licensed recreational hunting, management removals, and defense of life and property.

**Länsstyrelserna:** Swedish County Administrative Boards, in charge of the monitoring of large carnivores at the county level.

**MCMC:** Markov chain Monte Carlo.

**NGS:** Non-invasive genetic sampling.

**OPSCR:** Open-population spatial capture-recapture

**$p_0$ :** Baseline detection probability; probability of detecting an individual at a given detector, if the individual's AC is located exactly at the detector location.

**$\sigma$ :** Scale parameter of the detection function; related to the size of the circular home-range.

**SCR:** Spatial capture-recapture.

**SNO:** Statens naturoppsyn (Norwegian Nature Inspectorate) is the operative field branch of the Norwegian Environment Directorate (Miljødirektoratet).

**RovQuant:** Research group at the Norwegian University of Life Sciences (Ås, Norway) that develops and applies OPSCR models.

## 2 Methods

### 2.1 Data

We included data from multiple sources, the primary one being the Scandinavian large carnivore database Rovbase 3.0 (rovbase.se and rovbase.no; last extraction: 2022-10-28). This database is used jointly by Norway and Sweden to record detailed information associated with large carnivore monitoring, including, but not limited to, NGS data, dead recoveries, and GPS search tracks. In the following sections, we describe the various types of data used in the analysis. We used data collected during nine consecutive monitoring seasons from 2014 to 2022.

**Non-invasive genetic sampling** In Norway, the collection of wolverine scat, urine, glandular secretion, and hair is managed at the level of counties by SNO. Sample collection is conducted by SNO field officers, wardens at Statskog Fjelltjenesten (statskog.no), wardens at Fjellstyrene (fjellstyrene.no), local predator contacts, hunters and other members of the public. Rovdata (rovdata.no), a unit within the Norwegian Institute for Nature Research, has responsibility for the Norwegian large carnivore monitoring program. In Sweden, the collection of scat and hair is managed by Länsstyrelserna at the regional level and carried out by field officers from Länsstyrelserna. NGS collection was conducted primarily between December 1 and June 30 each year. NGS data collected late in the monitoring season and suspected to be from cubs were not included. This means that we only retained samples from individuals that were one year or older. DNA was isolated with an extraction robot (Maxwell 16, KingFisher or QIA Symphony instrument) and the samples were genotyped using 96 SNPs (Single Nucleotide Polymorphism) on a Fluidigm platform for sex determination and individual identification. For further details on the DNA analysis procedure see Flagstad et al. (2004), Flagstad et al. (2021), Kleven et al. (2022a), and Kleven et al. (2022b).

**Dead recoveries** In Scandinavia, all large carnivores killed legally (e.g., legal hunting, management kills, defense of life and property) have to be reported to the state authorities (Fylkesmannen or SNO in Norway and Länsstyrelserna or the police in Sweden). All wolverines found dead due to other reasons (e.g., natural deaths, vehicle and train collisions, illegal hunting) also have to be reported, but an unknown proportion remains undetected. Tissue is collected from all reported dead carnivores for DNA extraction and analysis.

**GPS search tracks** Government employees involved in systematic searches for wolverine DNA along roads and following wolverine tracks (via snowmobiles, skis, snowshoes, etc.) documented their effort with GPS track logs, which were registered in Rovbase 3.0. GPS search tracks were included in the OPSCR model to account for spatial and temporal variation in search effort during NGS.

**Observation reports in Skandobs** We used all observation records in the Skandobs database that were recorded during the wolverine monitoring seasons since 2012 (skandobs.se, skandobs.no; last extraction: 2022-06-09). Skandobs is a web application that allows anyone to anonymously register observations (visual, tracks, feces, etc.) of bears (*Ursus arctos*), lynx (*Lynx lynx*), wolves, and wolverines in Scandinavia. This data currently consists of more than 20 000 records of possible large carnivore observations. Although most observations are not verified, they offer the best available proxy for spatio-temporal variation in opportunistic effort at this time.



## 2.2 Open-population spatial capture-recapture model

We analysed the data using a Bayesian open-population spatial capture-recapture (OPSCR) model (Bischof et al., 2019b), which addresses three challenges associated with population-level wildlife inventories:

1. Detection is imperfect and sampling effort is heterogeneous in space and time: not all individuals present in the study area are detected (Kéry and Schaub, 2012).
2. Individuals that reside primarily outside the surveyed area may be detected within it. Without an explicit link between the population size parameter and the geographic area the population occupies, density cannot be estimated and population size is ill-defined (Efford, 2004).
3. Non-spatial population dynamic models usually estimate “apparent” survival and recruitment, as these parameters include the probability of permanent emigration and immigration, respectively. By explicitly modelling movement of individuals between years, the OPSCR model can help return unbiased estimates of demographic parameters (Ergon and Gardner, 2014; Schaub and Royle, 2014; Gardner et al., 2018).

The OPSCR model is composed of three sub-models:

1. A model for population dynamics and population size.
2. A model for density and individual movement.
3. A model for detections during DNA searches.

**Population dynamics and population size sub-model** We used a multi-state formulation (Lebreton and Pradel, 2002), where each individual’s life history is represented by a succession of up to 3 discrete states  $z_{i,t}$ : (1) “unborn” if the individual has not yet been recruited into the population (state “unborn” is required for the data augmentation procedure, see below); (2) “alive” if it is alive; (3) “dead” if it is dead. We then modelled the transition from one state to another between consecutive monitoring seasons ( $t$  to  $t + 1$ ) to estimate vital rates (recruitment and mortality). More details are available in Bischof et al. (2019b) and Bischof et al. (2020b). This formulation of the population dynamic model means that, contrary to previous analyses (Bischof et al., 2019b, 2020b; Flagstad et al., 2021; Milleret et al., 2022b), we did not use dead recoveries or model cause-specific mortality directly in the OPSCR model. Cause-specific mortality was instead derived after model fitting (see section “Other derived parameters”). We used data augmentation (Royle and Dorazio, 2012), whereby additional, undetected individuals are available for inclusion in the population at each time step.

**Density and movement sub-model** We used a Bernoulli point process to model the distribution of individual ACs (Zhang et al., 2022). In the first year, individuals were located according to an intensity surface, which was a function of the locations of known dens at time  $t - 1$  (see Bischof et al., 2019b and Bischof et al., 2020b for more details). For all subsequent years ( $t > 1$ ), the location of individual ACs was a function of the distance from previous locations of ACs (at time  $t - 1$ ) and the locations of known wolverine dens (at time  $t - 1$ ). Similar to the wolf abundance estimation by Milleret et al. (2022c), we used an exponential model to describe the movement of individuals between years, as it better accommodates distributions with long tails (i.e., a few individuals that make exceptionally long dispersal movements).

**Detection sub-model** SCR models take into account the spatial variation in individual detection probability based on the distance between AC locations (estimated by the density sub-model) and a given detector. A half-normal function was used to express the declining probability of detection with increasing distance between the AC and the detector (Royle et al., 2013).

In Scandinavia, DNA material from live wolverines is collected following two main processes. First, authorities collect genetic samples and record the corresponding search effort during official searches ("structured sampling" thereafter). Second, DNA material can be collected by any member of the public (e.g., hunters) or by the authorities in a more or less opportunistic manner, which means that search effort is not directly available ("unstructured sampling" thereafter). Currently, it is not possible to unambiguously distinguish between samples collected by the authorities during the structured or unstructured sampling in Rovbase. We therefore assigned each sample to structured or unstructured sampling based on whether a given sample matched in time and space with recorded search tracks: a sample was assigned to the "structured" sampling if it was collected by the authorities (marked as collected by "Statsforvalteren" or "SNO" in Rovbase) and located within 500 m from a GPS search track recorded the same day. All remaining samples were assigned to the unstructured sampling.

We assumed that both sampling processes could in theory occur within the entire study area and therefore used the same  $10 \times 10$  km detector grid for both observation processes. Samples were then assigned to the closest detector (see details in Bischof et al., 2019b, and Bischof et al., 2020b). However, spatial and temporal variation in the probability to detect a sample during structured or unstructured sampling were assumed to be driven by different processes.

We accounted for spatial, temporal and individual heterogeneity in detectability during *structured sampling* using:

- Spatio-temporal variation in search effort represented by the length of GPS search tracks in each detector grid cell.
- Spatio-temporal variation in snow cover during the monitoring period calculated as the average percentage of snow cover in each detector grid cell (MODIS at 0.1 degrees resolution, <https://cmr.earthdata.nasa.gov>, accessed 2022-09-29).
- Spatio-temporal variation in monitoring regimes between jurisdictions (groups of counties in Sweden, carnivore management regions in Norway, Figure A.6).
- Individual variation linked with a detection during the previous occasion that could be expected to influence the probability of being detected at the next occasion.

We accounted for spatial, temporal, and individual heterogeneity in detectability during *unstructured sampling* using:

- Spatio-temporal variation in unstructured sampling (Figure A.1). For each detector grid cell and during each monitoring season (Dec 1 - Jun 30), we identified whether a) any carnivore sample had been registered in Rovbase (excluding successfully genotyped wolverine samples already used in the OPSCR analysis) or b) any observation of carnivores had been registered in Skandobs. Roughly, this binary variable distinguishes areas with very low detection probability from those with a higher probability that carnivore DNA samples, if present in a detector grid cell, could have been detected and submitted for genetic analysis (Figure A.1).
- Spatio-temporal variation in snow cover during the monitoring period calculated as the average percentage of snow cover in each detector grid cell (MODIS at 0.1 degrees resolution, <https://cmr.earthdata.nasa.gov>, accessed 2022-09-29).
- Spatial variation in accessibility measured as the average distance to the nearest road.
- Spatio-temporal variation between countries Figure A.7).

- Individual and temporal variation linked with a previous detection that could influence the probability of being detected at subsequent occasions.

For years and areas without comprehensive sampling effort (i.e., Norrbotten county in Sweden in all years except 2017, 2018, and 2019), we removed all samples collected within the county and fixed detection probability to 0 for both structured and unstructured sampling. The different model components and data sources for covariates are described in detail in Bischof et al. (2019a), Bischof et al. (2019b), and Bischof et al. (2020b).

**Model fitting** We fitted sex-specific Bayesian OPSCR models using MCMC simulation with NIMBLE version 0.12.2 (de Valpine et al., 2017; Turek et al., 2021; de Valpine et al., 2022) and RovQuant’s R package nimbleSCR version 0.2.0 (Bischof et al., 2021) in R version 4.1.0 (R Core Team, 2021). We ran 4 chains each with 25 000 iterations, including a 10 000-iterations burn-in period. Due to the computing challenge associated with post-processing large amounts of data, we thinned chains by a factor of 10 before deriving abundance estimates. We considered models as converged when the Gelman-Rubin diagnostics (Rhat, Gelman and Rubin, 1992) was  $\leq 1.1$  for all parameters and when mixing between chains was satisfactory based on visual inspection of trace plots.

**Abundance estimates** To obtain an estimate of abundance for any given area, we summed the number of predicted AC locations (individuals detected during sampling or predicted to be alive by the model) that fell within that area for each iteration of the MCMC chains. This produced a posterior distribution of abundance for that area. From such posteriors, abundance estimates and the associated uncertainty can be extracted for any spatial unit, including countries, counties or management regions (Figure A.2). Individuals detected near a border can have their model-predicted AC placed on different sides of that border in different model iterations (even if detections are only made on one side of the border). As a result, the probability of designating such individuals to either side of the border can be integrated into jurisdiction-specific abundance estimates. This is especially relevant for wolverines detected along the Swedish and Norwegian border ("cross-boundary wolverines", Wabakken et al., 2022) as individual wolverines can be partially designated to both countries (Bischof, 2015; Bischof et al., 2016).

To ensure that abundance estimates for spatial sub-units (jurisdictions) add up to the overall abundance estimate, we used the mean and associated 95% credible interval limits to summarize posterior distributions of abundance. Combined (female and male) parameter estimates were obtained by merging posterior samples from the sex-specific models.

**Density maps** We used both the distribution of model-estimated AC positions and the scale parameter ( $\sigma$ ) of the detection function to construct density maps based on individual utilization distributions. These maps are not only based on the position of the activity center of an individual, but also take into account the area over which that individual’s activity is spread, i.e., its space use (Bischof et al., 2020b). To do so, we constructed raster maps (5 km resolution) of individual utilization distributions, scaled values in each raster to sum to one, and then summed rasters across individuals to create a single population-level raster map for each iteration. An overall density map was derived by calculating the mean across iterations in each cell (Bischof et al., 2020b). Note that this approach assumes circular home ranges of average size for all individuals of a given sex and does not take into account individual variation in home-range size and shape.

**Other derived parameters** We did not use dead recoveries and did not model cause-specific mortalities explicitly. Instead, we used the posterior distribution of the number of individuals that died between consecutive occasions, as estimated by the OPSCR model, and the recorded

number of legally culled individuals to derive cause-specific mortality estimates indirectly.

The average proportion of individuals detected and the associated uncertainty were obtained by dividing the number of individuals detected through NGS each year (Table A.2) by corresponding the abundance estimates and their associated credible intervals, respectively.

We derived the proportion of females in the population and the associated uncertainty by dividing the estimated number of females by the total abundance for each iteration, thus generating a posterior distribution of the proportion of females from which the median and credible interval could be derived (Table A.2). Yearly population growth rates ( $\lambda$ ; Table A.5) were calculated similarly as  $\lambda_t = N_{t+1}/N_t$  for each iteration of the MCMC chains.

**Focus on uncertainty** Although we reported median (or mean for abundance; see above) estimates for all parameters in the tables, we intentionally focused the main results of our report on the 95% credible interval limits of the estimates. We did so with the aim of drawing the reader's attention to the uncertainty around population size estimates, rather than a single point estimate (Milleret et al., 2022b).

## 3 Results

### 3.1 Non-invasive genetic samples and dead recoveries


A total of 18 745 (8 418 female; 10 327 male) genotyped wolverine genetic samples were included in the analysis, of which 43% originated from Sweden. These samples were associated with 2 550 (1360 female; 1190 male) individuals. We did not include individuals with unknown sex in this analysis. Among all genotyped samples, 12 021 (5298 female; 6723 male) were assigned to structured sampling and 6 724 (3120 female; 3604 male) to unstructured sampling. Annual total and country-specific tallies of detections and associated individuals, as well as dead recoveries are provided in the appendices (NGS samples: Table A.1, NGS individuals: Table A.2, dead recoveries: Table A.3)

### 3.2 Density and abundance

Wolverine abundance for the entire study area (612 350 km<sup>2</sup>, excluding the buffer area) was likely (95% credible interval) between 980 and 1 088 individuals in 2022 (Table 1, Figure 1). Estimates refer to the status of the population at the start of the annual sampling period (December 1). The proportion of females in the Scandinavian wolverine population was likely between 59% and 63% in 2022. Based on the model-predicted location of ACs, we estimated that in 2022, between 625 and 709 individuals could be attributed to Sweden and 349 to 391 to Norway (Table 1). See Table 1 for total and sex-specific estimates for each country and carnivore management region. See Table A.4 for annual estimates for all of Scandinavia and by region between 2014 and 2022. Note that estimates for different years (Figure A.3) shown here differ slightly from those provided in Bischof et al. (2020b) and Milleret et al. (2021a). This is due to the use of an updated OPSCR model and the inclusion of an additional year of data. The analysis yielded annual density maps, which illustrate changes in the distribution of wolverines over time (Figure A.4).

**Table 1:** Wolverine population size estimates in 2022 by sex at several spatial scales: the entire study area, by country, by management unit (carnivore management regions in Norway and "Rovdjursförvaltningsområden" in Sweden), and by counties ("Län" in Sweden); see Figure A.2 for a labelled map. Only counties and management units that are within or that intersect the study area are included in the table. The percentage of the total area of each unit included in the analysis is provided in the column "% Area". Readers should focus on the 95% credible interval provided in parentheses, as these - unlike mean values - convey uncertainty inherent in abundance estimates. Numbers are based on estimated AC locations of wolverines. Combined female-male estimates were obtained by joining sex-specific posterior distributions. Rounding may result in small deviations between total estimates and the sum of the estimates for constituent regions. Estimates for Norrbotten county in years without comprehensive non-invasive genetic sampling were derived solely using the prediction from the OPSCR model (shown in grey and marked with \*). Total estimates in Sweden and for the entire study area that includes estimates from Norrbotten without comprehensive NGS are shown in grey and marked with \*\*.


	Females	Males	Total	% Area
TOTAL	632.9 (585-680)**	402.3 (380-427)**	1035.2 (980-1088)**	82
NORWAY	229.6 (212-251)	137.7 (130-148)	367.2 (349-391)	93
Region 1	6 (2-11)	4.5 (2-8)	10.4 (6-16)	84
Region 2	3 (0-7)	2.1 (0-5)	5.1 (1-9)	73
Region 3	16.9 (13-22)	10.1 (7-13)	27 (21-33)	100
Region 4	0.7 (0-3)	0.5 (0-2)	1.2 (0-4)	75
Region 5	41.2 (36-48)	39.8 (36-44)	81 (75-89)	100
Region 6	58.5 (52-66)	33.2 (29-37)	91.7 (84-100)	100
Region 7	44.4 (39-51)	22.3 (20-25)	66.7 (61-74)	100
Region 8	58.8 (51-70)	25.3 (21-30)	84.1 (75-96)	100
SWEDEN	403.3 (367-441)**	264.6 (245-286)**	667.9 (625-709)**	74
Norra	349 (315-383)	208.7 (188-231)	557.6 (517-598)	100
Jämtland	122.4 (109-136)	71.8 (65-79)	194.1 (180-211)	100
Norrbotten*	114.3 (92-136)*	73 (56-92)*	187.3 (158-216)*	97
Västerbotten	77.9 (67-90)	39.1 (34-45)	117 (104-129)	96
Västernorrland	34.4 (28-41)	24.8 (20-29)	59.2 (51-67)	100
Mellersta	54.1 (45-63)	55.8 (50-62)	109.9 (99-122)	68
Dalarna	28.2 (22-34)	26.8 (23-31)	54.9 (48-62)	100
Gävleborg	14.2 (10-18)	15.9 (13-20)	30.1 (25-36)	100
Örebro	1.7 (0-4)	1.2 (0-4)	2.8 (0-7)	100
Uppsala	0.1 (0-1)	0 (0-1)	0.1 (0-1)	5
Värmland	8.9 (6-13)	11.2 (8-15)	20.1 (16-25)	100
Västmanland	0.6 (0-2)	0.4 (0-2)	0.9 (0-3)	62
VästraGötaland	0.5 (0-2)	0.4 (0-2)	0.9 (0-3)	10
Södra	0.3 (0-2)	0.2 (0-1)	0.4 (0-2)	1
Östergötland	0.1 (0-1)	0.1 (0-1)	0.2 (0-2)	4
Södermanland	0.1 (0-1)	0.1 (0-1)	0.2 (0-1)	7




**Figure 1:** Wolverine density based on individual utilization distributions in Scandinavia in 2022. Note that no comprehensive NGS was conducted in Norrbotten (polygon outlined in black) in 2022, which means that the results are solely based on the OPSCR model prediction and assumption. The grey area represents areas that were considered not searched and therefore were not included in the analysis. This map is freely available as a geo-referenced raster file at <https://github.com/richbi/RovQuantPublic>

### 3.3 Vital rates

The OPSCR model produced annual estimates of mortality and per capita recruitment rates (Figure 2; Table A.6). Mortality rates varied between years, with the risk of mortality from other causes than culling generally higher than the risk of mortality from legal culling. Overall, males had a higher mortality probability than females (Figure A.5).



**Figure 2:** Mortality probabilities due to legal culling (light green) and all other causes (dark green) for female and male wolverines. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Shown are overall estimates throughout the study area. Estimates refer to deaths occurring between the start of one sampling season and the start of the next.



**Figure 3:** Estimated annual number of male and female recruits in the Scandinavian wolverine population between the start of one sampling season and the start of the next. Recruitment represents the number of new individuals present in the population on Dec 1 (i.e., individuals that were born between the two consecutive monitoring seasons and survived to Dec 1 or that immigrated into the study area). Darker and lighter bars show the 50% and 95% credible intervals, respectively.



### 3.4 Detection probability

The overall proportion of detected individuals in the population was likely between 63% and 70% in 2022, and overall, larger in Norway than in Sweden (Table A.10). The baseline detection probability for the structured and unstructured sampling varied both in time and space (Figure A.6 and Figure A.7). More specifically, the length of recorded search tracks positively affected detection probability during structured sampling (2022; males:  $\beta = 0.45$ , CrI: 0.37 - 0.52; females:  $\beta = 0.42$ , CrI: 0.35 - 0.49; Table A.8). Detection of an individual during the previous year and the average proportion of snow cover had no significant effect on detection probability during structured sampling (Table A.8). The proxy for search effort during unstructured searches, derived using the observation data in Skandobs and Rovbase, had a strong positive effect on detection probability during unstructured sampling (2021/2022; males:  $\beta = 0.48$ , CrI: 0.22 - 0.76; females:  $\beta = 0.61$ , CrI: 0.29 - 0.94; Table A.9). Detection of an individual during the previous year also tended to have a positive effect on detection probability during unstructured sampling but the pattern was not consistent across years (Table A.9).

## 4 Summary of improvements made

The analysis described in this report includes the following adjustments compared with previous analyses of wolverine density in Scandinavia by RovQuant (Bischof et al., 2020b; Flagstad et al., 2021; Milleret et al., 2021b):

1. Addition of data from the 2022 monitoring season. We also excluded the 2013 winter monitoring season, as this enhanced model performance.
2. Due to the uncertainty about the dead recovery process, we excluded dead recoveries from this analysis.
3. Replaced the half-normal movement model with the exponential for modelling inter-annual movement. We found in an ongoing study that the latter is better suited for capturing the inter-annual movement patterns in populations that include long-distance dispersing individuals.
4. Used separate detection sub-models for structured and unstructured sampling to account for fundamental differences in how samples accumulate and better represent variation in search effort.
5. Allowed for annual variation in all covariates in the detection sub-model (i.e., covariates on  $p_0$  and  $\sigma$  are time-dependent).
6. Used carnivore observation reports in Skandobs and ancillary carnivore samples recorded in Rovbase to generate a new detection covariate that serves as a proxy for sampling effort (spatially and temporally varying) during unstructured sampling.

## 5 Suggestions for future improvements

RovQuant continues to work on improving the functionality and efficiency of OPSCR models. We plan to test and potentially implement the following developments in future analyses of the Scandinavian wolverine monitoring data:

1. Review and adjust spatial covariates on density. This may involve the addition of relevant spatial variables (Moqanaki et al., 2022).
2. Account for individual heterogeneity in detection for example by using a finite-mixture approach (Cubaynes et al., 2010).
3. Consider alternative detection models that do not assume a half-normal shape and/or circular home ranges (Sutherland et al., 2015; Dey et al., 2022).
4. Account for spatial variation in survival (Milleret et al., 2022a).

## 6 Other recommendations

In addition, we suggest the following:

1. Indicate sample association with the search tracks in Rovbase (if any) to unambiguously identify samples arising from structured vs unstructured sampling.
2. Consider full-coverage NGS in all regions for which estimates are desired (e.g., Norbotten or reindeer (*Rangifer tarandus*) herding areas).

3. Report information about how samples are selected for DNA analysis.
4. Record coarse measures of search conditions at the search track level (e.g., presence/absence of snow, days since last snowfall, experience level of searchers).
5. Unambiguously and consistently indicate the species targeted during searches when recording GPS search tracks.
6. Clearly identify and delineate areas excluded from structured and unstructured sampling and indicate the reason for exclusion (e.g., unable to search the area or low priority due to assumed lack of presence of the target species).
7. Explore the feasibility of using station-based detectors (e.g., hair snares or similar) for better control over the observation process.

## 7 Acknowledgements

This work was made possible by the large carnivore monitoring programs and the extensive monitoring data collected by Swedish (Länstyrelsen) and Norwegian (SNO) wildlife management authorities, as well as the public in both countries. Our analyses relied on genetic analyses conducted by the laboratory personnel at the DNA laboratories at the Swedish University of Agricultural Sciences, the Norwegian Institute for Nature Research, and Uppsala University. We also thank Swedish and Norwegian wildlife managers for feedback provided during project RovQuant and the Research Council of Norway for partial funding (NFR 286886; project WildMap). Computation was performed on resources provided by NMBU's computing cluster Orion, administered by the Centre for Integrative Genetics and by UNINETT Sigma2 - the National Infrastructure for High Performance Computing and Data Storage in Norway. We are grateful to P. de Valpine and D. Turek for help with the formulation of the OPSCR model in Nimble. J. Vermaat provided helpful comments on a draft of this report.

## 8 Data availability


Data, R code to reproduce the analysis, as well as figures, tables, and raster maps (Figure A.4) are available at <https://github.com/richbi/RovQuantPublic>.

## References


- Bischof, R. (2015). Approaches for assessing illegal hunting of brown bears and other large carnivores in Sweden. *Report 2015:5 from the Scandinavian Brown Bear Research Project to the Swedish Environmental Protection Agency. Norwegian University of Life Sciences, Ås, Norway.*
- Bischof, R., Brøseth, H., and Gimenez, O. (2016). Wildlife in a Politically Divided World: Insularism Inflates Estimates of Brown Bear Abundance. *Conservation Letters*, 9(2):122–130.
- Bischof, R., Dupont, P., Milleret, C., Chipperfield, J., and Royle, J. A. (2020a). Consequences of ignoring group association in spatial capture–recapture analysis. *Wildlife Biology*, 2020(1):1–10.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Åkesson, M., Brøseth, H., and Kindberg, J. (2019a). Estimating the size of the Scandinavian wolf population with spatial capture recapture and conversion factors. *MINAfaagrapport 57*, 80pp.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Brøseth, H., and Kindberg, J. (2019b). Estimating density, abundance and population dynamics of bears, wolverines, and wolves in Scandinavia. *MINAfaagrapport 63*, 79pp.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Tourani, M., Ordiz, A., de Valpine, P., Turek, D., Royle, J. A., Gimenez, O., Flagstad, , Åkesson, M., Svensson, L., Brøseth, H., and Kindberg, J. (2020b). Estimating and forecasting spatial population dynamics of apex predators using transnational genetic monitoring. *Proceedings of the National Academy of Sciences*, 117(48):30531–30538.
- Bischof, R., Turek, D., Milleret, C., Ergon, T., Dupont, P., Dey, S., and de Valpine, P. (2021). *nimbleSCR: Spatial Capture-Recapture (SCR) Methods Using 'nimble'*. R package version 0.1.2.
- Chandler, R. B., Hepinstall-Cymerman, J., Merker, S., Abernathy-Conners, H., and Cooper, R. J. (2018). Characterizing spatio-temporal variation in survival and recruitment with integrated population models. *The Auk*, 135(3):409–426.
- Cubaynes, S., Pradel, R., Choquet, R., Duchamp, C., Gaillard, J. M., Lebreton, J. D., Marboutin, E., Miquel, C., Reboulet, A. M., Poillot, C., Taberlet, P., and Gimenez, O. (2010). Importance of accounting for detection heterogeneity when estimating abundance: the case of French wolves. *Conservation Biology*, 24(2):621–626.
- de Valpine, P., Paciorek, C., Turek, D., Michaud, N., Anderson-Bergman, C., Obermeyer, F., Wehrhahn Cortes, C., Rodríguez, A., Temple Lang, D., and Paganin, S. (2022). *NIMBLE User Manual*. R package manual version 0.12.2.
- de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., and Bodik, R. (2017). Programming with models: writing statistical algorithms for general model structures with NIMBLE. *Journal of Computational and Graphical Statistics*, 26(2):403–413.
- Dey, S., Bischof, R., Dupont, P. P. A., and Milleret, C. (2022). Does the punishment fit the crime? consequences and diagnosis of misspecified detection functions in Bayesian spatial capture–recapture modeling. *Ecology and Evolution*, 12(2):e8600.
- Dupont, P., Milleret, C., Tourani, M., Brøseth, H., and Bischof, R. (2021). Integrating dead recoveries in open-population spatial capture–recapture models. *Ecosphere*, 12(7):e03571.
- Efford, M. (2004). Density estimation in live-trapping studies. *Oikos*, 106(3):598–610.
- Ergon, T. and Gardner, B. (2014). Separating mortality and emigration: modelling space use, dispersal and survival with robust-design spatial capture–recapture data. *Methods in Ecology and Evolution*, 5(12):1327–1336.
- Flagstad, , Hedmark, E., Landa, A., Brøseth, H., Persson, J., Andersen, R., Segerström, P., and Ellegren, H. (2004). Colonization history and noninvasive monitoring of a reestablished wolverine population. *Conservation Biology*, 18(3):676–688.
- Flagstad, , Kleven, O., Brandsegg, H., Spets, M., Eriksen, L., Andersskog, I., Johansson, M., Ekblom, R., Ellegren, H., and Brøseth, H. (2021). DNA-basert overvåking av den skandinaviske jervebestanden 2020. *Norsk institutt for naturforskning(NINA), Trondheim*, NINA Rapport nr 1956.
- Gardner, B., Sollmann, R., Kumar, N. S., Jathanna, D., and Karanth, K. U. (2018). State space and movement specification in open population spatial capture–recapture models. *Ecology and Evolution*, 8(20):10336–10344.
- Gelman, A. and Rubin, D. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7:457–511. <http://www.stat.columbia.edu/~gelman/research/published/itsim.pdf>.
- Kéry, M. and Schaub, M. (2012). *Bayesian population analysis using WinBUGS: a hierarchical perspective*. Academic Press, Waltham, MA.
- Kleven, O., Andersskog, I., Brandsegg, H., Eriksen, L., Spets, M., Königsson, H., Spong, a., Milleret, C., Dupont, P., Bischof, R., Flagstad, , and Brøseth, H. (2022a). DNA-basert overvåking av den skandinaviske jervebestanden 2021. *Norsk institutt for naturforskning(NINA), Trondheim*, NINA Rapport nr 2111.
- Kleven, O., Forfang, K., Opsahl, N. N., Königsson, H., Spong, G., Milleret, C., Dupont, P., Bischof, R., Flagstad, , and Brøseth, H. (2022b). DNA-based monitoring of the scandinavian wolverine population 2022. *Norsk institutt for naturforskning(NINA), Trondheim*, NINA Rapport nr 2236.

- Lebreton, J.-D. and Pradel, R. (2002). Multistate recapture models: modelling incomplete individual histories. *Journal of Applied Statistics*, 29:353–369.
- Milleret, C., Bischof, R., Dupont, P., Brøseth, H., Odden, J., and Mattisson, J. (2021a). GPS collars have an apparent positive effect on the survival of a large carnivore. *Biology Letters*, 17(6):20210128.
- Milleret, C., Dey, S., Dupont, P., Brøseth, H., Turek, D., de Valpine, P., and Bischof, R. (2022a). Estimating spatially variable and density-dependent survival using open-population spatial capture–recapture models. *Ecology*, n/a(n/a):e3934. e3934 ECY22-0819.
- Milleret, C., Dupont, P., Bonenfant, C., Brøseth, H., Flagstad, O., Sutherland, C., and Bischof, R. (2019). A local evaluation of the individual state-space to scale up Bayesian spatial capture recapture. *Ecology and Evolution*, 9(1):352–363.
- Milleret, C., Dupont, P., Brøseth, H., Flagstad, O., Kindberg, J., and Bischof, R. (2022b). Estimates of wolverine density, abundance, and population dynamics in Scandinavia, 2013–2021. *MINAflagrapport 74*, 30pp.
- Milleret, C., Dupont, P., Brøseth, H., Kindberg, J., Royle, J. A., and Bischof, R. (2018). Using partial aggregation in spatial capture recapture. *Methods in Ecology and Evolution*, 9(8):1896–1907.
- Milleret, C., Dupont, P., Åkesson, M., Brøseth, H., Kindberg, J., and Bischof, R. (2021b). Estimates of wolf density, abundance, and population dynamics in Scandinavia, 2012 - 2021. *MINAflagrapport 72*, 30pp.
- Milleret, C., Dupont, P., Åkesson, M., Svensson, L., Brøseth, H., Kindberg, J., and Bischof, R. (2022c). Estimates of wolf density, abundance, and population dynamics in Scandinavia, 2013–2022. *MINAflagrapport 77*, 35pp.
- Moqanaki, E., Milleret, C., Dupont, P., Brøseth, H., and Bischof, R. (2022). Wolverine density distribution reflects past persecution and current management in Scandinavia. *bioRxiv*.
- R Core Team (2021). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Royle, J. A., Chandler, R. B., Sollmann, R., and Gardner, B. (2013). *Spatial capture-recapture*. Elsevier/Academic Press, Amsterdam.
- Royle, J. A. and Dorazio, R. M. (2012). Parameter-expanded data augmentation for Bayesian analysis of capture–recapture models. *Journal of Ornithology*, 152(2):521–537.
- Schaub, M. and Royle, J. A. (2014). Estimating true instead of apparent survival using spatial Cormack–Jolly–Seber models. *Methods in Ecology and Evolution*, 5(12):1316–1326.
- Sutherland, C., Fuller, A. K., and Royle, J. A. (2015). Modelling non-Euclidean movement and landscape connectivity in highly structured ecological networks. *Methods in Ecology and Evolution*, 6(2):169–177.
- Turek, D., Milleret, C., Ergon, T., Brøseth, H., Dupont, P., Bischof, R., and de Valpine, P. (2021). Efficient estimation of large-scale spatial capture–recapture models. *Ecosphere*, 12(2):e03385.
- Wabakken, P., Svensson, L., Maartmann, E., Nordli, K., Flagstad, O., and Åkesson, M. (2022). Bestandsovervåking av ulv vinteren 2021-2022. inventering av varg vinteren 2021-2022. *Bestandsstatus for store rovdyr i Skandinavia. Bestandsstatus för stora rovdjur i Skandinavien 1-2022*, 1(59s).
- Zhang, W., Chipperfield, J. D., Illian, J. B., Dupont, P., Milleret, C., de Valpine, P., and Bischof, R. (2022). A flexible and efficient bayesian implementation of point process models for spatial capture-recapture data. *Ecology*, page e3887. e3887.

# Appendices




**Figure A.1:** Covariate used to account for spatio-temporal variation in unstructured sampling in the study area. Green cells ( $10 \times 10$  km) represent areas with at least one carnivore record from Rovbase (rovbase.no, rovbase.se, excluding the wolverine samples used in the OPSCR model) or an observation record from Skandobs (skandobs.se, skandobs.no) during a given monitoring season (Dec 1 – Jun 30).




**Figure A.2:** Management jurisdictions in Norway and Sweden. Polygons represent carnivore management regions in Norway and counties in Sweden. Thick outlines delineate Swedish carnivore management regions ("Rovdjursförvaltningsområden") encompassing multiple counties.







**Figure A.3:** Total (black) and country-specific (blue: Sweden, red: Norway) annual wolverine population size estimates in Scandinavia between 2014 and 2022. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Credible intervals indicate uncertainty in estimates given the model and data used to generate the estimates. Changes in the model and data can result in different estimates and associated uncertainty compared with estimates provided in previous reports by RovQuant.




**Figure A.4:** Wolverine density based on individual utilization distributions in Scandinavia between 2014 and 2022. Note that no comprehensive NGS was conducted in Norrbotten (polygon outlined in black) from 2014-2016 and from 2020-2022, which means that the results are solely based on the OPSCR model prediction and assumption. The grey area represents areas that were considered not searched and therefore were not included in the analysis. These maps are freely available as geo-referenced raster files at <https://github.com/richbi/RovQuantPublic>.



**Figure A.5:** Annual survival probabilities for female and male wolverines. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Shown are overall estimates for the entire study area between 2014 and 2022.



**Figure A.6:** Sex-specific baseline detection probabilities ( $p_{0,structured}$ ) for the different Scandinavian jurisdictions during structured sampling as estimated by the open-population spatial capture-recapture (OPSCR) model. Bars represent 95% credible intervals. Results are separated into panels based on regions. Estimates are shown for the mean values of the detection covariates. Note that baseline detection probability ( $p_0$ ) is a theoretical value of detection probability when a detector coincides with the location of an individual's AC; it is not to be confused with detectability, i.e., the overall probability of detecting an individual.



**Figure A.7:** Sex- and country-specific baseline detection probabilities ( $p_{0unstructured}$ ) of wolverines during unstructured sampling as estimated by the open-population spatial capture-recapture (OPSCR) model. Bars represent 95% credible intervals. Estimates are shown for the mean values of the detection covariates. Note that baseline detection probability ( $p_0$ ) is a theoretical value of detection probability when a detector coincides with the location of an individual's activity center; it is not to be confused with detectability, i.e., the overall probability of detecting an individual.

**Table A.1:** Annual number of wolverine non-invasive genetic samples included in the analysis. Numbers are reported by country, for females (F) and males (M), and for each type of sampling (structured and unstructured). We included only samples collected within the study area during the primary monitoring period (Dec 1 - June 30) between 2014 and 2022.

		2014		2015		2016		2017		2018		2019		2020		2021		2022	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
<b>Norway</b>	Structured	341	375	244	247	304	378	377	423	277	471	377	506	379	453	390	480	411	520
	Unstructured	180	203	168	198	164	201	229	247	185	276	215	225	196	231	249	259	157	227
<b>Sweden</b>	Structured	87	102	131	128	131	158	276	319	343	509	367	477	250	351	286	351	327	475
	Unstructured	99	52	97	93	105	117	211	233	293	330	137	141	105	187	174	211	156	173
<b>Total</b>	Structured	428	477	375	375	435	536	653	742	620	980	744	983	629	804	676	831	738	995
	Unstructured	279	255	265	291	269	318	440	480	478	606	352	366	301	418	423	470	313	400

**Table A.2:** Annual number of individual wolverines detected via non-invasive genetic sampling and included in the analysis. Numbers are reported by country, for females (F) and males (M), and for each type of sampling (structured and unstructured). We included only individuals associated with samples collected within the study area during the primary monitoring period (Dec 1 - Jun 30) between 2013/2014 and 2021/2022. Some individuals were detected in both countries during the same year, hence the sum of the national counts can exceed the total number of unique individuals detected in Scandinavia.

		2014		2015		2016		2017		2018		2019		2020		2021		2022	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
<b>Norway</b>	Structured	141	109	111	90	133	96	143	100	127	105	147	124	154	121	149	112	161	108
	Unstructured	102	91	93	78	106	91	109	100	103	87	108	93	107	97	102	91	95	93
<b>Sweden</b>	Structured	56	50	81	61	74	80	156	131	165	172	187	176	131	130	155	134	154	157
	Unstructured	45	33	57	52	49	60	115	119	143	128	89	84	72	88	98	91	72	92
<b>Total</b>	Structured	196	158	188	148	206	175	287	217	286	264	327	291	282	248	301	241	314	257
	Unstructured	145	124	149	126	155	148	220	208	243	207	196	176	178	182	200	179	167	184

**Table A.3:** Number of cause-specific dead recoveries of wolverines in Scandinavia between 2014 and 2022. Numbers are reported by country, for females (F) and males (M). Dead recovery data was only used to derive cause-specific mortality (Figure 2).

		Country		2014		2015		2016		2017		2018		2019		2020		2021		2022	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
<b>Other</b>	Norway	1	5	0	2	3	1	3	2	7	1	1	3	3	3	4	4	1	0		
	Sweden	1	3	1	2	0	2	0	1	0	1	1	1	0	1	1	1	1	2	1	
<b>Legal culling</b>	Norway	42	34	69	52	42	43	47	61	27	31	57	37	49	63	44	44	28	35		
	Sweden	13	8	22	18	6	9	5	2	2	5	6	3	7	19	3	10	5	1		
<b>Total</b>	Total	57	50	92	74	51	55	55	66	36	38	65	44	59	86	52	59	36	37		

**Table A.4:** Annual abundance estimates for wolverine at several spatial scales: the entire study area, by country, by management unit (carnivore management regions in Norway and "Rovdjursförvaltningsområden" in Sweden), and counties ("Län" in Sweden); see also Figure A.2. Only counties and management units that are within or that intersect the study area are included in the table. Estimates are based on model-estimated activity center locations. Credible intervals (95%) are shown in parentheses. Small deviations between the total estimate and the sum of abundance estimates from the constituent sub-regions may arise due to rounding. Note that the numbers reported here are predictions from a statistical model which always represents an oversimplification of reality and is based on available data (NGS). As a consequence, especially at the local scale, the model-estimated number of wolverines based on DNA sampling can deviate from the number of wolverines inferred from ancillary observations (e.g., camera traps). Estimates for Norrbotten county in years without comprehensive NGS were derived solely using the prediction from the OPSCR model (shown in grey and marked with \*). Total estimates in Sweden and for the entire study area that includes estimates from Norrbotten without comprehensive NGS are marked with \*\*.

	2014	2015	2016	2017	2018	2019	2020	2021	2022
TOTAL	970.3 (887-1065)**	827.6 (777-886)**	924.9 (876-975)**	905.6 (877-941)	952.6 (925-983)	1063 (1027-1101)	1077.6 (1033-1124)**	1068.4 (1028-1115)**	1035.2 (980-1088)**
NORWAY	427.7 (390-467)	351 (328-376)	379.7 (357-403)	355 (340-373)	360 (344-376)	411.8 (394-430)	400.6 (383-420)	368.8 (352-388)	367.2 (349-391)
Region 1	19.5 (12-27)	11.2 (7-16)	9.2 (5-14)	7.3 (4-12)	6.1 (3-10)	5.9 (2-10)	5.3 (2-10)	6.3 (2-11)	10.4 (6-16)
Region 2	8 (3-14)	3 (0-7)	3.1 (0-6)	2.7 (1-6)	2 (0-5)	2.2 (0-6)	2.3 (0-6)	4.3 (1-8)	5.1 (1-9)
Region 3	35.8 (29-43)	26.8 (22-32)	26.6 (22-32)	23.4 (19-28)	22.5 (18-27)	27.9 (23-33)	30.4 (26-35)	32.7 (28-38)	27 (21-33)
Region 4	2.6 (0-6)	1.1 (0-3)	0.8 (0-3)	0.7 (0-3)	0.5 (0-2)	0.6 (0-2)	0.9 (0-3)	0.8 (0-3)	1.2 (0-4)
Region 5	62.9 (55-72)	53.9 (47-62)	63.4 (57-70)	63.3 (57-70)	76.5 (70-84)	91.7 (85-98)	88 (81-96)	82.4 (75-90)	81 (75-89)
Region 6	82.9 (71-96)	70.5 (61-81)	80.7 (72-91)	79.8 (73-88)	90.6 (83-99)	102.3 (94-111)	85.4 (77-95)	79.2 (71-87)	91.7 (84-100)
Region 7	78.5 (69-89)	71.2 (64-79)	82.3 (75-90)	70.1 (64-76)	66.9 (62-74)	78.5 (72-86)	88.2 (81-95)	69.4 (64-75)	66.7 (61-74)
Region 8	137.5 (123-152)	113.3 (103-125)	113.5 (103-124)	107.7 (101-115)	94.9 (87-103)	102.6 (94-112)	100 (93-109)	93.7 (85-103)	84.1 (75-96)
SWEDEN	542.6 (484-608)**	476.7 (440-521)**	545.2 (506-581)**	550.5 (525-579)	592.6 (571-616)	651.2 (620-683)	677 (635-719)**	699.6 (663-740)**	667.9 (625-709)**
Norra	476.2 (427-536)	425 (391-464)	484.2 (448-519)	481.6 (460-505)	509.5 (490-530)	551.8 (523-580)	572.1 (536-610)	594.2 (559-630)	557.6 (517-598)
Jämtland	147.8 (126-171)	139.4 (125-155)	168.3 (155-184)	172.1 (159-187)	191.9 (180-205)	211.7 (198-227)	215.1 (200-230)	220.6 (207-233)	194.1 (180-211)
Norrbotten	181.6 (153-213)*	158 (134-182)*	173 (151-196)*	170.5 (159-183)	173.8 (163-185)	171 (154-188)	181.6 (158-206)*	188.8 (162-214)*	187.3 (158-216)*
Västerbotten	126.3 (112-144)	109.5 (100-121)	122.8 (111-135)	113.9 (104-124)	109 (99-119)	128.9 (118-141)	130.9 (120-143)	135.5 (124-146)	117 (104-129)
Västernorrland	20.6 (13-30)	18.1 (12-25)	20.1 (14-27)	25.1 (19-32)	34.8 (29-41)	40.1 (33-47)	44.5 (37-53)	49.3 (42-57)	59.2 (51-67)
Mellersta	65.8 (51-82)	51.2 (40-64)	60.6 (50-71)	68.6 (58-78)	82.8 (74-92)	99.1 (90-109)	104.6 (94-117)	105 (93-116)	109.9 (99-122)
Dalarna	30.4 (22-41)	26.3 (20-33)	30.8 (24-38)	35.4 (29-42)	46.4 (41-52)	48.4 (42-56)	50.1 (42-58)	53.2 (47-61)	54.9 (48-62)
Gävleborg	14.6 (7-22)	12.4 (7-18)	15 (10-21)	17.4 (12-24)	21.5 (17-27)	35.2 (30-41)	35.2 (30-42)	32.1 (27-38)	30.1 (25-36)
Örebro	4.9 (1-10)	2.5 (0-6)	2.1 (0-5)	1.7 (0-5)	2 (0-5)	1.5 (0-4)	2.5 (0-6)	2.7 (0-6)	2.8 (0-7)
Uppsala	0.3 (0-2)	0.1 (0-1)	0.2 (0-1)	0.1 (0-1)	0.1 (0-1)	0.2 (0-1)	0.1 (0-1)	0.1 (0-1)	0.1 (0-1)
Värmland	12 (7-19)	7.8 (4-13)	10.7 (7-15)	12.5 (9-17)	11.5 (8-15)	12.4 (9-17)	14.8 (10-20)	15.2 (11-20)	20.1 (16-25)
Västmanland	1.9 (0-5)	1.1 (0-3)	0.9 (0-3)	0.7 (0-3)	0.6 (0-3)	0.6 (0-2)	0.9 (0-3)	0.9 (0-3)	0.9 (0-3)
VästraGötaland	1.7 (0-5)	1 (0-3)	0.9 (0-3)	0.8 (0-3)	0.6 (0-2)	0.7 (0-3)	0.9 (0-3)	0.8 (0-3)	0.9 (0-3)
Södra	0.6 (0-2)	0.4 (0-2)	0.4 (0-2)	0.4 (0-2)	0.3 (0-2)	0.4 (0-2)	0.4 (0-2)	0.4 (0-2)	0.4 (0-2)
Östergötland	0.3 (0-2)	0.2 (0-1)	0.2 (0-1)	0.2 (0-1)	0.1 (0-1)	0.2 (0-1)	0.2 (0-1)	0.2 (0-1)	0.2 (0-2)
Södermanland	0.3 (0-2)	0.2 (0-1)	0.2 (0-1)	0.2 (0-1)	0.1 (0-1)	0.2 (0-1)	0.2 (0-1)	0.2 (0-2)	0.2 (0-1)



**Table A.5:** Annual population growth rate estimates for the wolverine population in Scandinavia ("Total") and separately for Norway and Sweden. Estimates were derived using the posterior distributions of annual abundance estimates (Table A.4). Credible intervals (95%) are shown in parentheses.

	2014-2015	2015-2016	2016-2017	2017-2018	2018-2019	2019-2020	2020-2021	2021-2022
Norway	0.82 (0.75-0.90)	1.08 (0.99-1.16)	0.94 (0.88-1.00)	1.01 (0.95-1.08)	1.14 (1.07-1.22)	0.97 (0.92-1.03)	0.92 (0.87-0.98)	1.00 (0.93-1.07)
Sweden	0.88 (0.79-0.98)	1.15 (1.05-1.26)	1.01 (0.94-1.10)	1.08 (1.02-1.14)	1.10 (1.04-1.17)	1.04 (0.98-1.12)	1.03 (0.97-1.10)	0.96 (0.89-1.02)
Total	0.85 (0.78-0.94)	1.12 (1.04-1.20)	0.98 (0.92-1.04)	1.05 (1.01-1.10)	1.12 (1.06-1.17)	1.01 (0.96-1.07)	0.99 (0.95-1.04)	0.97 (0.91-1.03)

**Table A.6:** Estimates of the demographic parameters obtained from the sex-specific wolverine open-population spatial capture-recapture (OPSCR) models. Parameters represent transition rates from Dec 1 to Nov 30 in the following year. Median estimates and 95% credible intervals (in parentheses) for per capita recruitment rate ( $\rho$ ), survival ( $\phi$ ), mortality due to legal culling ( $h$ ) and mortality due to other causes ( $w$ ) are presented for males (M) and females (F). Note that mortality due to legal culling was not estimated in the model, but derived from the recorded number of dead recoveries.

	2014-2015		2015-2016		2016-2017		2017-2018	
	M	F	M	F	M	F	M	F
$\rho$	0.26 (0.17-0.37)	0.15 (0.06-0.26)	0.44 (0.33-0.54)	0.35 (0.25-0.47)	0.35 (0.27-0.43)	0.19 (0.11-0.26)	0.41 (0.35-0.46)	0.27 (0.21-0.32)
$\phi$	0.56 (0.49-0.64)	0.71 (0.64-0.77)	0.64 (0.57-0.71)	0.78 (0.72-0.83)	0.65 (0.59-0.71)	0.77 (0.71-0.82)	0.66 (0.60-0.71)	0.76 (0.72-0.81)
$h$	0.06 (0.06-0.07)	0.06 (0.06-0.07)	0.15 (0.14-0.17)	0.10 (0.09-0.11)	0.11 (0.10-0.12)	0.06 (0.05-0.06)	0.12 (0.11-0.13)	0.06 (0.06-0.06)
$w$	0.37 (0.31-0.43)	0.23 (0.17-0.28)	0.20 (0.15-0.26)	0.12 (0.07-0.17)	0.23 (0.19-0.28)	0.18 (0.13-0.22)	0.22 (0.19-0.25)	0.17 (0.14-0.21)
	2018-2019		2019-2020		2020-2021		2021-2022	
	M	F	M	F	M	F	M	F
$\rho$	0.41 (0.36-0.48)	0.29 (0.24-0.36)	0.38 (0.32-0.44)	0.25 (0.19-0.32)	0.35 (0.30-0.41)	0.24 (0.18-0.30)	0.37 (0.32-0.44)	0.20 (0.15-0.25)
$\phi$	0.69 (0.64-0.75)	0.82 (0.77-0.87)	0.62 (0.57-0.68)	0.76 (0.70-0.81)	0.61 (0.56-0.67)	0.76 (0.70-0.82)	0.61 (0.55-0.67)	0.75 (0.68-0.83)
$h$	0.08 (0.07-0.08)	0.04 (0.04-0.04)	0.08 (0.07-0.08)	0.07 (0.06-0.07)	0.15 (0.14-0.16)	0.06 (0.05-0.06)	0.09 (0.09-0.10)	0.05 (0.05-0.05)
$w$	0.23 (0.20-0.26)	0.14 (0.10-0.18)	0.30 (0.27-0.34)	0.18 (0.13-0.22)	0.23 (0.19-0.27)	0.18 (0.13-0.23)	0.29 (0.26-0.33)	0.20 (0.13-0.26)

**Table A.7:** Estimates of the density and movement process parameters obtained from the sex-specific wolverine open-population spatial capture-recapture (OPSCR) models.  $\beta_{dens}$  represents the effect of the number of known wolverine dens on activity center locations (Bischof et al., 2020b). The scale parameter  $\sigma$  of the detection function is expressed in kilometers and estimated separately for each year.  $\lambda$  (in km) represents the mean of the exponential movement parameter, describing individual movement distances between years. Credible intervals (95%) are shown in parentheses. Parameters that were not estimated separately each year are marked with \*.

	2014		2015		2016		2017		2018	
	M	F	M	F	M	F	M	F	M	F
$\beta_{dens}^*$	0.44 (0.40-0.47)	0.48 (0.45-0.52)	0.44 (0.40-0.47)	0.48 (0.45-0.52)	0.44 (0.40-0.47)	0.48 (0.45-0.52)	0.44 (0.40-0.47)	0.48 (0.45-0.52)	0.44 (0.40-0.47)	0.48 (0.45-0.52)
$\sigma$	8.33 (7.91-8.79)	5.86 (5.52-6.25)	8.48 (8.05-8.93)	5.63 (5.32-5.98)	8.23 (7.86-8.63)	6.09 (5.77-6.45)	8.39 (8.08-8.73)	6.54 (6.27-6.82)	8.00 (7.74-8.26)	6.24 (5.98-6.51)
$\lambda^*$	14.63 (14.02-15.35)	7.99 (7.69-8.31)	14.63 (14.02-15.35)	7.99 (7.69-8.31)	14.63 (14.02-15.35)	7.99 (7.69-8.31)	14.63 (14.02-15.35)	7.99 (7.69-8.31)	14.63 (14.02-15.35)	7.99 (7.69-8.31)
	2019		2020		2021		2022			
	M	F	M	F	M	F	M	F		
$\beta_{dens}^*$	0.44 (0.40-0.47)	0.48 (0.45-0.52)	0.44 (0.40-0.47)	0.48 (0.45-0.52)	0.44 (0.40-0.47)	0.48 (0.45-0.52)	0.44 (0.40-0.47)	0.48 (0.45-0.52)		
$\sigma$	7.73 (7.45-8.03)	5.63 (5.39-5.88)	7.98 (7.68-8.31)	6.25 (5.96-6.55)	8.09 (7.80-8.40)	6.30 (6.02-6.60)	8.07 (7.79-8.38)	5.70 (5.43-5.99)		
$\lambda^*$	14.63 (14.02-15.35)	7.99 (7.69-8.31)	14.63 (14.02-15.35)	7.99 (7.69-8.31)	14.63 (14.02-15.35)	7.99 (7.69-8.31)	14.63 (14.02-15.35)	7.99 (7.69-8.31)		

**Table A.8:** Estimates of the detection process parameters for the structured sampling.  $\beta_{1Structured}$  corresponds to the effect of previous detection of an individual,  $\beta_{2Structured}$  to the effect of search-effort (track length), and  $\beta_{3Structured}$  to the effect of average snow cover during the monitoring period on baseline detection probability ( $p_{0Structured}$ ). Coefficients are associated with scaled covariates. Credible intervals (95%) are shown in parentheses.

	2014		2015		2016		2017		2018	
	M	F	M	F	M	F	M	F	M	F
$\beta_{1Structured}$	0.91 (0.66-1.16)	0.61 (0.35-0.87)	0.50 (0.23-0.77)	0.03 (-0.24-0.31)	0.53 (0.31-0.76)	0.27 (0.01-0.52)	0.45 (0.26-0.64)	0.04 (-0.16-0.23)	0.54 (0.37-0.71)	0.11 (-0.09-0.31)
$\beta_{2Structured}$	0.48 (0.40-0.56)	0.39 (0.32-0.47)	0.44 (0.35-0.54)	0.43 (0.34-0.53)	0.43 (0.36-0.51)	0.49 (0.41-0.59)	0.44 (0.37-0.50)	0.39 (0.32-0.45)	0.42 (0.36-0.48)	0.41 (0.34-0.48)
$\beta_{3Structured}$	0.43 (0.13-0.75)	0.50 (0.19-0.83)	-0.02 (-0.23-0.19)	0.20 (-0.02-0.42)	0.22 (0.00-0.43)	0.10 (-0.13-0.33)	0.32 (0.14-0.50)	0.20 (0.02-0.37)	0.12 (0.00-0.24)	0.04 (-0.09-0.18)
	2019		2020		2021		2022			
	M	F	M	F	M	F	M	F		
$\beta_{1Structured}$	0.57 (0.39-0.75)	0.14 (-0.06-0.33)	0.17 (0.00-0.36)	-0.01 (-0.21-0.19)	0.48 (0.29-0.67)	-0.15 (-0.36-0.06)	0.42 (0.23-0.60)	0.06 (-0.15-0.28)		
$\beta_{2Structured}$	0.47 (0.40-0.55)	0.49 (0.41-0.57)	0.43 (0.37-0.49)	0.46 (0.39-0.53)	0.50 (0.43-0.58)	0.55 (0.48-0.63)	0.44 (0.37-0.52)	0.42 (0.35-0.49)		
$\beta_{3Structured}$	-0.09 (-0.24-0.06)	-0.06 (-0.22-0.12)	0.34 (0.16-0.52)	0.33 (0.10-0.56)	0.13 (-0.01-0.27)	0.24 (0.08-0.40)	0.08 (-0.07-0.22)	0.08 (-0.08-0.24)		

**Table A.9:** Estimates of the detection process parameters for the unstructured sampling.  $\beta_{1Unstructured}$  corresponds to the effect of previous detection,  $\beta_{2Unstructured}$  to the effect of distance to the nearest road,  $\beta_{3Unstructured}$  to the effect of average snow cover during the monitoring period, and  $\beta_{4Unstructured}$  to the effect of spatio-temporal heterogeneity in unstructured sampling derived using the observation data in Skandobs and Rovbase on baseline detection probability ( $p_{0Unstructured}$ ). Coefficients are associated with scaled covariates. Credible intervals (95%) are shown in parentheses.

	2014		2015		2016		2017		2018	
	M	F	M	F	M	F	M	F	M	F
$\beta_{1Unstructured}$	0.73 ( 0.43-1.03)	0.41 ( 0.10-0.73)	0.16 (-0.13-0.45)	0.13 (-0.19-0.43)	0.56 ( 0.29-0.83)	0.37 ( 0.06-0.67)	0.25 ( 0.03-0.46)	-0.16 (-0.39-0.07)	0.85 ( 0.64-1.06)	0.22 ( 0.00-0.45)
$\beta_{2Unstructured}$	0.56 ( 0.25-0.88)	0.61 ( 0.31-0.94)	0.29 ( 0.09-0.51)	0.44 ( 0.22-0.67)	-0.03 (-0.22-0.17)	0.30 ( 0.08-0.52)	0.42 ( 0.23-0.61)	0.36 ( 0.17-0.56)	0.18 ( 0.04-0.32)	0.14 (-0.01-0.30)
$\beta_{3Unstructured}$	0.09 (-0.08-0.26)	0.18 ( 0.02-0.35)	0.02 (-0.17-0.21)	0.13 (-0.05-0.31)	0.04 (-0.15-0.22)	-0.12 (-0.32-0.06)	0.13 ( 0.04-0.21)	0.15 ( 0.06-0.23)	0.02 (-0.07-0.10)	0.04 (-0.04-0.13)
$\beta_{4Unstructured}$	0.74 (0.44-1.04)	0.53 (0.24-0.83)	0.48 (0.20-0.76)	0.73 (0.41-1.04)	0.77 (0.50-1.05)	0.30 (0.01-0.59)	0.58 (0.36-0.80)	0.55 (0.32-0.78)	0.61 (0.40-0.83)	0.82 (0.58-1.07)
	2019		2020		2021		2022			
	M	F	M	F	M	F	M	F		
$\beta_{1Unstructured}$	0.55 ( 0.29-0.81)	0.08 (-0.17-0.35)	0.32 ( 0.09-0.56)	-0.05 (-0.32-0.22)	0.63 ( 0.40-0.86)	0.40 ( 0.14-0.66)	0.88 ( 0.63-1.15)	0.14 (-0.14-0.43)		
$\beta_{2Unstructured}$	0.06 (-0.14-0.26)	0.20 (-0.01-0.41)	0.29 ( 0.09-0.50)	0.61 ( 0.32-0.93)	0.18 ( 0.02-0.34)	0.01 (-0.17-0.20)	0.19 ( 0.01-0.37)	0.04 (-0.17-0.24)		
$\beta_{3Unstructured}$	0.02 (-0.11-0.15)	-0.02 (-0.15-0.10)	-0.11 (-0.30-0.06)	-0.18 (-0.38-0.00)	-0.12 (-0.29-0.05)	-0.06 (-0.24-0.10)	-0.14 (-0.34-0.04)	-0.12 (-0.33-0.08)		
$\beta_{4Unstructured}$	0.77 (0.50-1.06)	0.88 (0.59-1.18)	0.36 (0.12-0.61)	0.36 (0.09-0.64)	0.57 (0.33-0.84)	0.80 (0.51-1.11)	0.48 (0.21-0.76)	0.61 (0.29-0.95)		

**Table A.10:** Average proportion of individuals detected via non-invasive genetic sampling (NGS) in Sweden and Norway for males (M) and females (F). Values were calculated as the number of individuals detected with NGS (Table A.2) divided by the total and sex-specific abundance estimates obtained from the open-population spatial capture-recapture (OPSCR) models (Table A.4). Credible intervals (95%) are shown in parentheses. Note that in some years in Norway, male wolverines detected exceeded the estimated number of wolverines. This is possible when wolverine detection probability was very high and wolverines with activity centers in Sweden were detected on the Norwegian side of the border.

	2014		2015		2016		2017		2018	
	F	M	F	M	F	M	F	M	F	M
Norway	0.70 (0.62-0.79)	0.86 (0.78-0.96)	0.69 (0.63-0.75)	0.92 (0.85-0.99)	0.72 (0.67-0.78)	0.96 (0.89-1.02)	0.79 (0.74-0.83)	1.07 (1.01-1.14)	0.73 (0.69-0.77)	0.99 (0.94-1.04)
Sweden	0.26 (0.23-0.30)	0.32 (0.27-0.38)	0.37 (0.33-0.41)	0.46 (0.40-0.51)	0.32 (0.29-0.36)	0.50 (0.45-0.55)	0.64 (0.60-0.69)	0.80 (0.75-0.84)	0.67 (0.63-0.71)	0.86 (0.81-0.90)
Total	0.46 (0.41-0.51)	0.54 (0.48-0.60)	0.50 (0.45-0.54)	0.62 (0.57-0.67)	0.49 (0.45-0.53)	0.66 (0.62-0.71)	0.67 (0.64-0.71)	0.83 (0.80-0.87)	0.67 (0.64-0.70)	0.85 (0.82-0.88)
	2019		2020		2021		2022			
	F	M	F	M	F	M	F	M		
Norway	0.74 (0.69-0.79)	0.99 (0.93-1.04)	0.78 (0.73-0.83)	0.99 (0.93-1.06)	0.81 (0.76-0.87)	0.99 (0.93-1.05)	0.84 (0.77-0.91)	1.00 (0.93-1.05)		
Sweden	0.58 (0.54-0.61)	0.75 (0.70-0.80)	0.42 (0.38-0.45)	0.59 (0.55-0.64)	0.48 (0.45-0.52)	0.62 (0.58-0.67)	0.48 (0.44-0.52)	0.67 (0.62-0.72)		
Total	0.63 (0.60-0.66)	0.81 (0.77-0.84)	0.55 (0.51-0.58)	0.72 (0.69-0.76)	0.59 (0.55-0.62)	0.73 (0.69-0.77)	0.61 (0.56-0.66)	0.76 (0.71-0.80)		