

**Norwegian University of Life Sciences**

Faculty of Environmental Sciences and Natural Resource Management


**2023**

ISSN 2535-2806

MINA fagrapport 85

# **Estimates of wolf density, abundance, and population dynamics in Scandinavia, 2013–2023**

Cyril Milleret  
Pierre Dupont  
Henrik Brøseth  
Øystein Flagstad  
Jonas Kindberg  
Linn Svensson  
Richard Bischof



Milleret, C., Dupont, P., Brøseth, H., Flagstad, Ø., Kindberg, J., Svensson, L., and Bischof, R., 2023.  
**Estimates of wolf density, abundance, and population dynamics in Scandinavia, 2013–2023**  
- MINA fagrapport 85. 34 pp.

Ås, July 2023

ISSN: 2535-2806

COPYRIGHT

© Norwegian University of Life Sciences (NMBU)

The publication may be freely cited where the source is acknowledged

AVAILABILITY

Open

PUBLICATION TYPE

Digital document (pdf)

QUALITY CONTROLLED BY

The Research committee (FU), MINA, NMBU

PRINCIPAL

Naturvårdsverket, Ref: NV-02444-23, Contact person: Robert Ekblom

Miljødirektoratet, Contact person: Terje Bø

COVER PICTURE

Ridgeline plot of wolf density.

NØKKEORD

*Canis lupus*, ulv, tetthet, populasjonsdynamikk, deteksjonssannsynlighet, ikke-invaderende innsamling av genetisk materiale, åpen populasjon romlig fangst-gjenfangst, rovdyrforvaltning

KEY WORDS

*Canis lupus*, wolf, population density, population dynamics, detection probability, non-invasive genetic sampling, open-population spatial capture-recapture, carnivore management

Cyril Milleret, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Pierre Dupont, Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

Henrik Brøseth, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway.

Øystein Flagstad, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway

Jonas Kindberg, Norwegian Institute for Nature Research, PO Box 5685, NO-7485 Trondheim, Norway.

Linn Svensson, Grimsö Wildlife Research Station, Department of Ecology, Swedish University of Agricultural Sciences, Grimsö 152, SE-739 93 Riddarhyttan, Sweden


Richard Bischof ([richard.bischof@nmbu.no](mailto:richard.bischof@nmbu.no)), Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, PO Box 5003, NO-1432 Ås, Norway.

## Summary

**Background** The Scandinavian wolf (*Canis lupus*) population is being monitored annually using non-invasive genetic sampling (NGS) and recovery of dead individuals. DNA extracted from faeces, urine, hair, and tissue is used to identify the species, sex, and individual from which each sample originated. These data are compiled in the Scandinavian large carnivore database Rovbase 3.0.

**Approach** Using the Bayesian open-population spatial capture-recapture (OPSCR) model developed by RovQuant, we estimated annual density and vital rates of the Scandinavian wolf population for ten consecutive seasons from 2013/2014 to 2022/2023.

**Results** We generated annual density maps and estimated total and jurisdiction-specific population sizes for wolf from the winter 2013/2014 to 2022/2023 within the main population range. Based on the OPSCR model, the size of the Scandinavian wolf population was likely (95% credible interval) between 441 and 488 individuals in 2022/2023, with 376 to 415 individuals attributed to Sweden and 59 to 80 to Norway. In addition to annual density and jurisdiction-specific abundance estimates, we report annual estimates of cause-specific mortalities, recruitment, and detection probabilities.




Density map and ranges of abundance estimated for wolves in 2022/2023

## Sammendrag

**Bakgrunn** Den skandinaviske bestanden av ulv (*Canis lupus*) blir overvåket årlig ved bruk av ikke-invasiv genetisk prøveinnsamling (NGS) og gjenfunn av døde individer. DNA ekstrahert fra skit, urin, hår og vev brukes til å identifisere art, kjønn og individ for hver enkelt prøve. Denne informasjonen samles og ivaretas i den skandinaviske databasen for store rovdyr; Rovbase 3.0.

**Tilnærming** Ved bruk av en Bayesiansk åpen romlig fangst-gjenfangst populasjons modell (OPSCR), utviklet av RovQuant, estimerte vi årlige tettheter og demografiske rater hos den skandinaviske ulvebestanden i ti sesonger fra 2013/2014 til 2022/2023.

**Resultater** Innenfor hovedutbredelsesområdet til ulvebestanden laget vi årlige kart over tettheten av ulv fra 2013/2014 til 2022/2023, som viser både total bestandsstørrelse og bestandsstørrels innenfor ulike administrative enheter. Basert på OPSCR modellen var den skandinaviske ulvebestanden mellom 441 og 488 individer i 2022/2023 (95% kredibelt intervall), med 376 til 415 individer i Sverige og 59 til 80 individer i Norge. I tillegg til årlige tettheter og områdespesifikke bestandsestimater, gir rapporten estimater på dødlighetsfaktorer, rekruttering og oppdagbarhet.



Kart som viser tetthet av ulv i 2022/2023 sammen med intervaller for estimert antall ulv

# Contents

<b>1</b>	<b>Introduction</b>	<b>6</b>
<b>2</b>	<b>Methods</b>	<b>8</b>
2.1	Data . . . . .	8
2.2	Open-population spatial capture-recapture model . . . . .	9
<b>3</b>	<b>Results</b>	<b>13</b>
3.1	Non-invasive genetic samples and dead recoveries . . . . .	13
3.2	Density and abundance . . . . .	13
3.3	Vital rates . . . . .	15
3.4	Detection probability . . . . .	17
<b>4</b>	<b>Summary of improvements made</b>	<b>18</b>
<b>5</b>	<b>Suggestions for future improvements</b>	<b>18</b>
<b>6</b>	<b>Acknowledgements</b>	<b>18</b>
<b>7</b>	<b>Data availability</b>	<b>18</b>
	<b>References</b>	<b>20</b>
	<b>Appendices</b>	<b>21</b>

# 1 Introduction

Non-invasive genetic sampling (NGS) and dead recoveries, are a centerpiece of national and regional large carnivore monitoring in Norway and Sweden. Both countries have accumulated an extensive individual-based data set for wolf (*Canis lupus*) and plan to continue such transnational monitoring in the future.

Since 2017, project RovQuant has been developing statistical methods that allow a comprehensive assessment of the status and dynamics of large carnivore populations using NGS data and other sources of information collected and stored in Rovbase ([www.rovbase.se](http://www.rovbase.se), [www.rovbase.no](http://www.rovbase.no)) by the national monitoring programs in Sweden and Norway. At the core of the analytical framework developed by RovQuant (Bischof et al., 2019b, 2020) are Bayesian open-population spatial capture-recapture (OPSCR) models (Ergon and Gardner, 2014; Bischof et al., 2016; Chandler et al., 2018). These models use the spatial and temporal information contained in the repeated genetic detections of individuals to estimate various population parameters, including spatially-explicit abundance and vital rates. Importantly, the approach accounts for imperfect detection (i.e., the fact that some individuals are not detected at all) and animal movement (i.e., the fact that individuals may use and be detected in multiple management units or countries).

The conventional approach for wolf population size estimation in Scandinavia since 2013/2014 entails the use of a static conversion factor to calculate the size of the wolf population from the number of observed reproductions (Svensson et al., 2023). The conversion factor approach assumes that the relationship between total population size and the number of observed reproduction remains unchanged over time and between regions, which has led to criticism of the method (Chapron et al., 2016; Bischof et al., 2019a). Aside from relaxing this and several other assumptions (Bischof et al., 2019a), the OPSCR method brings along several advantages, including the ability to map density, derive jurisdiction-specific abundance, estimate survival and recruitment (which are needed for making population projections), and yield tractable measures of uncertainty (Bischof et al., 2019a, 2020).

RovQuant reported its first results for wolves in 2019 (Bischof et al., 2019a,b, 2020) and on an annual basis since 2021 (Milleret et al., 2021, 2022c). In the present report we summarize the analysis of a 10-year time series (2013/2014–2022/2023) using the latest available wolf monitoring data and the most recent version of the OPSCR model. We provide the following information:

- Annual estimates of the number of wolves (with credible intervals) for Sweden, Norway, and both countries combined, as well as estimates by county in Sweden and by large carnivore management regions in both countries.
- Annual maps of wolf density throughout the species’ primary range in Scandinavia.
- Annual estimates of survival, cause-specific mortality, recruitment, and population growth rate (with credible intervals).
- Estimated proportion of individuals detected through non-invasive genetic sampling (with credible intervals).

### Box 1: Terms and acronyms used

**AC:** Activity center. Model-based equivalent to the center of an individual's home range during the monitoring period. "AC location" refers to the spatial coordinates of an individual AC in a given year, and "AC movement" to the movement of an individual AC between consecutive years.

**CrI:** 95% credible interval associated with a posterior sample distribution.

**Detectors:** Potential detection locations in the spatial capture-recapture framework. These can refer to fixed locations (e.g., camera-trap locations) or to areas searched (e.g., habitat grid cells where searches for genetic samples were conducted; used in this project).

**Statsforvalteren:** Norwegian state's representative in the county, responsible for following up decisions, goals, and guidelines from the legislature and the government.

**Habitat buffer:** Buffer surrounding the searched area that is considered suitable habitat but was not searched.

**Legal culling:** Lethal removal of individuals by legal means including licensed recreational hunting, management removals, defense of life and property.

**Länsstyrelserna:** Swedish County Administrative Boards, in charge of the monitoring of large carnivores at the county level.

**MCMC:** Markov Chain Monte Carlo.

**NGS:** Non-invasive genetic sampling.

**OPSCR:** Open-population spatial capture-recapture.

**p0:** Baseline detection probability; probability of detecting an individual at a given detector, if the individual's AC is located exactly at the detector location.

**$\sigma$ :** Scale parameter of the detection function; related to the size of the circular home-range.

**SCR:** Spatial capture-recapture.

**SNO:** Statens naturoppsyn (Norwegian Nature Inspectorate) is the operative field branch of the Norwegian Environment Directorate (Miljødirektoratet).

## 2 Methods

### 2.1 Data

We included data from multiple sources, the primary one being the Scandinavian large carnivore database Rovbase 3.0 (rovbase.se and rovbase.no; last extraction: 2023-05-18). This database is used jointly by Norway and Sweden to record detailed information associated with large carnivore monitoring, including, but not limited to, non-invasive genetic sampling (NGS) data, dead recoveries, and GPS search tracks. In the following sections, we describe the various types of data used in the analysis. We used data collected during winters 2013/2014 to 2022/2023.

**Non-invasive genetic sampling** Swedish and Norwegian management authorities, SNO (Norway) and Länsstyrelserna (Sweden) conduct annual searches for sources of DNA (primarily scats and urine) throughout the Scandinavian wolf range (Svensson et al., 2023). Although samples may be collected throughout the year, the official survey period starting with the 2014/2015 season has been Oct 1 – Mar 31. About one third (30%) of DNA samples originated from opportunistic searches (without associated information on search effort) conducted by hunters or other members of the public (see Liberg et al. 2012 and Bischof et al. 2019a for further details about the data collection procedure). For individual identification, at least 17 microsatellite markers were amplified with PCR throughout the study period. Samples were amplified four times to account for the occurrence of allelic dropout and false alleles (Taberlet et al., 1996). Consensus genotypes were constructed from the replicated PCR runs using the threshold rule that the same alleles had to appear at least twice for a heterozygous genotype and three times for a homozygous genotype (Åkesson et al., 2016). Since 2017/2018, 90 single nucleotide polymorphisms (SNPs) were used for individual identification, in addition to microsatellites. Samples were amplified and genotypes visualized twice using nanofluid arrays (Fluidigm Inc.). Consensus SNP genotypes consisted of the markers with the same allele composition from the two amplifications (Åkesson et al., 2018).

**Dead recoveries** In Scandinavia, all dead large carnivores killed legally (e.g., legal hunting, management kills, defense of life and property) have to be reported to the state authorities (Fylkesmannen or SNO in Norway and Länsstyrelserna or the police in Sweden). All wolves found dead due to other reasons (e.g., natural deaths, vehicle and train collisions, illegal hunting) also have to be reported, but an unknown proportion remains undetected. Tissue is collected from all reported dead carnivores for DNA extraction and analysis. DNA from dead carnivores, if genotyped, can be linked with NGS data via individual IDs and provide definite information about the fate of individuals. Dead recoveries associated with GPS-coordinates also provide additional information for the estimation of individual locations, especially valuable for individuals with few or no NGS detections (Dupont et al., 2021).

**GPS search tracks** Government employees involved in systematic searches for wolf DNA along roads and following wolf tracks (e.g., via snowmobiles, skis, snowshoes) documented their effort with GPS track logs which were registered in Rovbase 3.0. GPS search tracks were included in the OPSCR model to account for spatial and temporal variation in search effort.

**Observation reports in Skandobs** We used all observation records available from Skandobs that were recorded during the wolf monitoring seasons since 2012, skandobs.se; last extraction: 2023-05-08). Skandobs is a web application that allows anyone to anonymously register observations (e.g., visual, tracks, faeces) of bears, lynx, wolves, and wolverines in Scandinavia. This data consists of more than 50 000 records of possible large carnivore observations. Although most observations are not verified, they offer the best available proxy for spatio-temporal variation in opportunistic effort.



**Parentage analysis and social status assignation** Each year, field observations, genetic sampling data, dead recoveries, and parentage analysis (which resulted in a near-complete pedigree of the Scandinavian wolf population) are used to establish counts of pairs, packs, and reproductions (Svensson et al., 2014, 2015, 2017, 2019, 2021, 2023; Wabakken et al., 2016, 2018, 2020, 2022). The social status (adult scent-marking member in a pack/pair, or subadult member of a pack and vagrant individuals) was determined on a yearly basis for genetically identified individuals using the pedigree and tracking information from the monitoring (Åkesson et al., 2022). The identity of known adult scent-marking individuals was reported in the yearly monitoring reports (Svensson et al., 2014, 2015, 2017, 2019, 2021, 2023; Wabakken et al., 2016, 2018, 2020, 2022) and a detailed description of the method used for parental assignment was provided in Åkesson et al. (2016). We assigned individuals to two different states depending on whether the individual had been identified as an adult scent-marking member of a pair or a pack (“adult scent-marking”) or not (“other”). See Bischof et al. (2019a) for further details.

**Study area** Although wolves can be occasionally present within the entire Scandinavian range, we restricted the analysis to the core area (Figure 1). Following the recent wolf range expansion towards the south of Sweden, the Swedish Environmental Protection Agency requested an expansion of the study area (Milleret et al., 2022c). The RovQuant models therefore included an additional 68 400 km<sup>2</sup> in the study area (total= 322 400 km<sup>2</sup>), compared to the previous analyses of the Scandinavian monitoring wolf data by RovQuant (Bischof et al., 2019b, 2020; Milleret et al., 2021).

## 2.2 Open-population spatial capture-recapture model

We analysed the data using a Bayesian open-population spatial capture-recapture (OPSCR) model (Bischof et al., 2019b), which addresses three challenges associated with population-level wildlife inventories:

1. Detection is imperfect and sampling effort heterogeneous in space and time: not all individuals present in the study area are detected (Kéry and Schaub, 2012).
2. Individuals that reside primarily outside the surveyed area may be detected within it. Without an explicit link between the population size parameter and geographic space or area, density cannot be estimated and population size is ill-defined (Efford, 2004).
3. Non-spatial population dynamic models usually estimate “apparent” survival and recruitment, as these parameters include the probability of permanent emigration and immigration, respectively. By explicitly modelling movement of individuals between years, the OPSCR model can help return unbiased estimates of demographic parameters (Ergon and Gardner, 2014; Schaub and Royle, 2014; Gardner et al., 2018; Efford and Schofield, 2022).

The OPSCR model (Ergon and Gardner, 2014; Bischof et al., 2016; Chandler et al., 2018) is composed of three sub-models:

1. A model for population dynamics and population size.
2. A model for density and movements.
3. A model for detections during DNA searches.

**Population dynamics and population size sub-model** We used a multi-state formulation (Lebreton and Pradel, 2002) where each individual life history is represented by a succession of up to five discrete states  $z_{i,t}$ : 1) “unborn” if the individual has not been recruited in the population (state “unborn” is required for the data augmentation procedure, see below); 2) “alive other” if it is alive and not assigned as an “adult scent-marking” individual; 3) “alive adult scent-marking” if it is alive and assigned as an “adult scent-marking” individual; 4) “culled” if it was

culled and therefore recovered dead between the start of the previous and current monitoring season; or 5) “dead”: if it has died but was not recovered dead. We then modelled the transition between states from one monitoring season to the next ( $t$  to  $t + 1$ ) to estimate vital rates (recruitment and mortality). More details are available in Bischof et al. (2019b), and Bischof et al. (2020). We used data augmentation (Royle and Dorazio, 2012), whereby additional, undetected individuals are available for inclusion in the population at each time step.

**Density and movement sub-model** We used a Bernoulli point process to model the distribution of individual activity centers. In the first year, individuals were located according to an intensity surface, which was a function of the locations of known packs at time  $t - 1$  (see Bischof et al., 2019b and Bischof et al., 2020 for more details). For all subsequent years ( $t > 1$ ), the location of individual activity centers was a function of the distance from previous activity centers (at time  $t - 1$ ), as well as a function of the locations of known packs (at time  $t - 1$ ). As in the previous analysis (Milleret et al., 2022c), we used an exponential model to describe the movement of individuals between years, as it better accommodates distributions with long tails (i.e., a few individuals that make exceptionally long dispersal movements).

**Detection sub-model** SCR models take into account the spatial-variation in individual detection probability based on the distance between activity center locations (estimated by the density sub-model) and a given detector. A half-normal function was used to express the declining probability of detection with increasing distance between the activity center and the detector.

In Scandinavia, DNA material from live wolves are collected following two main processes. First, authorities collect genetic samples and record the corresponding search effort during official searches ("structured sampling" thereafter). Second, DNA material can be collected by any member of the public (e.g., hunters) or by the authorities in a more or less opportunistic manner and search effort is not directly available ("unstructured" sampling thereafter). Currently, it is not possible to unambiguously distinguish between samples collected by the authorities during the structured or unstructured sampling in Rovbase. We therefore assigned each sample to structured or unstructured sampling based on whether a given sample matched in time and space with recorded search tracks: a sample was assigned to the "structured" sampling if it was collected by the authorities (marked as collected by "Statsforvalteren" or "SNO" in Rovbase) and if it was located within 500 m from a search track recorded on the same day. All remaining samples were assigned to the unstructured sampling.

We assumed that both sampling processes could in theory occur within the entire study area and therefore used the same 10 km detector grid for both observation processes. Samples were then assigned to the closest detector (see details in Bischof et al., 2019b, and Bischof et al., 2020). However, spatial and temporal variation in the probability to detect a sample during the structured or unstructured sampling were assumed to be driven by different processes.

We accounted for spatio-temporal heterogeneity in detectability during *structured sampling* using:

- Spatial-temporal variation in effort using the length of GPS search tracks in each detector grid cell.
- Spatio-temporal variation in snow cover.
- Spatio-temporal variation in monitoring regimes between jurisdictions (groups of counties in Sweden, monitoring regions in Norway).
- Individual variation linked with the state of the individual (i.e., adult scent-marking individuals, other individuals).

- Individual variation linked with a previous detection that could be expected to positively or negatively influence the probability of being detected at subsequent occasions, depending on the state of knowledge in the territory or neighboring territories.

We accounted for spatio-temporal heterogeneity in detectability during *unstructured sampling* using:

- Spatio-temporal variation in unstructured sampling or carnivore observations (Figure A.1). For each detector grid cell and during each monitoring season (Oct 1 – Mar 31), we identified whether a) any carnivore sample had been registered in Rovbase (excluding successfully genotyped wolf samples already used in the OPSCR analysis) or b) any observation of carnivores had been registered in Skandobs. Roughly, the variable distinguishes areas with very low detection probability from those with some reasonable probability that carnivore DNA samples, if present in a detector grid cell, could have been detected and submitted for genetic analysis (Figure A.1).
- Spatio-temporal variation in snow cover.
- Spatial variation in accessibility measured as the average distance to the nearest road.
- Spatio-temporal variation between countries.
- Individual and temporal variation linked with the state of the individual (i.e., adult scent-marking individuals, other individuals).
- Individual and temporal variation linked with a previous detection that could be expected to positively influence the probability of being detected at subsequent occasions.

The different model components and data sources for covariates are described in detail in Bischof et al. (2019a), Bischof et al. (2019b), and Bischof et al. (2020).

**Model fitting** We fitted sex-specific Bayesian OPSCR models using Markov chain Monte Carlo (MCMC) simulation with NIMBLE version 0.12.2 (Turek et al., 2021; de Valpine et al., 2017; de Valpine et al., 2022) and nimbleSCR (Bischof et al., 2021) in R version 4.1.0 (R Core Team, 2021). We ran 4 chains each with 36 000 iterations, including a 7 000-iterations burn-in period. Due to the computing challenge associated with post-processing large amounts of data, we thinned chains by a factor of 10 from which abundance estimates were derived. We considered models as converged when the Gelman-Rubin diagnostics (Rhat, Gelman and Rubin, 1992) was  $\leq 1.1$  for all parameters and by visually inspecting the trace plots.

**Abundance estimates** To obtain an estimate of abundance for any given area, we summed the number of predicted AC locations (live individuals) that fell within that area for each iteration of the MCMC chains, thus generating a posterior distribution of abundance for that area. In this fashion, abundance estimates and the associated uncertainty can be extracted for any spatial unit, including countries or counties (Figure A.2). Individuals detected near a border can have their model-predicted AC placed on different sides of that border in different model iterations (even if detections are only made on one side of the border). As a result, the probability of designating such individuals to either side of the border can be integrated into jurisdiction-specific abundance estimates. This is especially relevant for wolves detected along the Swedish and Norwegian border ("cross-boundary wolves", Wabakken et al., 2022); individual wolves can be partially designated to both countries (Bischof, 2015).

To ensure that abundance estimates for spatial sub-units (jurisdictions) add up to the overall abundance estimate, we used the mean and associated 95% credible interval limits to summarize posterior distributions of abundance. Combined (female/male) parameter estimates were

obtained by merging posterior samples from the sex-specific models.

**Density maps** We used both the distribution of model-estimated AC positions and the scale parameter ( $\sigma$ ) of the detection function to construct density maps based on individual utilization distributions. These maps are not only based on the position of the center of an individual's home range, but also take into account the area over which that individual's activity is spread, i.e., its space use (Bischof et al., 2020). To do so, we constructed raster maps (5 km resolution) of individual utilization distributions, scaled values in each raster to sum to one, and then summed rasters across individuals to create a single population-level raster map for each iteration. An overall density map was derived by calculating the mean across iterations in each cell (Bischof et al., 2020).

**Other derived parameters** The average proportion of individuals detected and the associated uncertainty were obtained by dividing the number of individuals detected through NGS sampling (Table A.3) by the abundance estimates and their associated credible intervals, respectively. We derived the proportion of females in the population and the associated uncertainty by dividing the posterior distribution representing the number of females by the combined abundance estimate for males and females (Table A.3). Yearly population growth rates ( $\lambda$ ; Table A.6) were calculated as  $\lambda_t = N_{t+1}/N_t$ .

**Focus on uncertainty** Although we reported median (or mean for abundance; see above) estimates for all parameters in the tables, we intentionally focused the main results of this report on the 95% credible interval limits of the estimates. We did so with the aim of drawing the reader's attention to the uncertainty around population size estimates, rather than a single point estimate (Milleret et al., 2022b).

## 3 Results

### 3.1 Non-invasive genetic samples and dead recoveries


A total of 18 693 (8 197 female; 10 496 male) genotyped wolf genetic samples were included in the analysis, of which 75% originated from Sweden. These samples were associated with 1 767 (958 female; 809 male) individuals. We did not include individuals with unknown sex in this analysis. Among all genotyped samples, 8 053 (3358 female; 4695 male) were assigned to the structured sampling and 10 640 (4839 female; 5801 male) to the unstructured sampling. We also used 684 (306 female; 378 male) dead recoveries of wolves in the OPSCR model, of which 559 (248 female; 311 male) were due to legal culling and 125 (58 female; 67 male) due to other causes of mortality. The majority of dead recoveries (68%) originated from Sweden. Annual total and country-specific tallies of NGS samples and associated individuals, as well as dead recoveries included in the analysis are provided in the Appendices (NGS samples: Table A.2, NGS individuals: Table A.3, dead recoveries: Table A.4)

### 3.2 Density and abundance

Wolf abundance for the entire study area (322 400 km<sup>2</sup>, excluding the buffer area) was likely (95% credible interval) between 441 and 488 individuals in 2022/2023 (Table 1, Figure 1). Estimates refer to the status of the population at the start of the annual sampling period (Oct. 1). The proportion of females in the Scandinavian wolf population was likely between 47% and 52% in 2022/2023. Based on the model-predicted location of ACs, we estimated that in 2022/2023, between 376 to 415 individuals could be attributed to Sweden and 59 to 80 to Norway (Table 1). See Table 1 for total and sex-specific abundance estimates for each country and management area. See Table A.5 for annual wolf abundance estimates for all of Scandinavia and by region between 2012/2013 and 2022/2023. Note that estimates for different years (Figure A.3) shown here differ slightly from those provided in Bischof et al. (2020), Milleret et al. (2021), and Milleret et al. (2022c). This is due to the use of an updated OPSCR model and the inclusion of additional years of data. The analysis yielded annual density maps, which illustrate changes in the distribution of wolves over time (Figure A.4).

**Table 1:** Wolf population size estimates in 2022/2023 by sex at several spatial scales: the entire study area, by country, by management unit (carnivore management regions in Norway and "Rovdjursförvaltningsområden" in Sweden), and counties ("Län" in Sweden); see also Figure A.2. Only counties and management units that are within or that intersect the study area are included in the table. Readers should focus on the 95% credible interval provided in parentheses, as these - unlike mean values - convey uncertainty inherent in abundance estimates. Numbers are based on estimated activity center locations of wolves. Combined female-male estimates were obtained by joining sex-specific posterior distributions. Rounding may result in small deviations between total estimates and the sum of the estimates for constituent regions. Note that the numbers reported here are predictions from a statistical model which always represents an oversimplification of reality and is based on available data (NGS and dead recoveries). As a consequence, especially at the local scale, the model-estimated number of wolves based on DNA sampling can deviate from the number of wolves inferred from ancillary observations (e.g., camera traps).


	Females	Males	Total
TOTAL	230.4 (217-246)	232.3 (216-253)	462.7 (441-488)
NORWAY	38.9 (32-48)	29.7 (24-38)	68.6 (59-80)
Region 2	0.3 (0-2)	0.3 (0-2)	0.5 (0-2)
Region 3	1 (0-3)	1 (0-3)	2 (0-5)
Region 4	13.8 (11-17)	10.9 (8-14)	24.7 (21-29)
Region 5	22.3 (17-29)	16.7 (12-23)	39 (32-48)
Region 6	1.4 (0-4)	0.9 (0-3)	2.3 (0-6)
SWEDEN	191.5 (181-204)	202.7 (189-220)	394.1 (376-415)
Norra	11.3 (8-16)	11.9 (8-17)	23.2 (18-30)
Jämtland	7.1 (4-11)	8.4 (5-13)	15.5 (11-21)
Västerbotten	0 (0-0)	0 (0-0)	0 (0-1)
Västernorrland	4.2 (2-7)	3.4 (2-6)	7.6 (5-11)
Mellersta	152.8 (144-163)	157.8 (146-172)	310.5 (295-328)
Dalarna	34.1 (29-40)	22.6 (18-29)	56.7 (50-65)
Gävleborg	19.6 (16-23)	14.8 (12-19)	34.4 (30-40)
Örebro	20.9 (17-25)	29.8 (25-35)	50.7 (44-57)
Stockholm	2.3 (1-4)	1.6 (0-4)	3.8 (1-7)
Uppsala	4.2 (3-6)	3 (1-6)	7.2 (5-11)
Värmland	42.6 (37-49)	56 (51-62)	98.6 (91-108)
Västmanland	14.3 (11-18)	15.8 (12-20)	30.1 (25-36)
VästraGötaland	14.8 (12-18)	14.2 (11-18)	29 (25-34)
Södra	27.4 (24-32)	33 (29-38)	60.4 (55-67)
Blekinge	0.1 (0-1)	0.1 (0-1)	0.2 (0-1)
Halland	1.1 (0-2)	0.3 (0-2)	1.3 (0-3)
Jönköping	4.4 (3-6)	7.3 (5-10)	11.7 (9-15)
Kalmar	0.4 (0-2)	0.5 (0-2)	0.8 (0-3)
Kronoberg	1.2 (0-3)	2.2 (1-4)	3.5 (2-6)
Östergötland	3.4 (2-5)	5.8 (5-8)	9.1 (8-12)
Skåne	3.4 (3-5)	5.4 (5-7)	8.8 (8-11)
Södermanland	13.5 (12-16)	11.5 (10-14)	24.9 (22-28)




**Figure 1:** Wolf density based on individual utilization distributions throughout the study area (white background) in Scandinavia in 2022/2023. This map is freely available as a geo-referenced raster file at <https://github.com/richbi/RovQuantPublic>

### 3.3 Vital rates

The OPSCR model produced annual estimates of legal hunting mortality, mortality associated with all other causes, and per capita recruitment rates (Figure 2; Table A.7). There was temporal variation in mortality rates, with the risk of mortality from other causes than culling generally higher than the risk of mortality from culling. Overall, scent-marking individuals had a higher survival compared to other individuals, with no pronounced systematic difference between sexes (Figure A.5).



**Figure 2:** Mortality probabilities due to legal culling (light green) and all other causes (dark green) for female and male wolves. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Shown are overall estimates throughout the study area. Estimates refer to deaths occurring between the start of one sampling season and the start of the next.



**Figure 3:** Estimated annual number of recruits for female and male wolves. Recruitment represents the number of new individuals present in the population on Oct 1 (i.e., individuals that were born between the two consecutive monitoring seasons and survived to Oct 1 or that immigrated in the study area). Darker and lighter bars show the 50% and 95% credible intervals, respectively.



### 3.4 Detection probability

The overall proportion of detected individuals in the study area was likely between 81% and 90% in 2022/23 (Table A.11). The baseline detection probability for the structured and unstructured sampling varied both in time and space (Figure A.6 and Figure A.7). More specifically, the length of recorded search tracks positively affected detection probability of the structured sampling (2022/2023; males:  $\beta = 0.35$ , CrI: 0.27 - 0.44; females:  $\beta = 0.34$ , CrI: 0.28 - 0.41; Table A.9). However, the previous detection of individuals and the average proportion of snow cover had no strong effect on detection probability for the structured sampling (Table A.9). Unstructured search effort derived using the observation data in Skandobs and Rovbase had a strong positive effect on detection probability during unstructured sampling (2022/2023; males:  $\beta = 1.09$ , CrI: 0.56 - 1.68; females:  $\beta = 1.61$ , CrI: 1.04 - 2.33; Table A.10). The effect of previous detection also tended to have a positive effect on detection probability during unstructured sampling, but the pattern was not consistent across years (Table A.10).

## 4 Summary of improvements made

The analysis described in this report includes the following adjustments compared with previous analyses of wolf density in Scandinavia by RovQuant (Milleret et al., 2022c):

1. Addition of data from the 2022/2023 monitoring season.
2. Correct a previous error in the assignment of some samples to structured and unstructured sampling.

## 5 Suggestions for future improvements

RovQuant continues to work on improving the functionality and efficiency of OPSCR models. We plan to test and potentially implement the following developments in future analyses of the Scandinavian wolf monitoring data:

1. Review and adjust spatial covariates on density. This may involve the addition of land cover and topographic variables.
2. Account for individual heterogeneity in detectability/space use, for example by using a finite-mixture approach (Cubaynes et al., 2010).
3. Consider alternative detection models that do not assume a half-normal shape and/or circular home ranges (Sutherland et al., 2015).
4. Account for spatial variation in vital rates (i.e., survival, recruitment; Milleret et al. (2022a)).

## 6 Acknowledgements

This work was made possible by the large carnivore monitoring programs and the extensive monitoring and observation data collected by Swedish (Länstyrelsen) and Norwegian (SNO) wildlife management authorities, as well as the public in both countries. Our analysis relied on genetic analyses conducted by the laboratory personnel at the DNA laboratories at the Swedish University of Agricultural Sciences, and the Norwegian Institute for Nature Research. We also thank Swedish and Norwegian wildlife managers for feedback provided during project RovQuant and the Research Council of Norway for partial funding (NFR 286886; project WildMap). The computations/simulations were performed on resources provided by NMBU's computing cluster "Orion", administered by the Centre for Integrative Genetics and by UNINETT Sigma2 - the National Infrastructure for High Performance Computing and Data Storage in Norway. We are grateful to P. de Valpine and D. Turek for help with the formulation of the OPSCR model in Nimble. J. Vermaat provided helpful comments on a draft of this report.

## 7 Data availability


Data, R code to reproduce the analysis, as well as figures, tables, and raster maps (Figure A.4) are available at <https://github.com/richbi/RovQuantPublic>.

## References


- Åkesson, M., Danielsson, A., Hedmark, E., and Öhrn (2018). Teknisk rapport över genetiska analyser på varg i sverige år 2017. *Rapport från SLU*, (162 s).
- Åkesson, M., Liberg, O., Sand, H., Wabakken, P., Bensch, S., and Flagstad, O. (2016). Genetic rescue in a severely inbred wolf population. *Molecular Ecology*, 25(19):4745–4756.
- Bischof, R. (2015). Approaches for assessing illegal hunting of brown bears and other large carnivores in sweden. *Report 2015:5 from the Scandinavian Brown Bear Research Project to the Swedish Environmental Protection Agency. Norwegian University of Life Sciences, Ås, Norway.*
- Bischof, R., Brøseth, H., and Gimenez, O. (2016). Wildlife in a Politically Divided World: Insularism Inflates Estimates of Brown Bear Abundance. *Conservation Letters*, 9(2):122–130.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Åkesson, M., BrOseth, H., and Kindberg, J. (2019a). Estimating the size of the scandinavian wolf population with spatial capture recapture and conversion factors. *MINAflagrapport57*, 80pp.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., BrOseth, H., and Kindberg, J. (2019b). Estimating density, abundance and population dynamics of bears, wolverines, and wolves in scandinavia. *MINAflagrapport63*, 79pp.
- Bischof, R., Milleret, C., Dupont, P., Chipperfield, J., Tourani, M., Ordiz, A., de Valpine, P., Turek, D., Royle, J. A., Gimenez, O., Øystein Flagstad, Åkesson, M., Svensson, L., Brøseth, H., and Kindberg, J. (2020). Estimating and forecasting spatial population dynamics of apex predators using transnational genetic monitoring. *Proceedings of the National Academy of Sciences*, 117(48):30531–30538.
- Bischof, R., Turek, D., Milleret, C., Ergon, T., Dupont, P., Dey, S., and de Valpine, P. (2021). *nimbleSCR: Spatial Capture-Recapture (SCR) Methods Using 'nimble'*. R package version 0.1.2.
- Chandler, R. B., Hepinstall-Cymerman, J., Merker, S., Abernathy-Conners, H., and Cooper, R. J. (2018). Characterizing spatio-temporal variation in survival and recruitment with integrated population models. *The Auk*, 135(3):409–426.
- Chapron, G., Wikenros, C., Liberg, O., Wabakken, P., Flagstad, O., Milleret, C., Månsson, J., Svensson, L., Zimmermann, B., Åkesson, M., and Sand, H. (2016). Estimating wolf (*canis lupus*) population size from number of packs and an individual based model. *Ecological Modelling*, 339:33 – 44.
- Cubaynes, S., Pradel, R., Choquet, R., Duchamp, C., Gaillard, J. M., Lebreton, J. D., Marboutin, E., Miquel, C., Reboulet, A. M., Poillot, C., Taberlet, P., and Gimenez, O. (2010). Importance of accounting for detection heterogeneity when estimating abundance: the case of French wolves. *Conservation Biology*, 24(2):621–626.
- de Valpine, P., Paciorek, C., Turek, D., Michaud, N., Anderson-Bergman, C., Obermeyer, F., Wehrhahn Cortes, C., Rodriguez, A., Temple Lang, D., and Paganin, S. (2022). *NIMBLE User Manual*. R package manual version 0.12.2.
- de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., and Bodik, R. (2017). Programming with models: writing statistical algorithms for general model structures with nimble. *Journal of Computational and Graphical Statistics*, 26(2):403–413.
- Dupont, P., Milleret, C., Tourani, M., BrOseth, H., and Bischof, R. (2021). Integrating dead recoveries in open-population spatial capture–recapture models. *Ecosphere*, 12(7):e03571.
- Efford, M. (2004). Density estimation in live-trapping studies. *Oikos*, 106(3):598–610.
- Efford, M. G. and Schofield, M. R. (2022). A review of movement models in open population capture–recapture. *Methods in Ecology and Evolution*, 13(10):2106–2118.
- Ergon, T. and Gardner, B. (2014). Separating mortality and emigration: modelling space use, dispersal and survival with robust-design spatial capture–recapture data. *Methods in Ecology and Evolution*, 5(12):1327–1336.
- Gardner, B., Sollmann, R., Kumar, N. S., Jathanna, D., and Karanth, K. U. (2018). State space and movement specification in open population spatial capture–recapture models. *Ecology and Evolution*, 8(20):10336–10344.
- Gelman, A. and Rubin, D. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7:457–511. <http://www.stat.columbia.edu/~gelman/research/published/itsim.pdf>.
- Kéry, M. and Schaub, M. (2012). *Bayesian population analysis using WinBUGS: a hierarchical perspective*. Academic Press, Waltham, MA.
- Lebreton, J.-D. and Pradel, R. (2002). Multistate recapture models: modelling incomplete individual histories. *Journal of Applied Statistics*, 29:353–369.
- Liberg, O., Aronson, Å., Sand, H., Wabakken, P., Maartmann, E., Svensson, L., and Åkesson, M. (2012). Monitoring of wolves in scandinavia. *Hystrix, the Italian Journal of Mammalogy*, 23(1):29–34.
- Milleret, C., Dey, S., Dupont, P., Turek, D., de Valpine, P., and Bischof, R. (2022a). Estimating spatially variable and density-dependent survival using open-population spatial capture-recapture models. *bioRxiv*.
- Milleret, C., Dupont, P., BrOseth, H., Flagstad, O., Kindberg, J., and Bischof, R. (2022b). Estimates of wolverine density, abundance, and population dynamics in scandinavia, 2013–2021. *MINAflagrapport 74*, 30pp.
- Milleret, C., Dupont, P., Åkesson, M., BrOseth, H., Kindberg, J., and Bischof, R. (2021). Estimates of wolf

- density, abundance, and population dynamics in scandinavia, 2012 - 2021. *MINAfaagrapport 72*, 30pp.
- Milleret, C., Dupont, P., Åkesson, M., BrOseth, H., Kindberg, J., and Bischof, R. (2022c). Estimates of wolf density, abundance, and population dynamics in scandinavia, 2013-2022. *MINAfaagrapport 77*, 35pp.
- R Core Team (2021). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Royle, J. A. and Dorazio, R. M. (2012). Parameter-expanded data augmentation for bayesian analysis of capture–recapture models. *Journal of Ornithology*, 152(2):521–537.
- Schaub, M. and Royle, J. A. (2014). Estimating true instead of apparent survival using spatial Cormack–Jolly–Seber models. *Methods in Ecology and Evolution*, 5(12):1316–1326.
- Sutherland, C., Fuller, A. K., and Royle, J. A. (2015). Modelling non-Euclidean movement and landscape connectivity in highly structured ecological networks. *Methods in Ecology and Evolution*, 6(2):169–177.
- Svensson, L., Wabakken, P., Kojola, I., Maartmann, E., Strømseth, T., Åkesson, M., and Flagstad, Ø. (2014). Varg i Skandinavien och Finland: Slutrapport från inventering av varg vintern 2013–2014. Technical report.
- Svensson, L., Wabakken, P., Maartmann, E., Åkesson, M., and Flagstad, Ø. (2015). Inventering av varg vintern 2014-2015. Inventeringsresultat för stora rovdjur i Skandinavien 1-2015. Technical report.
- Svensson, L., Wabakken, P., Maartmann, E., Åkesson, M., and Flagstad, Ø. (2017). Inventering av varg vintern 2016-2017 / Bestandsövervakning av ulv vintern 2016-2017. Technical report.
- Svensson, L., Wabakken, P., Maartmann, E., Cardoso Palacios, C., Flagstad, O., and Åkesson, M. (2021). Inventering av varg vintern 2020-2021. bestandsövervakning av ulv vintern 2020-2021. *Bestandsstatus för stora rovdjur i Skandinavien*, 1(55s).
- Svensson, L., Wabakken, P., Maartmann, E., Nordli, K., Flagstad, , Danielsson, A., Hensel, H., Pöchhacker, K., and Åkesson, M. (2023). Inventering av varg vintern 2022- 2023. bestandsövervakning av ulv vintern 2022-2023. *Bestandsstatus för stora rovdjur i Skandinavien*, 1(65s).
- Svensson, L., Wabakken, P., Maartmann, E., Åkesson, M., Flagstad, O., and Hedmark, E. (2019). Inventering av varg vintern 2018-2019. bestandsövervakning av ulv vintern 2018-2019. *Bestandsstatus för stora rovdjur i Skandinavien*, 1(53s).
- Taberlet, P., Griffin, S., Goossens, B., Questiau, S., Manceau, V., Escaravage, N., Waits, L. P., and Bouvet, J. (1996). Reliable genotyping of samples with very low dna quantities using pcr. *Nucleic acids research*, 24(16):3189–3194.
- Turek, D., Milleret, C., Ergon, T., Brøseth, H., Dupont, P., Bischof, R., and de Valpine, P. (2021). Efficient estimation of large-scale spatial capture–recapture models. *Ecosphere*, 12(2):e03385.
- Wabakken, P., Svensson, L., , Maartmann, E., Nordli, K., Flagstad, O., and Åkesson, M. (2022). Bestandsövervakning av ulv vintern 2021-2022. inventering av varg vintern 2021-2022. *Bestandsstatus för stora rovdjur i Skandinavien*, 1(59s).
- Wabakken, P., Svensson, L., Maartmann, E., Åkesson, M., and Flagstad, Ø. (2016). Bestandsövervakning av ulv vintern 2015-2016. Technical report.
- Wabakken, P., Svensson, L., Maartmann, E., Nordli, K., Flagstad, O., and Åkesson, M. (2020). Bestandsövervakning av ulv vintern 2019-2020. inventering av varg vintern 2019-2020. *Bestandsstatus för stora rovdjur i Skandinavien*, 1(55s).
- Wabakken, P., Svensson, L., Maartmann, E., Åkesson, M., and Flagstad, (2018). Bestandsövervakning av ulv vintern 2017-2018 / Inventering av varg vintern 2017-2018. Technical report.
- Åkesson, M., Svensson, L., Flagstad, , Wabakken, P., and Frank, J. (2022). Wolf monitoring in scandinavia: evaluating counts of packs and reproduction events. *The Journal of Wildlife Management*, 86(4):e22206.


# Appendices



**Figure A.1:** Covariate used to account for spatio-temporal variation in unstructured sampling in the study area. Green cells (10 x 10 km) represent areas with at least one carnivore record in Rovbase (rovbase.no, rovbase.se, excluding the ones used in the OPSCR model) or an observation record from Skandobs (skandobs.se, skandobs.no) during a given monitoring season (Oct 1 – Mar 31).




**Figure A.2:** Management jurisdictions in Norway and Sweden. Shaded polygons show carnivore management regions in Norway and counties in Sweden. Thick outlines delineate Swedish carnivore management regions ("Rovdjursförvaltningsområden") encompassing multiple counties.




**Figure A.3:** Total (black) and country-specific (blue: Sweden, red: Norway) annual wolf population size estimates in Scandinavia between 2013/2014 and 2022/2023. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Credible intervals indicate uncertainty in estimates given the model and data used to generate the estimates. Changes in the model and the data can result in different estimates and associated uncertainty compared with estimates provided in previous reports by RovQuant. Note the higher precision (lower uncertainty) in the three monitoring seasons (2016/17, 2017/18, 2019/20) with higher sampling intensity.







**Figure A.4:** Wolf density based on individual utilization distributions throughout the study area (white background) in Scandinavia between 2013/2014 and 2022/2023. These maps are freely available as geo-referenced raster files at <https://github.com/richbi/RovQuantPublic>.



**Figure A.5:** Annual survival probabilities for scent-marking adults and other individuals and for female and male wolves. Darker and lighter bars show the 50% and 95% credible intervals, respectively. Combined mortality probabilities (and therefore overall survival) are obtained by adding the mortality probabilities due to legal culling and all other causes (Figure 2). Shown are overall estimates for the entire study area between 2013/2014 and 2022/2023.



**Figure A.6:** Sex-specific baseline detection probabilities ( $p_{0structured}$ ) of scent-marking adults and other individuals for the different Scandinavian jurisdictions during structured sampling as estimated by the open-population spatial capture-recapture model. Bars represent 95% credible intervals. Results are separated into panels based on regions. Estimates are shown for the mean values of the detection covariates. Note that baseline detection probability ( $p_0$ ) is a theoretical value of detection probability when a detector coincides with the location of an individual's activity center. It is not to be confused with detectability, i.e., the overall probability of detecting an individual.



**Figure A.7:** Sex- and country-specific baseline detection probabilities ( $p_{0unstructured}$ ) of scent-marking adults and others individuals during unstructured sampling as estimated by the open-population spatial capture-recapture model. Bars represent 95% credible intervals. Estimates are shown for the mean values of the detection covariates. Note that baseline detection probability ( $p_0$ ) is a theoretical value of detection probability when a detector coincides with the location of an individual’s activity center. It is not to be confused with detectability, i.e., the overall probability of detecting an individual.

**Table A.1:** Annual number of wolf non-invasive genetic samples included in the analysis. Numbers are reported by country, for females (F) and males (M), and for each type of sampling (structured and unstructured). We included only samples collected within the study area during the primary monitoring period (Oct 1 - Mar 31) between 2013/2014 and 2022/2023.

		2013/2014		2014/2015		2015/2016		2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		2021/2022		2022/2023	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
<b>Norway</b>	Structured	0	1	4	5	6	11	22	11	36	42	52	93	47	81	51	133	71	162	59	110
	Unstructured	53	83	91	105	142	173	245	197	237	240	180	252	196	252	275	330	137	228	99	93
<b>Sweden</b>	Structured	80	110	98	120	137	200	373	582	474	666	222	293	597	716	281	457	419	499	329	403
	Unstructured	103	125	83	107	102	178	470	549	607	836	127	166	842	832	254	342	286	387	310	326
<b>Total</b>	Structured	80	111	102	125	143	211	395	593	510	708	274	386	644	797	332	590	490	661	388	513
	Unstructured	156	208	174	212	244	351	715	746	844	1076	307	418	1038	1084	529	672	423	615	409	419

**Table A.2:** Annual number of wolf non-invasive genetic samples included in the analysis. Numbers are reported by country, for females (F) and males (M), and for each type of sampling (structured and unstructured). We included only samples collected within the study area during the primary monitoring period (Oct 1 - March 31) between 2013/2014 and 2022/2023.

		2013/2014		2014/2015		2015/2016		2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		2021/2022		2022/2023	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
Norway		53	84	95	110	148	184	267	208	273	282	232	345	243	333	326	463	208	390	158	203
Sweden		183	235	181	227	239	378	843	1131	1081	1502	349	459	1439	1548	535	799	705	886	639	729
<b>Total</b>		236	319	276	337	387	562	1110	1339	1354	1784	581	804	1682	1881	861	1262	913	1276	797	932

**Table A.3:** Annual number of individual wolves detected via non-invasive genetic sampling and included in the analysis. Numbers are reported by country, for females (F) and males (M), and for each type of sampling (structured and unstructured). We included only individuals associated with samples collected within the study area during the primary monitoring period (Oct 1 - Mar 31) between 2013/2014 and 2022/2023. Some individuals were detected in both countries during the same year, hence the sum of the national counts can exceed the total number of unique individuals detected in Scandinavia.

		2013/2014		2014/2015		2015/2016		2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		2021/2022		2022/2023	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
<b>Norway</b>	Structured	0	1	3	4	5	6	11	7	18	17	16	21	17	23	22	25	30	34	21	23
	Unstructured	18	27	27	31	29	43	45	36	47	44	31	44	36	45	37	53	42	47	27	19
<b>Sweden</b>	Structured	49	64	59	71	68	82	111	130	115	126	82	91	136	153	107	115	130	147	120	129
	Unstructured	59	72	47	59	54	87	105	124	120	144	68	71	146	158	97	115	127	129	128	124
<b>Total</b>	Structured	49	65	61	75	73	88	121	136	130	140	93	108	150	171	124	135	153	173	137	147
	Unstructured	72	95	73	87	80	124	142	156	161	178	96	110	178	194	134	163	160	165	151	140

**Table A.4:** Number of cause-specific dead recoveries of wolves in Scandinavia between 2013 and 2023 that were included in the OPSCR analysis. Numbers are reported by country, for females (F) and males (M). Note that dead recovery data from the final season (2022/2023) were not used in this analysis because survival is modeled as part of the transition from one season to the next.

		Country		2013/2014		2014/2015		2015/2016		2016/2017		2017/2018		2018/2019		2019/2020		2020/2021		2021/2022		2022/2023	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
<b>Other</b>	Norway	1	1	0	0	1	3	0	0	1	3	1	1	1	1	0	2	1	2	0	0		
	Sweden	4	6	9	10	8	6	4	5	8	2	4	5	5	7	8	5	1	6	1	2		
<b>Legal culling</b>	Norway	4	4	3	5	5	1	6	6	18	16	10	8	6	12	11	16	13	23	8	5		
	Sweden	12	12	32	32	7	20	17	23	12	31	4	4	6	12	18	30	22	18	34	33		
<b>Total</b>	Total	21	23	44	47	21	30	27	34	39	52	19	18	18	32	37	53	37	49	43	40		

**Table A.5:** Annual abundance estimates for wolf at several spatial scales: the entire study area, by country, by management unit (carnivore management regions in Norway and "Rovdjursförvaltningsområden" in Sweden), and counties ("Län" in Sweden); see also Figure A.2. Only counties and management units that are within or that intersect the study area are included in the table. Estimates are based on model-estimated activity center locations. Credible intervals (95%) are shown in parentheses. Small deviations between the total estimate and the sum of abundance estimates from the constituent subregions may arise due to rounding. Note that the numbers reported here are predictions from a statistical model which always represents an oversimplification of reality and is based on available data (NGS and dead recoveries). As a consequence, especially at the local scale, the model-estimated number of wolves based on DNA sampling can deviate from the number of wolves inferred from ancillary observations (e.g., camera traps).

	2013/2014	2014/2015	2015/2016	2016/2017	2017/2018	2018/2019	2019/2020	2020/2021	2021/2022	2022/2023
TOTAL	394.7 (359-434)	421.8 (396-452)	363.7 (348-382)	408 (399-418)	410.4 (403-419)	373.3 (358-392)	433.4 (427-441)	449.4 (436-465)	501 (488-516)	462.7 (441-488)
NORWAY	70.3 (58-84)	72 (62-83)	70 (65-76)	82 (75-90)	94.1 (87-102)	79.7 (74-86)	85 (78-92)	88.3 (81-96)	92.8 (84-102)	68.6 (59-80)
Region 2	2.3 (1-5)	0.5 (0-2)	0.2 (0-1)	0.6 (0-2)	0.7 (0-3)	0.8 (0-2)	1 (0-3)	0.2 (0-1)	0.5 (0-2)	0.5 (0-2)
Region 3	5.3 (2-10)	2.9 (0-6)	0.5 (0-2)	1.1 (0-3)	1.7 (0-5)	0.7 (0-3)	2.7 (1-5)	1.1 (0-3)	1.6 (0-4)	2 (0-5)
Region 4	10.2 (6-15)	7 (4-11)	11.9 (9-15)	16.1 (13-20)	21.9 (18-26)	23.9 (20-28)	28 (24-33)	24.2 (20-29)	27.2 (23-32)	24.7 (21-29)
Region 5	47.5 (39-57)	58.7 (51-67)	56.8 (53-61)	63.1 (57-70)	68.5 (63-75)	52.5 (48-58)	51.3 (46-57)	61.9 (56-68)	61.5 (54-69)	39 (32-48)
Region 6	5.1 (1-10)	3 (0-7)	0.6 (0-3)	1.1 (0-4)	1.3 (0-4)	1.7 (1-4)	2.1 (1-5)	0.9 (0-3)	2 (1-4)	2.3 (0-6)
SWEDEN	324.4 (294-358)	349.8 (326-377)	293.7 (278-312)	326 (316-336)	316.3 (308-325)	293.6 (278-312)	348.4 (342-355)	361.2 (347-377)	408.2 (394-423)	394.1 (376-415)
Norra	27.8 (19-37)	23.8 (16-33)	20.3 (15-27)	17 (12-23)	7.7 (4-12)	15.7 (10-22)	16.9 (13-21)	16.5 (12-21)	10 (5-15)	23.2 (18-30)
Jämtland	19.9 (13-28)	18.2 (11-26)	15.8 (11-22)	14.4 (10-20)	6.9 (3-11)	12.9 (8-18)	15.8 (12-20)	14.7 (11-19)	8.4 (4-13)	15.5 (11-21)
Västerbotten	0.1 (0-1)	0.1 (0-1)	0 (0-1)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-1)
Västernorrland	7.8 (4-13)	5.5 (2-10)	4.4 (1-9)	2.6 (1-5)	0.7 (0-3)	2.8 (0-6)	1.1 (0-3)	1.7 (0-5)	1.7 (0-4)	7.6 (5-11)
Mellersta	273.8 (250-301)	310.5 (291-333)	264.1 (250-279)	298.9 (289-309)	289.1 (281-297)	266 (252-282)	310.8 (304-318)	313 (300-327)	351.6 (339-365)	310.5 (295-328)
Dalarna	67.5 (57-80)	72 (61-83)	74.8 (66-84)	79.8 (73-87)	62.9 (57-69)	55.9 (47-65)	69 (64-74)	67 (59-75)	75.5 (68-84)	56.7 (50-65)
Gävleborg	25.3 (18-34)	29.8 (23-38)	40.4 (34-47)	50.9 (45-57)	55.4 (51-60)	46.4 (39-54)	69.2 (65-74)	62.5 (57-68)	52.2 (46-58)	34.4 (30-40)
Örebro	32.5 (27-39)	51.1 (42-61)	31.9 (24-40)	39.5 (34-45)	35.1 (31-39)	38.4 (31-46)	49.3 (44-55)	50 (43-57)	60.7 (53-69)	50.7 (44-57)
Stockholm	3.8 (1-8)	3.6 (1-7)	4.2 (2-7)	2.5 (2-4)	4.3 (3-6)	2.2 (0-5)	3.5 (2-5)	7.3 (4-11)	6.1 (3-9)	3.8 (1-7)
Uppsala	4.7 (1-9)	4.3 (1-8)	4.2 (2-7)	5.5 (3-9)	8.2 (5-11)	6.7 (4-10)	4.9 (3-7)	7.7 (5-10)	12.6 (10-16)	7.2 (5-11)
Värmland	108.4 (96-122)	114.2 (102-128)	87.2 (79-96)	89.9 (82-98)	88.3 (82-94)	87.1 (78-96)	82.3 (77-88)	80.4 (71-90)	101.2 (92-111)	98.6 (91-108)
Västmanland	17.1 (12-23)	20.1 (15-26)	14.2 (10-19)	26 (23-30)	26.9 (23-31)	17.9 (13-24)	27.5 (23-32)	21.3 (16-27)	25.6 (20-32)	30.1 (25-36)
VästraGötaland	14.6 (9-21)	15.4 (11-21)	7.3 (4-11)	4.7 (2-8)	7.9 (5-12)	11.5 (7-16)	5.1 (2-8)	16.8 (12-22)	17.8 (13-23)	29 (25-34)
Södra	22.7 (14-33)	15.5 (9-23)	9.4 (4-15)	10.1 (7-14)	19.6 (17-23)	11.9 (7-17)	20.7 (18-24)	31.7 (26-38)	46.6 (41-53)	60.4 (55-67)
Blekinge	0.7 (0-3)	0.5 (0-2)	0.3 (0-2)	1.1 (1-2)	0.1 (0-1)	0.1 (0-1)	0.2 (0-1)	0.1 (0-1)	0.2 (0-1)	0.2 (0-1)
Halland	1.4 (0-4)	1 (0-3)	0.4 (0-2)	0.3 (0-2)	0.1 (0-1)	0.3 (0-2)	0.2 (0-1)	1.3 (0-3)	0.5 (0-2)	1.3 (0-3)
Jönköping	3.8 (1-8)	3.3 (1-7)	2.1 (1-5)	0.7 (0-3)	0.4 (0-2)	0.9 (0-3)	1.4 (0-3)	2.4 (1-5)	8.3 (5-12)	11.7 (9-15)
Kalmar	2.4 (0-6)	1.5 (0-4)	0.8 (0-3)	0.4 (0-2)	1 (0-2)	1.5 (1-3)	0.4 (0-2)	1.5 (1-3)	0.7 (0-3)	0.8 (0-3)
Kronoberg	2.9 (0-6)	2.6 (0-6)	1.4 (0-4)	1.3 (1-3)	0.4 (0-2)	0.6 (0-2)	0.6 (0-2)	1.5 (1-3)	1.6 (1-3)	3.5 (2-6)
Östergötland	3.6 (0-8)	2.4 (0-6)	1.4 (0-4)	2 (1-4)	3.4 (3-5)	3.6 (2-6)	9.9 (9-12)	7.8 (6-11)	11 (9-14)	9.1 (8-12)
Skåne	3.6 (1-7)	1.9 (0-5)	1.2 (0-3)	1.3 (1-3)	2 (1-3)	0.5 (0-2)	0.3 (0-2)	2.6 (2-4)	4.5 (4-6)	8.8 (8-11)
Södermanland	4.4 (1-8)	2.2 (0-5)	1.8 (0-5)	3 (2-5)	12.2 (11-14)	4.4 (2-7)	7.9 (6-10)	14.6 (11-19)	19.7 (16-24)	24.9 (22-28)

**Table A.6:** Annual population growth rate estimates for the wolf population in Scandinavia ("Total") and separately for Norway and Sweden. Estimates were derived using the posterior distributions of annual abundance estimates (Table 1). Credible intervals (95%) are shown in parentheses.

	2013-2014	2014-2015	2015-2016	2016-2017	2017-2018	2018-2019	2019-2020	2020-2021	2021-2022
Norway	1.03 (0.82-1.29)	0.98 (0.82-1.15)	1.17 (1.04-1.32)	1.15 (1.01-1.30)	0.85 (0.76-0.95)	1.07 (0.95-1.20)	1.04 (0.92-1.16)	1.05 (0.93-1.19)	0.74 (0.62-0.89)
Sweden	1.08 (0.95-1.22)	0.84 (0.76-0.92)	1.11 (1.04-1.18)	0.97 (0.93-1.01)	0.93 (0.87-0.99)	1.19 (1.12-1.26)	1.04 (0.99-1.09)	1.13 (1.07-1.19)	0.97 (0.91-1.03)
Total	1.07 (0.95-1.21)	0.86 (0.79-0.94)	1.12 (1.06-1.18)	1.01 (0.98-1.04)	0.91 (0.87-0.96)	1.16 (1.10-1.22)	1.04 (1.00-1.08)	1.12 (1.07-1.16)	0.92 (0.87-0.98)

**Table A.7:** Estimates of the demographic parameters obtained from the sex-specific wolf OPSCR models. Parameters represent transition rates from Oct 1 to Sep 30 in the following year. State 2 and 3 represents other and scent-marking individuals, respectively. Median estimates and 95% credible intervals (in parentheses) for per capita recruitment rate ( $\rho$ ), survival ( $\phi$ ), transition probability from other to scent-marking adult state ( $\psi$ ), mortality due to legal culling ( $h$ ) and mortality due to other causes ( $w$ ) are presented for males (M) and females (F).

	State	2013-2014		2014-2015		2015-2016		2016-2017		2017-2018	
		M	F	M	F	M	F	M	F	M	F
$\rho$	-	0.53 (0.30-0.80)	1.25 (0.90-1.65)	0.58 (0.44-0.74)	0.41 (0.27-0.56)	0.69 (0.57-0.81)	1.16 (0.97-1.35)	0.66 (0.60-0.72)	0.73 (0.66-0.81)	0.64 (0.55-0.74)	0.49 (0.41-0.60)
$\phi$	2	0.50 (0.39-0.62)	0.50 (0.36-0.64)	0.41 (0.32-0.51)	0.45 (0.34-0.56)	0.51 (0.41-0.60)	0.48 (0.37-0.60)	0.48 (0.39-0.56)	0.58 (0.49-0.67)	0.47 (0.38-0.56)	0.50 (0.41-0.59)
	3	0.66 (0.52-0.79)	0.67 (0.53-0.79)	0.65 (0.51-0.76)	0.68 (0.55-0.79)	0.68 (0.56-0.78)	0.65 (0.54-0.76)	0.55 (0.43-0.66)	0.57 (0.46-0.69)	0.57 (0.45-0.68)	0.61 (0.50-0.72)
$\psi$	-	0.19 (0.10-0.31)	0.40 (0.23-0.59)	0.41 (0.28-0.55)	0.40 (0.26-0.56)	0.34 (0.24-0.47)	0.55 (0.39-0.70)	0.42 (0.30-0.54)	0.45 (0.34-0.57)	0.34 (0.23-0.46)	0.28 (0.18-0.39)
$h$	2	0.07 (0.04-0.12)	0.12 (0.06-0.21)	0.18 (0.13-0.25)	0.21 (0.15-0.30)	0.12 (0.08-0.19)	0.07 (0.03-0.14)	0.15 (0.09-0.21)	0.12 (0.07-0.18)	0.25 (0.18-0.32)	0.15 (0.10-0.22)
	3	0.06 (0.02-0.14)	0.09 (0.04-0.16)	0.10 (0.04-0.19)	0.11 (0.05-0.19)	0.06 (0.02-0.13)	0.09 (0.04-0.17)	0.11 (0.06-0.20)	0.12 (0.06-0.20)	0.19 (0.11-0.29)	0.14 (0.07-0.23)
$w$	2	0.42 (0.31-0.53)	0.38 (0.24-0.52)	0.40 (0.31-0.50)	0.34 (0.23-0.45)	0.37 (0.28-0.46)	0.45 (0.33-0.56)	0.38 (0.30-0.46)	0.29 (0.22-0.38)	0.28 (0.20-0.37)	0.35 (0.26-0.44)
	3	0.27 (0.14-0.41)	0.25 (0.13-0.38)	0.25 (0.15-0.38)	0.21 (0.11-0.33)	0.25 (0.16-0.37)	0.25 (0.15-0.36)	0.34 (0.23-0.45)	0.30 (0.20-0.42)	0.23 (0.15-0.34)	0.25 (0.16-0.35)
	State	2018-2019		2019-2020		2020-2021		2021-2022			
		M	F	M	F	M	F	M	F		
$\rho$	-	0.82 (0.68-0.94)	0.87 (0.71-1.00)	0.76 (0.68-0.84)	0.63 (0.56-0.71)	0.74 (0.62-0.85)	0.86 (0.75-0.97)	0.63 (0.53-0.76)	0.65 (0.57-0.76)		
$\phi$	2	0.59 (0.50-0.69)	0.55 (0.45-0.65)	0.46 (0.38-0.54)	0.53 (0.44-0.61)	0.52 (0.43-0.60)	0.55 (0.45-0.64)	0.37 (0.30-0.45)	0.48 (0.39-0.57)		
	3	0.61 (0.49-0.73)	0.65 (0.53-0.76)	0.70 (0.59-0.80)	0.66 (0.55-0.76)	0.64 (0.53-0.74)	0.73 (0.62-0.82)	0.59 (0.48-0.69)	0.61 (0.50-0.72)		
$\psi$	-	0.36 (0.25-0.47)	0.48 (0.36-0.61)	0.39 (0.29-0.51)	0.39 (0.28-0.50)	0.36 (0.26-0.47)	0.36 (0.25-0.48)	0.50 (0.37-0.63)	0.48 (0.36-0.60)		
$h$	2	0.06 (0.03-0.11)	0.09 (0.05-0.15)	0.13 (0.08-0.18)	0.06 (0.03-0.10)	0.18 (0.13-0.25)	0.17 (0.11-0.24)	0.17 (0.12-0.23)	0.15 (0.10-0.21)		
	3	0.07 (0.03-0.15)	0.07 (0.02-0.14)	0.07 (0.02-0.14)	0.07 (0.03-0.15)	0.18 (0.11-0.28)	0.10 (0.04-0.17)	0.11 (0.06-0.19)	0.14 (0.07-0.22)		
$w$	2	0.34 (0.25-0.43)	0.35 (0.26-0.46)	0.41 (0.34-0.49)	0.42 (0.33-0.51)	0.30 (0.22-0.38)	0.28 (0.20-0.37)	0.45 (0.37-0.53)	0.37 (0.28-0.46)		
	3	0.31 (0.20-0.43)	0.27 (0.17-0.39)	0.23 (0.14-0.33)	0.26 (0.17-0.36)	0.17 (0.10-0.27)	0.17 (0.10-0.27)	0.29 (0.20-0.40)	0.25 (0.16-0.35)		



**Table A.8:** Estimates of the spatial and movement process parameters obtained from the sex-specific wolf OPSCR models.  $\beta_{dens}$  represents the effect of the number of known wolf packs on AC locations (Bischof et al., 2020). The scale parameter  $\sigma$  of the detection function is expressed in kilometers and estimated separately for other ( $\sigma_1$ ) and scent-marking ( $\sigma_2$ ) individuals.  $\lambda$  represents the mean of the exponential movement parameter, describing individual movement distances between years, estimated separately for other ( $\lambda_1$ ) and scent-marking ( $\lambda_2$ ) individuals. Credible intervals (95%) are shown in parentheses. Parameters that were not estimated every year are marked with \*.

	2013/2014		2014/2015		2015/2016		2016/2017		2017/2018	
	M	F	M	F	M	F	M	F	M	F
$\beta_{dens}^*$	0.46 (0.44-0.48)	0.48 (0.45-0.50)	0.46 (0.44-0.48)	0.48 (0.45-0.50)	0.46 (0.44-0.48)	0.48 (0.45-0.50)	0.46 (0.44-0.48)	0.48 (0.45-0.50)	0.46 (0.44-0.48)	0.48 (0.45-0.50)
$\sigma_1$	7.16 (6.46-8.06)	6.59 (5.73-7.67)	7.77 (6.93-8.80)	6.76 (6.02-7.67)	8.09 (7.47-8.79)	6.44 (5.79-7.20)	7.44 (7.02-7.92)	7.93 (7.48-8.43)	7.63 (7.28-8.01)	8.17 (7.76-8.61)
$\sigma_2$	7.51 (6.75-8.47)	6.27 (5.61-7.08)	8.50 (7.73-9.38)	8.35 (7.53-9.26)	8.36 (7.77-9.07)	7.33 (6.75-8.01)	7.92 (7.59-8.29)	7.92 (7.56-8.31)	7.88 (7.58-8.20)	7.35 (7.02-7.69)
$\lambda_1^*$	74.66 (69.99-80.04)	51.28 (47.79-55.73)	74.66 (69.99-80.04)	51.28 (47.79-55.73)	74.66 (69.99-80.04)	51.28 (47.79-55.73)	74.66 (69.99-80.04)	51.28 (47.79-55.73)	74.66 (69.99-80.04)	51.28 (47.79-55.73)
$\lambda_2^*$	6.95 (6.42-7.53)	4.68 (4.27-5.14)	6.95 (6.42-7.53)	4.68 (4.27-5.14)	6.95 (6.42-7.53)	4.68 (4.27-5.14)	6.95 (6.42-7.53)	4.68 (4.27-5.14)	6.95 (6.42-7.53)	4.68 (4.27-5.14)
	2018/2019		2019/2020		2020/2021		2021/2022		2022/2023	
	M	F	M	F	M	F	M	F	M	F
$\beta_{dens}^*$	0.46 (0.44-0.48)	0.48 (0.45-0.50)	0.46 (0.44-0.48)	0.48 (0.45-0.50)	0.46 (0.44-0.48)	0.48 (0.45-0.50)	0.46 (0.44-0.48)	0.48 (0.45-0.50)	0.46 (0.44-0.48)	0.48 (0.45-0.50)
$\sigma_1$	7.32 (6.78-7.93)	6.64 (6.08-7.29)	7.91 (7.61-8.24)	7.58 (7.27-7.91)	7.84 (7.38-8.33)	7.32 (6.81-7.89)	6.98 (6.56-7.45)	7.78 (7.31-8.32)	6.89 (6.38-7.49)	7.72 (7.16-8.34)
$\sigma_2$	8.45 (7.99-8.95)	8.22 (7.70-8.82)	8.03 (7.70-8.38)	7.76 (7.44-8.10)	7.65 (7.29-8.03)	7.08 (6.71-7.48)	7.63 (7.30-7.98)	7.77 (7.35-8.22)	7.47 (7.11-7.86)	6.94 (6.56-7.36)
$\lambda_1^*$	74.66 (69.99-80.04)	51.28 (47.79-55.73)	74.66 (69.99-80.04)	51.28 (47.79-55.73)	74.66 (69.99-80.04)	51.28 (47.79-55.73)	74.66 (69.99-80.04)	51.28 (47.79-55.73)	74.66 (69.99-80.04)	51.28 (47.79-55.73)
$\lambda_2^*$	6.95 (6.42-7.53)	4.68 (4.27-5.14)	6.95 (6.42-7.53)	4.68 (4.27-5.14)	6.95 (6.42-7.53)	4.68 (4.27-5.14)	6.95 (6.42-7.53)	4.68 (4.27-5.14)	6.95 (6.42-7.53)	4.68 (4.27-5.14)

**Table A.9:** Estimates of the detection process parameters for the structured sampling  $\beta_{1Structured}$  corresponds to the effect of previous detection of an individual on detection probability,  $\beta_{2Structured}$  to the effect of search-effort (track length) on detection probability, and  $\beta_{3Structured}$  to the effect of average snow cover during the monitoring period on detection probability ( $p_{0Structured}$ ). Coefficients are associated with scaled covariates. Credible intervals (95%) are shown in parentheses.

	2013/2014		2014/2015		2015/2016		2016/2017		2017/2018	
	M	F	M	F	M	F	M	F	M	F
$\beta_{1Structured}$	0.40 (-0.13-0.93)	0.62 (0.00-1.23)	-0.12 (-0.60-0.36)	-0.17 (-0.67-0.33)	-0.24 (-0.62-0.14)	-0.26 (-0.69-0.20)	0.09 (-0.20-0.40)	0.25 (-0.05-0.55)	-0.12 (-0.38-0.15)	0.34 (0.06-0.61)
$\beta_{2Structured}$	0.26 (0.16-0.36)	0.26 (0.16-0.37)	0.40 (0.30-0.50)	0.45 (0.34-0.56)	0.40 (0.30-0.49)	0.40 (0.26-0.53)	0.24 (0.18-0.31)	0.32 (0.25-0.39)	0.33 (0.27-0.39)	0.32 (0.25-0.38)
$\beta_{3Structured}$	-0.22 (-0.62- 0.18)	0.19 (-0.32-0.69)	-0.05 (-0.44- 0.35)	-0.27 (-0.70-0.15)	0.10 (-0.29- 0.50)	-0.47 (-0.96-0.00)	-0.26 (-0.50-0.02)	-0.21 (-0.50-0.07)	0.03 (-0.26- 0.34)	-0.01 (-0.38-0.37)
	2018/2019		2019/2020		2020/2021		2021/2022		2022/2023	
	M	F	M	F	M	F	M	F	M	F
$\beta_{1Structured}$	-0.37 (-0.85-0.06)	-0.01 (-0.41-0.39)	0.06 (-0.16-0.28)	0.03 (-0.21-0.26)	0.76 (0.43-1.09)	0.27 (-0.12-0.65)	0.36 (0.08-0.66)	-0.02 (-0.29-0.25)	0.40 (0.08-0.72)	-0.03 (-0.33-0.29)
$\beta_{2Structured}$	0.34 (0.25-0.42)	0.37 (0.28-0.46)	0.39 (0.33-0.46)	0.35 (0.28-0.42)	0.25 (0.19-0.31)	0.24 (0.17-0.32)	0.28 (0.22-0.34)	0.28 (0.22-0.34)	0.35 (0.27-0.44)	0.34 (0.28-0.41)
$\beta_{3Structured}$	-0.11 (-0.45- 0.22)	-0.02 (-0.43-0.40)	-0.04 (-0.25- 0.16)	0.05 (-0.18-0.29)	0.21 (0.01- 0.42)	0.25 (0.00-0.51)	0.25 (0.05- 0.46)	0.07 (-0.17-0.30)	0.03 (-0.16- 0.25)	0.15 (-0.09-0.39)

**Table A.10:** Estimates of the detection process parameters for the unstructured sampling  $\beta_{1Unstructured}$  corresponds to the effect of previous detection of an individual on baseline detection probability,  $\beta_{2Unstructured}$  to the effect of distance to the nearest roads on detection probability,  $\beta_{3Unstructured}$  to the effect of average snow cover during the monitoring period on detection probability, and  $\beta_{4Unstructured}$  to the effect of spatio-temporal heterogeneity in unstructured sampling derived using the observation data in Skandobs and Rovbase. Coefficients are associated with scaled covariates. Credible intervals (95%) are shown in parentheses.

	2013/2014		2014/2015		2015/2016		2016/2017		2017/2018	
	M	F	M	F	M	F	M	F	M	F
$\beta_{1Unstructured}$	1.24 ( 0.84-1.64)	0.26 (-0.30-0.79)	0.42 ( 0.03-0.81)	0.17 (-0.30-0.61)	0.42 ( 0.11-0.74)	-0.23 (-0.58-0.13)	0.11 (-0.13-0.34)	0.31 ( 0.09-0.54)	0.21 ( 0.02-0.41)	0.66 ( 0.45-0.87)
$\beta_{2Unstructured}$	-0.45 (-1.02-0.02)	-0.68 (-1.79- 0.05)	-1.05 (-2.24-0.18)	-0.92 (-2.42- 0.11)	-0.09 (-0.57- 0.32)	-0.47 (-1.23- 0.13)	-0.33 (-0.76- 0.01)	-0.48 (-1.03-0.04)	-1.25 (-2.07-0.59)	-0.76 (-1.34-0.27)
$\beta_{3Unstructured}$	0.33 ( 0.05- 0.60)	0.22 (-0.13- 0.57)	-0.10 (-0.39- 0.19)	-0.48 (-0.80-0.17)	0.12 (-0.17- 0.40)	0.03 (-0.32- 0.39)	-0.18 (-0.35-0.01)	-0.06 (-0.24- 0.13)	-0.27 (-0.43-0.11)	-0.28 (-0.48-0.09)
$\beta_{4Unstructured}$	1.15 (0.81-1.50)	1.30 (0.87-1.73)	0.80 (0.45-1.16)	0.69 (0.32-1.08)	0.73 (0.44-1.03)	0.79 (0.43-1.19)	1.03 (0.74-1.34)	1.13 (0.80-1.47)	1.19 (0.88-1.55)	1.59 (1.21-2.02)
	2018/2019		2019/2020		2020/2021		2021/2022		2022/2023	
	M	F	M	F	M	F	M	F	M	F
$\beta_{1Unstructured}$	0.02 (-0.30-0.36)	0.96 ( 0.55-1.38)	0.21 ( 0.02-0.40)	0.55 ( 0.36-0.73)	0.82 ( 0.55-1.11)	0.54 ( 0.21-0.87)	0.12 (-0.16-0.40)	0.43 ( 0.15-0.71)	0.74 ( 0.42-1.08)	0.39 ( 0.08-0.70)
$\beta_{2Unstructured}$	-0.20 (-0.79- 0.25)	0.46 (-0.20- 1.08)	-0.65 (-1.47-0.02)	-0.42 (-1.19- 0.24)	-0.84 (-1.73-0.17)	2.46 ( 0.75- 3.95)	0.16 (-0.34- 0.59)	-1.04 (-2.95- 0.39)	0.17 (-0.52- 0.87)	-0.34 (-1.60- 0.67)
$\beta_{3Unstructured}$	-0.15 (-0.41- 0.11)	-0.49 (-0.75-0.21)	-0.02 (-0.13- 0.10)	0.08 (-0.06- 0.21)	-0.19 (-0.36-0.03)	-0.31 (-0.51-0.12)	-0.12 (-0.27- 0.03)	-0.25 (-0.42-0.08)	-0.19 (-0.35-0.02)	-0.05 (-0.22- 0.11)
$\beta_{4Unstructured}$	1.09 (0.63-1.60)	1.73 (1.13-2.39)	1.57 (1.22-1.96)	1.79 (1.41-2.20)	1.22 (0.85-1.62)	1.47 (0.99-2.10)	1.92 (1.30-2.66)	1.40 (0.85-2.13)	1.09 (0.56-1.68)	1.61 (1.04-2.33)

**Table A.11:** Average proportion of individuals detected via non-invasive genetic sampling (NGS) in Scandinavia. Values were calculated as the number of individuals detected with NGS (Table A.3) divided by the total and sex-specific abundance estimates obtained from the OPSCR models (Table A.5). Credible intervals (95%) are shown in parentheses.

	2013/2014	2014/2015	2015/2016	2016/2017	2017/2018	2018/2019	2019/2020	2020/2021	2021/2022	2022/2023
M	0.54 (0.47-0.62)	0.56 (0.50-0.61)	0.80 (0.75-0.85)	0.90 (0.86-0.92)	0.93 (0.91-0.96)	0.79 (0.74-0.84)	0.95 (0.93-0.97)	0.84 (0.80-0.88)	0.87 (0.83-0.91)	0.85 (0.77-0.91)
F	0.62 (0.54-0.69)	0.57 (0.51-0.62)	0.73 (0.68-0.78)	0.89 (0.86-0.91)	0.91 (0.88-0.94)	0.76 (0.70-0.80)	0.94 (0.92-0.96)	0.86 (0.83-0.89)	0.88 (0.85-0.91)	0.87 (0.82-0.93)
Total	0.57 (0.52-0.62)	0.56 (0.52-0.60)	0.77 (0.73-0.80)	0.89 (0.87-0.91)	0.92 (0.90-0.94)	0.77 (0.74-0.81)	0.95 (0.93-0.96)	0.85 (0.82-0.88)	0.88 (0.85-0.90)	0.86 (0.81-0.90)